

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/14, 14/46	A2	(11) International Publication Number: WO 98/20034 (43) International Publication Date: 14 May 1998 (14.05.98)
(21) International Application Number: PCT/US97/20678 (22) International Filing Date: 5 November 1997 (05.11.97) (30) Priority Data: 08/745,995 7 November 1996 (07.11.96) US (71) Applicant (for all designated States except US): HESKA CORPORATION [US/US]; 1825 Sharp Point Drive, Fort Collins, CO 80525 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WISNEWSKI, Nancy [US/US]; 4219 Beaver Creek Drive, Fort Collins, CO 80526 (US). BRANDT, Kevin, S. [US/US]; 706 Walnut Street, Windsor, CO 80550 (US). SILVER, Gary, M. [US/US]; 1424 Fleetwood Court, Fort Collins, CO 80521 (US). MADDUX, Joely, D. [US/US]; 4533 Seaway Circle, Fort Collis, CO 80525 (US). (74) Agents: ROTHENBERGER, Scott, D. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NOVEL SERINE PROTEASE INHIBITOR NUCLEIC ACID MOLECULES, PROTEINS AND USES THEREOF (57) Abstract The present invention relates to flea serine protease inhibitor proteins; to flea serine protease inhibitor nucleic acid molecules, including those that encode such serine protease inhibitor proteins; to antibodies raised against such serine protease inhibitor proteins; and to compounds that inhibit flea serine protease inhibitor activity. The present invention also includes methods to obtain such proteins, nucleic acid molecules, antibodies, and inhibitory compounds. Also included in the present invention are therapeutic compositions comprising such proteins, nucleic acid molecules, antibodies and/or inhibitory compounds as well as the use of such therapeutic compositions to protect animals from hematophagous ectoparasite infestation.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NOVEL SERINE PROTEASE INHIBITOR NUCLEIC ACID MOLECULES, PROTEINS AND USES THEREOF

FIELD OF THE INVENTION

The present invention relates to flea serine protease inhibitor nucleic acid
5 molecules, proteins encoded by such nucleic acid molecules, antibodies raised against
such proteins, and inhibitors of such proteins. The present invention also includes
therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies,
and/or other inhibitors, as well as their use to protect an animal from flea infestation.

BACKGROUND OF THE INVENTION

10 Hematophagous ectoparasite infestation of animals is a health and economic
concern because hematophagous ectoparasites are known to cause and/or transmit a
variety of diseases. Hematophagous ectoparasites directly cause a variety of diseases,
including allergies, and also carry a variety of infectious agents including, but not
limited to, endoparasites (e.g., nematodes, cestodes, trematodes and protozoa), bacteria
15 and viruses. In particular, the bites of hematophagous ectoparasites are a problem for
animals maintained as pets because the infestation becomes a source of annoyance not
only for the pet but also for the pet owner who may find his or her home generally
contaminated with insects. As such, hematophagous ectoparasites are a problem not
only when they are on an animal but also when they are in the general environment of
20 the animal.

Bites from hematophagous ectoparasites are a particular problem because they
not only can lead to disease transmission but also can cause a hypersensitive response in
animals which is manifested as disease. For example, bites from fleas can cause an
allergic disease called flea allergic (or allergy) dermatitis (FAD). A hypersensitive
25 response in animals typically results in localized tissue inflammation and damage,
causing substantial discomfort to the animal.

The medical importance of hematophagous ectoparasite infestation has prompted
the development of reagents capable of controlling hematophagous ectoparasite
infestation. Commonly encountered methods to control hematophagous ectoparasite
30 infestation are generally focused on use of insecticides. While some of these products
are efficacious, most offer protection of a very limited duration at best. Furthermore,

many of the methods are often not successful in reducing hematophagous ectoparasite populations. In particular, insecticides have been used to prevent hematophagous ectoparasite infestation of animals by adding such insecticides to shampoos, powders, sprays, foggers, collars and liquid bath treatments (i.e., dips). Reduction of

5 hematophagous ectoparasite infestation on the pet has been unsuccessful for one or more of the following reasons: (1) failure of owner compliance (frequent administration is required); (2) behavioral or physiological intolerance of the pet to the pesticide product or means of administration; and (3) the emergence of hematophagous ectoparasite populations resistant to the prescribed dose of pesticide.

10 Prior investigators have described sequences of a few insect serine protease inhibitors: *Bombyx mori* nucleic acid and amino acid sequences have been disclosed by Narumi et al., *Eur. J. Biochem.*, 214:181-187, 1993; Takagi et al., *J. Biochem.*, 108:372-378, 1990; and amino acid sequence has been disclosed by Sasaki, *Eur. J Biochem*, 202:255-261, 1991. *Manduca sexta* nucleic acid and amino acid sequences have been

15 disclosed by Kanost et al., *J. Biol. Chem*, 264:965-972, 1989; U.S. Patent No. 5,436,392, to Thomas et al., issued July 25, 2085, 1990; U.S. Patent No. 5,196,304, to Kanost et al., issued March 23, 1993; Jiang et al., *J. Biol. Chem.*, 269:55-58, 1994; and *Manduca sexta* peptide sequences have been disclosed by Fox et al., *Peptides*, 12:937-944, 1991. *Locusta migratoria* peptide sequences have been disclosed by Kellenberger et al., *J.*

20 *Biol. Chem*, 270:25514-25519, 1995. *Rhodnius prolixus* peptide sequences have been disclosed by Van De Loch, *EMBO*, 14:5149-5157, 1995. *Lymantria dispar* peptide sequences have been disclosed by Valaitis, *Insect Biochem Molec Biol*, 25:139-149, 1995. *Lucilia cuprina* nucleic acid and amino acid sequences have been disclosed by Casu et al., *Insect Molecular Biology*, 3:159-170, 1994. Identification of a serine

25 protease inhibitor of the present invention is unexpected because the most identical amino acid or nucleic acid sequence identified by previous investigators could not be used to identify a flea serine protease inhibitor of the present invention.

In summary, there remains a need to develop a reagent and a method to protect animals from hematophagous ectoparasite infestation.

SUMMARY OF THE INVENTION

The present invention relates to a novel product and process for protection of animals from hematophagous ectoparasite infestation. According to the present invention there are provided flea serine protease inhibitor proteins and mimetopes thereof; flea nucleic acid molecules, including those that encode such proteins; antibodies raised against such serine protease inhibitor proteins (i.e., anti-flea serine protease inhibitor antibodies); and other compounds that inhibit flea serine protease inhibitor activity (i.e, inhibitory compounds or inhibitors).

The present invention also includes methods to obtain such proteins, mimetopes, nucleic acid molecules, antibodies and inhibitory compounds. Also included in the present invention are therapeutic compositions comprising such proteins, mimetopes, nucleic acid molecules, antibodies, and/or inhibitory compounds, as well as use of such therapeutic compositions to protect animals from hematophagous ectoparasite infestation.

Identification of a serine protease inhibitor protein of the present invention is unexpected because the most identical amino acid or nucleic acid sequence identified by previous investigators could not be used to identify a flea serine protease inhibitor protein of the present invention. In addition, identification of a flea serine protease inhibitor protein of the present invention is unexpected because a protein fraction from flea prepupal larvae that was obtained by monitoring for carboxylesterase activity surprisingly also contained flea serine protease inhibitor molecular epitopes of the present invention.

One embodiment of the present invention is an isolated flea serine protease nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene, including, but not limited to, nucleic acid molecules that hybridize under stringent conditions with a nucleic acid molecule having at least one of the following nucleic acid sequences: SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID

NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90. Particularly preferred flea serine protease inhibitor nucleic acid molecules include nucleic acid sequences SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, and/or nucleic acid sequences encoding proteins having amino acid sequences SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90, as well as allelic variants of any of the listed nucleic acid sequences or complements of any of the listed nucleic acid sequences.

25 The present invention also includes an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a nucleic acid sequence encoding a protein comprising an amino acid sequence including SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID

NO:89, SEQ ID NO:90, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO: 97, and SEQ ID NO:98.

The present invention also relates to recombinant molecules, recombinant viruses and recombinant cells that include flea serine protease inhibitor nucleic acid molecules
5 of the present invention. Also included are methods to produce such nucleic acid molecules, recombinant molecules, recombinant viruses and recombinant cells.

Another embodiment of the present invention includes an isolated flea serine protease inhibitor protein. A preferred flea serine protease inhibitor protein is capable of eliciting an immune response when administered to an animal and/or of having serine
10 protease inhibitor activity. A preferred flea serine protease inhibitor protein is encoded by a nucleic acid molecule that hybridizes under stringent hybridization conditions to a nucleic acid sequence including SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:15, SEQ ID NO:21, SEQ ID NO:27, and SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID
15 NO:68 and SEQ ID NO:71. Particularly preferred flea serine protease inhibitor proteins include at least one of the following amino acid sequences: SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID
20 NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO: 97, and SEQ ID NO:98.

Yet another embodiment of the present invention is a therapeutic composition that is capable of reducing hematophagous ectoparasite infestation. Such a therapeutic
25 composition includes one or more of the following protective compounds: an isolated flea serine protease inhibitor protein or a mimetope thereof; an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene; an isolated antibody that selectively binds to a flea *Ctenocephalides felis* serine protease inhibitor protein; and an inhibitor of
30 serine protease inhibitor protein activity identified by its ability to inhibit flea serine protease inhibitor activity, such as, but not limited to, a substrate analog of a flea serine

protease inhibitor protein. A preferred therapeutic composition of the present invention also includes an excipient, an adjuvant and/or a carrier. Also included in the present invention is a method to reduce flea infestation. The method includes the step of administering to the animal a therapeutic composition of the present invention.

5 The present invention also includes an inhibitor of serine protease inhibitor protein activity identified by its ability to inhibit the activity of a flea serine protease inhibitor protein. An example of such an inhibitor is a substrate analog of a flea serine protease inhibitor protein. Also included in the present invention are mimetopes of flea serine protease inhibitor proteins of the present invention identified by their ability to
10 inhibit flea serine protease activity.

 Yet another embodiment of the present invention is a method to identify a compound capable of inhibiting flea serine protease inhibitor activity. The method includes the steps of: (a) contacting an isolated flea serine protease inhibitor protein with a putative inhibitory compound under conditions in which, in the absence of the
15 compound, the protein has serine protease inhibitor activity; and (b) determining if the putative inhibitory compound inhibits the activity. Also included in the present invention is a test kit to identify a compound capable of inhibiting flea serine protease inhibitor activity. Such a kit includes an isolated flea serine protease inhibitor protein having serine protease inhibitor activity and a means for determining the extent of
20 inhibition of the activity in the presence of a putative inhibitory compound.

 Yet another embodiment of the present invention is a method to produce a flea serine protease inhibitor protein, the method comprising culturing a cell transformed with a nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene.

25 BRIEF DESCRIPTION OF THE FIGURES

 Fig. 1 depicts proteins from tissue extracts that bind to a polyclonal antiserum made against a serine protease inhibitor protein.

DETAILED DESCRIPTION OF THE INVENTION

 The present invention provides for isolated flea serine protease inhibitor (SPI)
30 proteins, isolated flea serine protease inhibitor nucleic acid molecules, antibodies directed against flea serine protease inhibitor proteins and other inhibitors of flea serine

protease inhibitor activity. As used herein, the terms isolated flea serine protease inhibitor proteins and isolated flea serine protease inhibitor nucleic acid molecules refers to serine protease inhibitor proteins and serine protease inhibitor nucleic acid molecules derived from fleas and, as such, can be obtained from their natural source or can be
5 produced using, for example, recombinant nucleic acid technology or chemical synthesis. A SPI protein can have the ability to inhibit the proteolytic activity of a serine protease protein. A protein denoted as a SPI protein can also possess cysteine protease activity, in addition to serine protease activity. Also included in the present invention is the use of these proteins, nucleic acid molecules, antibodies and other inhibitors as
10 therapeutic compositions to protect animals from hematophagous ectoparasite infestation as well as in other applications, such as those disclosed below.

Flea serine protease inhibitor proteins and nucleic acid molecules of the present invention have utility because they represent novel targets for anti-hematophagous ectoparasite vaccines and drugs. The products and processes of the present invention are
15 advantageous because they enable the inhibition of hematophagous ectoparasite serine protease activity necessary for hematophagous ectoparasite survival or the inhibition of serine protease inhibitors, thereby deregulating serine protease activity, leading to uncontrolled proteolysis of an hematophagous ectoparasite.

One embodiment of the present invention is an isolated protein comprising a flea
20 SPI protein. It is to be noted that the term "a" or "an" entity refers to one or more of that entity; for example, a protein refers to one or more proteins or at least one protein. As such, the terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. Furthermore, a compound "selected from the
25 group consisting of" refers to one or more of the compounds in the list that follows, including mixtures (i.e., combinations) of two or more of the compounds. According to the present invention, an isolated, or biologically pure, protein, is a protein that has been removed from its natural milieu. As such, "isolated" and "biologically pure" do not necessarily reflect the extent to which the protein has been purified. An isolated protein
30 of the present invention can be obtained from its natural source, can be produced using recombinant DNA technology or can be produced by chemical synthesis.

As used herein, an isolated flea SPI protein can be a full-length protein or any homolog of such a protein. An isolated protein of the present invention, including a homolog, can be identified in a straight-forward manner by the protein's ability to elicit an immune response against flea SPI proteins and/or ability to inhibit, or reduce, serine protease activity. Examples of serine protease inhibitor homologs include SPI proteins in which amino acids have been deleted (e.g., a truncated version of the protein, such as a peptide), inserted, inverted, substituted and/or derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation and/or addition of glycerophosphatidyl inositol) such that the homolog includes at least one epitope capable of eliciting an immune response against a flea protein or has at least some serine protease inhibitor activity. For example, when the homolog is administered to an animal as an immunogen, using techniques known to those skilled in the art, the animal will produce an immune response against at least one epitope of a natural flea SPI protein. The ability of a protein to effect an immune response, can be measured using techniques known to those skilled in the art. Techniques to measure serine protease inhibitor activity are also known to those skilled in the art; see, for example, Jiang et al., 1995, *Insect Biochem. Molec. Biol.* 25, 1093-1100.

Flea SPI protein homologs can be the result of natural allelic variation or natural mutation. SPI protein homologs of the present invention can also be produced using techniques known in the art including, but not limited to, direct modifications to the protein or modifications to the gene encoding the protein using, for example, classic or recombinant nucleic acid techniques to effect random or targeted mutagenesis.

Isolated SPI proteins of the present invention have the further characteristic of being encoded by nucleic acid molecules that hybridize under stringent hybridization conditions to a gene encoding a *Ctenocephalides felis* SPI protein (i.e., a *C. felis* SPI gene). As used herein, stringent hybridization conditions refer to standard hybridization conditions under which nucleic acid molecules, including oligonucleotides, are used to identify similar nucleic acid molecules. Such standard conditions are disclosed, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press, 1989; Sambrook et al., *ibid.*, is incorporated by reference herein in its entirety. Stringent hybridization conditions typically permit isolation of nucleic acid

molecules having at least about 70% nucleic acid sequence identity with the nucleic acid molecule being used to probe in the hybridization reaction. Formulae to calculate the appropriate hybridization and wash conditions to achieve hybridization permitting 30% or less mismatch of nucleotides are disclosed, for example, in Meinkoth et al., 1984, 5 *Anal. Biochem.* 138, 267-284; Meinkoth et al., *ibid.*, is incorporated by reference herein in its entirety.

As used herein, a *C. felis* SPI gene includes all nucleic acid sequences related to a natural *C. felis* SPI gene such as regulatory regions that control production of the *C. felis* SPI protein encoded by that gene (such as, but not limited to, transcription, translation or 10 post-translation control regions) as well as the coding region itself. In one embodiment, a *C. felis* SPI gene of the present invention includes the nucleic acid sequence SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:3, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID 15 NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69 and/or SEQ ID NO:71. Nucleic acid sequence SEQ ID NO:1 represents the deduced sequence of the coding strand of a complementary DNA (cDNA) nucleic acid molecule denoted herein 20 as nfSPI1₁₅₈₄, the production of which is disclosed in the Examples. The complement of SEQ ID NO:1 (represented herein by SEQ ID NO:3) refers to the nucleic acid sequence of the strand complementary to the strand having SEQ ID NO:1, which can easily be determined by those skilled in the art. Likewise, a nucleic acid sequence complement of any nucleic acid sequence of the present invention refers to the nucleic acid sequence of 25 the nucleic acid strand that is complementary to (i.e., can form a complete double helix with) the strand for which the sequence is cited.

Nucleic acid sequence SEQ ID NO:7 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI2₁₃₅₈, the production of which is disclosed in the Examples. The complement of SEQ ID NO:7 is 30 represented herein by SEQ ID NO:9.

Nucleic acid sequence SEQ ID NO:13 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI3₁₈₃₈, the production of which is disclosed in the Examples. The complement of SEQ ID NO:13 is represented herein by SEQ ID NO:15.

5 Nucleic acid sequence SEQ ID NO:19 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI4₁₄₁₄, the production of which is disclosed in the Examples. The complement of SEQ ID NO:19 is represented herein by SEQ ID NO:21.

10 Nucleic acid sequence SEQ ID NO:25 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI5₁₄₉₂, the production of which is disclosed in the Examples. The complement of SEQ ID NO:25 is represented herein by SEQ ID NO:27.

15 Nucleic acid sequence SEQ ID NO:31 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI6₁₄₅₄, the production of which is disclosed in the Examples. The complement of SEQ ID NO:31 is represented herein by SEQ ID NO:33.

20 Nucleic acid sequence SEQ ID NO:45 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI7₅₄₉, the production of which is disclosed in the Examples. The complement of SEQ ID NO:45 is represented herein by SEQ ID NO:47.

Nucleic acid sequence SEQ ID NO:48 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI8₅₄₉, the production of which is disclosed in the Examples. The complement of SEQ ID NO:48 is represented herein by SEQ ID NO:50.

25 Nucleic acid sequence SEQ ID NO:51 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI9₅₈₁, the production of which is disclosed in the Examples. The complement of SEQ ID NO:51 is represented herein by SEQ ID NO:53.

30 Nucleic acid sequence SEQ ID NO:54 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI10₆₅₄, the

production of which is disclosed in the Examples. The complement of SEQ ID NO:54 is represented herein by SEQ ID NO:56.

Nucleic acid sequence SEQ ID NO:57 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI1₆₇₀, the
5 production of which is disclosed in the Examples. The complement of SEQ ID NO:57 is represented herein by SEQ ID NO:59.

Nucleic acid sequence SEQ ID NO:60 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI2₇₀₆, the
production of which is disclosed in the Examples. The complement of SEQ ID NO:60 is
10 represented herein by SEQ ID NO:62.

Nucleic acid sequence SEQ ID NO:63 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI3₆₂₃, the
production of which is disclosed in the Examples. The complement of SEQ ID NO:63 is represented herein by SEQ ID NO:65.

15 Nucleic acid sequence SEQ ID NO:66 represents the deduced sequence of the coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI4₇₃₁, the
production of which is disclosed in the Examples. The complement of SEQ ID NO:66 is represented herein by SEQ ID NO:68.

Nucleic acid sequence SEQ ID NO:69 represents the deduced sequence of the
20 coding strand of a cDNA nucleic acid molecule denoted herein as nfSPI5₆₈₅, the
production of which is disclosed in the Examples. The complement of SEQ ID NO:69 is represented herein by SEQ ID NO:71.

It should be noted that since nucleic acid sequencing technology is not entirely error-free, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:13, SEQ ID NO:19, SEQ ID
25 NO:25, SEQ ID NO:31, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:57, SEQ ID NO:60, SEQ ID NO:63, SEQ ID NO:66 and SEQ ID NO:69, and complements thereof (as well as other nucleic acid and protein sequences presented herein), at best, represent apparent nucleic acid sequences of certain nucleic acid molecules encoding *C. felis* SPI proteins of the present invention.

30 In another embodiment, a *C. felis* SPI gene can be an allelic variant that includes a similar but not identical sequence to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4,

SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33. SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90. An allelic variant of a *C. felis* SPI gene is a gene that occurs at essentially the same locus (or loci) in the genome as the gene including SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33. SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90, but which, due to natural variations caused by, for example, mutation or recombination, has a similar but not identical sequence. Allelic variants typically encode proteins having similar activity to that of the protein encoded by the gene to which they are being compared. Allelic variants can also comprise alterations in the 5' or 3' untranslated regions of the gene (e.g., in regulatory control regions). Allelic variants are well known to those skilled in the art and would be expected to be found within a given flea since the genome is diploid and/or among a group of two or more fleas.

The minimal size of a SPI protein homolog of the present invention is a size sufficient to be encoded by a nucleic acid molecule capable of forming a stable hybrid

(i.e., hybridize under stringent hybridization conditions) with the complementary sequence of a nucleic acid molecule encoding the corresponding natural protein. As such, the size of the nucleic acid molecule encoding such a protein homolog is dependent on nucleic acid composition and percent homology between the nucleic acid molecule and complementary sequence. It should also be noted that the extent of homology required to form a stable hybrid can vary depending on whether the homologous sequences are interspersed throughout the nucleic acid molecules or are clustered (i.e., localized) in distinct regions on the nucleic acid molecules. The minimal size of such nucleic acid molecules is typically at least about 12 to about 15 nucleotides in length if the nucleic acid molecules are GC-rich and at least about 15 to about 17 bases in length if they are AT-rich. As such, the minimal size of a nucleic acid molecule used to encode a SPI protein homolog of the present invention is from about 12 to about 18 nucleotides in length. Thus, the minimal size of a SPI protein homolog of the present invention is from about 4 to about 6 amino acids in length. There is no limit, other than a practical limit, on the maximal size of such a nucleic acid molecule in that the nucleic acid molecule can include a portion of a gene, an entire gene, multiple genes, or portions thereof. The preferred size of a protein encoded by a nucleic acid molecule of the present invention depends on whether a full-length, fusion, multivalent, or functional portion of such a protein is desired.

Suitable fleas from which to isolate SPI proteins of the present invention (including isolation of the natural protein or production of the protein by recombinant or synthetic techniques) include *Ctenocephalides*, *Ceratophyllus*, *Diamanus*, *Echidnophaga*, *Nosopsyllus*, *Pulex*, *Tunga*, *Oropsylla*, *Orchopeus* and *Xenopsylla*. More preferred fleas from which to isolate SPI proteins include *Ctenocephalides felis*, *Ctenocephalides canis*, *Ceratophyllus pulicidae*, *Pulex irritans*, *Oropsylla (Thrassis) bacchi*, *Oropsylla (Diamanus) montana*, *Orchopeus howardi*, *Xenopsylla cheopis* and *Pulex simulans*, with *C. felis* being even more preferred.

Suitable flea tissues from which to isolate a SPI protein of the present invention includes tissues from unfed fleas or tissue from fleas that recently consumed a blood meal (i.e., blood-fed fleas). Such flea tissues are referred to herein as, respectively, unfed flea tissues and fed flea tissues. Preferred flea tissues from which to obtain a SPI

protein of the present invention includes unfed or fed pre-pupal larval, 1st instar larval, 2nd instar larval, 3rd instar larval, and/or adult flea tissues. More preferred flea tissue includes prepupal larval tissue. A SPI of the present invention is also preferably obtained from hemolymph.

5 A preferred flea SPI protein of the present invention is a compound that when administered to an animal in an effective manner, is capable of protecting that animal from a hematophagous ectoparasite infestation. In accordance with the present invention, the ability of a SPI protein of the present invention to protect an animal from a hematophagous ectoparasite infestation refers to the ability of that protein to, for
10 example, treat, ameliorate and/or prevent infestation caused by a hematophagous ectoparasite. In particular, the phrase "to protect an animal from hematophagous ectoparasite infestation" refers to reducing the potential for hematophagous ectoparasite population expansion on and around the animal (i.e., reducing the hematophagous ectoparasite burden). Preferably, the hematophagous ectoparasite population size is
15 decreased, optimally to an extent that the animal is no longer bothered by hematophagous ectoparasites. A host animal, as used herein, is an animal from which hematophagous ectoparasites can feed by attaching to and feeding through the skin of the animal. Hematophagous ectoparasites, and other ectoparasites, can live on a host animal for an extended period of time or can attach temporarily to an animal in order to
20 feed. At any given time, a certain percentage of a hematophagous ectoparasite population can be on a host animal whereas the remainder can be in the environment of the animal. Such an environment can include not only adult hematophagous ectoparasites, but also hematophagous ectoparasite eggs and/or hematophagous ectoparasite larvae. The environment can be of any size such that hematophagous
25 ectoparasite in the environment are able to jump onto and off of a host animal. For example, the environment of an animal can include plants, such as crops, from which hematophagous ectoparasites infest an animal. As such, it is desirable not only to reduce the hematophagous ectoparasite burden on an animal per se, but also to reduce the hematophagous ectoparasite burden in the environment of the animal. In one
30 embodiment, a SPI protein of the present invention can elicit an immune response

(including a humoral and/or cellular immune response) against a hematophagous ectoparasite.

Suitable hematophagous ectoparasites to target include any hematophagous ectoparasite that is essentially incapable of infesting an animal administered a SPI protein of the present invention. As such, a hematophagous ectoparasite to target includes any hematophagous ectoparasite that produces a protein having one or more epitopes that can be targeted by a humoral and/or cellular immune response against a SPI protein of the present invention, that can be targeted by a compound that otherwise inhibits SPI activity, and/or that can be targeted by a SPI protein (e.g., a peptide) or mimetope of a SPI protein of the present invention in such a manner as to inhibit serine protease activity, thereby resulting in the decreased ability of the hematophagous ectoparasite to infest an animal. Preferred hematophagous ectoparasite to target include insects and acarines. A SPI protein of the present invention preferably protects an animal from infestation by hematophagous ectoparasites including, but are not limited to, agricultural pests, stored product pests, forest pests, structural pests or animal health pests. Suitable agricultural pests of the present invention include, but are not limited to, Colorado potato beetles, corn earworms, fleahoppers, weevils, pink boll worms, cotton aphids, beet armyworms, lygus bugs, hessian flies, sod webworms, whites grubs, diamond back moths, white flies, planthoppers, leafhoppers, mealy bugs, mormon crickets and mole crickets. Suitable stored product pests of the present invention include, but are not limited to, dermestids, anobeids, saw toothed grain beetles, indian mealmoths, flour beetles, long-horn wood boring beetles and metallic wood boring beetles. Suitable forest pests of the present invention include, but are not limited to, southern pine bark beetles, gypsy moths, elm beetles, ambrosia beetles, bag worms, tent worms and tussock moths. Suitable structural pests of the present invention include, but are not limited to, bess beetles, termites, fire ants, carpenter ants, wasps, hornets, cockroaches, silverfish, *Musca domestica* and *Musca autumnalis*. Suitable animal health pests of the present invention include, but are not limited to, fleas, ticks, mosquitoes, black flies, lice, true bugs, sand flies, *Psychodidae*, tsetse flies, sheep blow flies, cattle grub, mites, horn flies, heel flies, deer flies, *Culicoides* and warble flies. A SPI protein of the present invention more preferably protects an animal from infestation by

hematophagous ectoparasites including fleas, midges, mosquitos, sand flies, black flies, horse flies, snipe flies, louse flies, horn flies, deer flies, tsetse flies, buffalo flies, blow flies, stable flies, myiasis-causing flies, biting gnats, lice, mites, bee, wasps, ants, true bugs and ticks, even more preferably fleas and ticks, and even more preferably fleas.

- 5 Preferred fleas from which to protect an animal from flea infestation include those disclosed herein for the isolation of a SPI of the present invention.

The present invention also includes mimetopes of SPI proteins of the present invention. As used herein, a mimetope of a SPI protein of the present invention refers to any compound that is able to mimic the activity of such a SPI protein (e.g., ability to
10 elicit an immune response against a SPI protein of the present invention and/or ability to inhibit serine protease activity), often because the mimetope has a structure that mimics the SPI protein. It is to be noted, however, that the mimetope need not have a structure similar to an SPI protein as long as the mimetope functionally mimics the protein. Mimetopes can be, but are not limited to: peptides that have been modified to decrease
15 their susceptibility to degradation; anti-idiotypic and/or catalytic antibodies, or fragments thereof; non-proteinaceous immunogenic portions of an isolated protein (e.g., carbohydrate structures); synthetic or natural organic or inorganic molecules, including nucleic acids; and/or any other peptidomimetic compounds. Mimetopes of the present invention can be designed using computer-generated structures of SPI proteins of the
20 present invention. Mimetopes can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides or other organic molecules, and screening such samples by affinity chromatography techniques using the corresponding binding partner, (e.g., a flea serine protease or anti-flea serine protease inhibitor antibody). A preferred mimetope is a peptidomimetic compound that is structurally and/or
25 functionally similar to a SPI protein of the present invention, particularly to the active site of the SPI protein.

One embodiment of a flea SPI protein of the present invention is a fusion protein that includes a flea SPI protein-containing domain attached to one or more fusion segments. Suitable fusion segments for use with the present invention include, but are
30 not limited to, segments that can: enhance a protein's stability; act as an immunopotentiator to enhance an immune response against a SPI protein; and/or assist

-17-

purification of a SPI protein (e.g., by affinity chromatography). A suitable fusion segment can be a domain of any size that has the desired function (e.g., imparts increased stability, imparts increased immunogenicity to a protein, and/or simplifies purification of a protein). Fusion segments can be joined to amino and/or carboxyl

5 termini of the SPI-containing domain of the protein and can be susceptible to cleavage in order to enable straight-forward recovery of a SPI protein. Fusion proteins are preferably produced by culturing a recombinant cell transformed with a fusion nucleic acid molecule that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of a SPI-containing domain. Preferred fusion

10 segments include a metal binding domain (e.g., a poly-histidine segment); an immunoglobulin binding domain (e.g., Protein A; Protein G; T cell; B cell; Fc receptor or complement protein antibody-binding domains); a sugar binding domain (e.g., a maltose binding domain); and/or a "tag" domain (e.g., at least a portion of β -galactosidase, a strep tag peptide, other domains that can be purified using compounds

15 that bind to the domain, such as monoclonal antibodies). More preferred fusion segments include metal binding domains, such as a poly-histidine segment; a maltose binding domain; a strep tag peptide, such as that available from Biometra in Tampa, FL; and an S10 peptide. Examples of particularly preferred fusion proteins of the present invention include PHis-PfSPI2₃₇₆, PHis-PfSPI3₃₉₀, PHis-PfSPI4₃₇₆, PHis-PfSPI6₃₇₆, PHis-

20 PfSPIC4:V7, PHis-PfSPIC4:V8, PHis-PfSPIC4:V9, PHis-PfSPIC4:V10, PHis-PfSPIC4:V12, PHis-PfSPIC4:V13 and PHis-PfSPIC4:V15, production of which are disclosed herein.

In another embodiment, a flea SPI protein of the present invention also includes at least one additional protein segment that is capable of protecting an animal from

25 hematophagous ectoparasite infestations. Such a multivalent protective protein can be produced by culturing a cell transformed with a nucleic acid molecule comprising two or more nucleic acid domains joined together in such a manner that the resulting nucleic acid molecule is expressed as a multivalent protective compound containing at least two protective compounds, or portions thereof, capable of protecting an animal from

30 hematophagous ectoparasite infestation by, for example, targeting two different flea proteins.

Examples of multivalent protective compounds include, but are not limited to, a SPI protein of the present invention attached to one or more compounds protective against one or more flea compounds. Preferred second compounds are proteinaceous compounds that effect active immunization (e.g., antigen vaccines), passive immunization (e.g., antibodies), or that otherwise inhibit a hematophagous ectoparasite activity that when inhibited can reduce hematophagous ectoparasite burden on and around an animal. Examples of second compounds include a compound that inhibits binding between a flea protein and its ligand (e.g., a compound that inhibits flea ATPase activity or a compound that inhibits binding of a peptide or steroid hormone to its receptor), a compound that inhibits hormone (including peptide or steroid hormone) synthesis, a compound that inhibits vitellogenesis (including production of vitellin and/or transport and maturation thereof into a major egg yolk protein), a compound that inhibits fat body function, a compound that inhibits muscle action, a compound that inhibits the nervous system, a compound that inhibits the immune system and/or a compound that inhibits flea feeding. Particular examples of second compounds include, but are not limited to, serine proteases, cysteine proteases, aminopeptidases, calreticulins and esterases, as well as antibodies and inhibitors of such proteins. In one embodiment, a flea SPI protein of the present invention is attached to one or more additional compounds protective against hematophagous ectoparasite infestation. In another embodiment, one or more protective compounds, such as those listed above, can be included in a multivalent vaccine comprising a flea SPI protein of the present invention and one or more other protective molecules as separate compounds.

A preferred flea SPI protein of the present invention is encoded by a nucleic acid molecule that hybridizes under stringent hybridization conditions with at least one of the following nucleic acid molecules: nfSPI1₁₅₈₄, nfSPI1₁₁₉₁, nfSPI1₃₇₆, nfSPI2₁₃₅₈, nfSPI2₁₁₉₇, nfSPI2₃₇₆, nfSPI3₁₈₃₈, nfSPI3₁₂₆₀, nfSPI3₃₉₁, nfSPI4₁₄₁₄, nfSPI4₁₁₇₉, nfSPI4₃₇₆, nfSPI5₁₄₉₂, nfSPI5₁₁₉₄, nfSPI5₃₇₆, nfSPI6₁₄₅₄, nfSPI6₁₁₉₁, nfSPI6₃₇₆, nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI11₆₇₀, nfSPI12₇₀₆, nfSPI13₆₂₃, nfSPI14₇₃₁, nfSPI15₆₈₅, nfSPI3₁₂₂₂, nfSPI6₁₁₅₅, nfSPI2₁₀₆₅ and nfSPI4₁₀₇₀. A further preferred isolated protein is encoded by a nucleic acid molecule that hybridizes under stringent hybridization conditions with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:3,

SEQ ID NO:9, SEQ ID NO:15, SEQ ID NO:21, SEQ ID NO:27, and SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68 and SEQ ID NO:71.

Translation of SEQ ID NO:1 suggests that nucleic acid molecule nfSPII₁₅₈₄ encodes a full-length flea protein of about 397 amino acids, referred to herein as PfSPII₃₉₇, represented by SEQ ID NO:2, assuming an open reading frame having an initiation (start) codon spanning from about nucleotide 136 through about nucleotide 138 of SEQ ID NO:1 and a termination (stop) codon spanning from about nucleotide 1327 through about nucleotide 1329 of SEQ ID NO:1. The coding region encoding PfSPII₃₉₇ is represented by nucleic acid molecule nfSPII₁₁₉₁, having a coding strand with the nucleic acid sequence represented by SEQ ID NO:4 and a complementary strand with the nucleic acid sequence represented by SEQ ID NO:5. The deduced amino acid sequence SEQ ID NO:2 suggests a protein having a molecular weight of about 44.4 kilodaltons (kD) and an estimated pI of about 4.97. Analysis of SEQ ID NO:2 suggests the presence of a signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 21. The proposed mature protein, denoted herein as PfSPII₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:6. The amino acid sequence of flea PfSPII₃₇₆ (i.e. SEQ ID NO:6) predicts that PfSPII₃₇₆ has an estimated molecular weight of about 42.1 kD, an estimated pI of about 4.90, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

Comparison of amino acid sequence SEQ ID NO:2 (i.e., the amino acid sequence of PfSPII₃₉₇) with amino acid sequences reported in GenBank indicates that SEQ ID NO:2 showed the most homology, i.e., about 36% identity, with GenBank accession number 1378131, a serpin protein from *Manduca sexta*.

Translation of SEQ ID NO:7 suggests that nucleic acid molecule nfSPI2₁₃₅₈ encodes a non-full-length flea SPI protein of about 399 amino acids, referred to herein as PfSPI2₃₉₉, represented by SEQ ID NO:8, assuming an open reading frame having a first in-frame codon spanning from about nucleotide 2 through about nucleotide 4 of SEQ ID NO:7 and a termination codon spanning from about nucleotide 1199 through about nucleotide 1201 of SEQ ID NO:7. The coding region encoding PfSPI2₃₉₉ is represented

by nucleic acid molecule nfSPI2₁₁₉₇, having a coding strand with the nucleic acid sequence represented by SEQ ID NO:10 and a complementary strand with the nucleic acid sequence represented by SEQ ID NO:11. Analysis of SEQ ID NO:8 suggests the presence of a partial signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 23. The proposed mature protein, denoted herein as PfSPI2₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:12. The amino acid sequence of flea PfSPI1₃₇₆ (i.e. SEQ ID NO:12) predicts that PfSPI2₃₇₆ has an estimated molecular weight of about 42.1 kD, an estimated pI of about 4.87, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

Comparison of amino acid sequence SEQ ID NO:8 (i.e., the amino acid sequence of PfSPI2₃₉₉) with amino acid sequences reported in GenBank indicates that SEQ ID NO:8, showed the most homology, i.e., about 36% identity, with GenBank accession number 1345616, a serpin protein from *Homo sapiens*.

Translation of SEQ ID NO:13 suggests that nucleic acid molecule nfSPI3₁₈₃₈ encodes a full-length flea SPI protein of about 420 amino acids, referred to herein as PfSPI3₄₂₀, represented by SEQ ID NO:14, assuming an open reading frame having an initiation codon spanning from about nucleotide 306 through about nucleotide 308 of SEQ ID NO:13 and a termination codon spanning from about nucleotide 1566 through about nucleotide 1568 of SEQ ID NO:13. The coding region encoding PfSPI3₄₂₀ is represented by nucleic acid molecule nfSPI3₁₂₆₀, having a coding strand with the nucleic acid sequence represented by SEQ ID NO:16 and a complementary strand with the nucleic acid sequence represented by SEQ ID NO:17. The deduced amino acid sequence SEQ ID NO:14 suggests a protein having a molecular weight of about 47.1 kilodaltons (kD) and an estimated pI of about 4.72. Analysis of SEQ ID NO:14 suggests the presence of a signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 30. The proposed mature protein, denoted herein as PfSPI3₃₉₀, contains about 390 amino acids which is represented herein as SEQ ID NO:18. The amino acid sequence of flea PfSPI3₃₉₀ (i.e. SEQ ID NO:18) predicts that PfSPI3₃₉₀ has an estimated molecular weight of about 43.7 kD, an estimated pI of about 4.63, and two predicted asparagine-linked glycosylation sites extending from about

amino acid 252 to about amino acid 254 and from about amino acid 369 to about amino acid 371.

Comparison of amino acid sequence SEQ ID NO:14 (i.e., the amino acid sequence of PfSPI3₄₂₀) with amino acid sequences reported in GenBank indicates that
5 SEQ ID NO:14, showed the most homology, i.e., about 35% identity, with GenBank accession number 1345616, a serpin protein from *Homo sapiens*.

Translation of SEQ ID NO:19 suggests that nucleic acid molecule nfSPI4₁₄₁₄ encodes a non-full-length flea SPI protein of about 393 amino acids, referred to herein as PfSPI4₃₉₃, represented by SEQ ID NO:20, assuming an open reading frame having a first
10 in-frame codon spanning from about nucleotide 2 through about nucleotide 4 of SEQ ID NO:19 and a termination codon spanning from about nucleotide 1181 through about nucleotide 1183 of SEQ ID NO:19. The coding region encoding PfSPI4₃₉₃, is represented by nucleic acid molecule nfSPI4₁₁₇₉, having a coding strand with the nucleic acid sequence represented by SEQ ID NO:22 and a complementary strand with the
15 nucleic acid sequence represented by SEQ ID NO:23. Analysis of SEQ ID NO:20 suggests the presence of a partial signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 17. The proposed mature protein, denoted herein as PfSPI4₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:24. The amino acid sequence of flea PfSPI4₃₇₆ (i.e.
20 SEQ ID NO:24) predicts that PfSPI4₃₇₆ has an estimated molecular weight of about 42.2 kD, an estimated pI of about 5.31, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

Comparison of amino acid sequence SEQ ID NO:20 (i.e., the amino acid sequence of PfSPI4₃₉₃) with amino acid sequences reported in GenBank indicates that
25 SEQ ID NO:20, showed the most homology, i.e., about 38% identity, with GenBank accession number 1345616, a serpin protein from *Homo sapiens*.

Translation of SEQ ID NO:25 suggests that nucleic acid molecule nfSPI5₁₄₉₂ encodes a non-full-length flea SPI protein of about 398 amino acids, referred to herein as PfSPI5₃₉₈, represented by SEQ ID NO:26, assuming an open reading frame having a first
30 in-frame codon spanning from about nucleotide 3 through about nucleotide 5 of SEQ ID NO:25 and a termination codon spanning from about nucleotide 1197 through about

nucleotide 1199 of SEQ ID NO:25. The coding region encoding PfSPI5₃₉₈, is represented by nucleic acid molecule nfSPI5₁₁₉₄, having a coding strand with the nucleic acid sequence represented by SEQ ID NO:28 and a complementary strand with the nucleic acid sequence represented by SEQ ID NO:29. Analysis of SEQ ID NO:26 suggests the presence of a partial signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 22. The proposed mature protein, denoted herein as PfSPI5₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:30. The amino acid sequence of flea PfSPI5₃₇₆ (i.e. SEQ ID NO:30) predicts that PfSPI5₃₇₆ has an estimated molecular weight of about 42.3 kD, an estimated pI of about 5.31 and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

Comparison of amino acid sequence SEQ ID NO:26 (i.e., the amino acid sequence of PfSPI5₃₉₈) with amino acid sequences reported in GenBank indicates that SEQ ID NO:26 showed the most homology, i.e., about 38% identity with GenBank accession number 1345616, a serpin protein from *Homo sapiens*.

Translation of SEQ ID NO:31 suggests that nucleic acid molecule nfSPI6₁₄₅₄ encodes a full-length flea SPI protein of about 397 amino acids, referred to herein as PfSPI6₃₉₇, represented by SEQ ID NO:32, assuming an open reading frame having an initiation codon spanning from about nucleotide 20 through about nucleotide 22 of SEQ ID NO:31 and a termination codon spanning from about nucleotide 1211 through about nucleotide 1213 of SEQ ID NO:31. The coding region encoding PfSPI6₃₉₇ is represented by nucleic acid molecule nfSPI6₁₁₉₁, having a coding strand with the nucleic acid sequence represented by SEQ ID NO:34 and a complementary strand with the nucleic acid sequence represented by SEQ ID NO:35. The deduced amino acid sequence SEQ ID NO:32 suggests a protein having a molecular weight of about 44.4 kilodaltons (kD) and an estimated pI of about 4.90. Analysis of SEQ ID NO:32 suggests the presence of a signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 21. The proposed mature protein, denoted herein as PfSPI6₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:36. The amino acid sequence of flea PfSPI6₃₇₆ (i.e. SEQ ID NO:36) predicts that PfSPI6₃₇₆ has an estimated molecular weight of about 42.1 kD, an estimated pI of about 4.84, and a

predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

Comparison of amino acid sequence SEQ ID NO:32 (i.e., the amino acid sequence of PfSPI6₃₉₇) with amino acid sequences reported in GenBank indicates that
5 SEQ ID NO:32 showed the most homology, i.e., about 36% identity with GenBank accession number 1378131, a serpin protein from *Manduca sexta*.

Translation of SEQ ID NO:45 suggests that nucleic acid molecule nfSPI7₅₄₉ encodes a portion of a serine protease inhibitor protein of about 134 amino acids, referred to herein as PfSPI7₁₃₄, having amino acid sequence SEQ ID NO:46, assuming
10 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:45 and the last codon spans from nucleotide 402 through nucleotide 404 of SEQ ID NO:45. The complement of SEQ ID NO:45 is represented herein by SEQ ID NO:47.

Comparison of amino acid sequence SEQ ID NO:46 (i.e., the amino acid sequence of PfSPI7₁₃₄) with amino acid sequences reported in SwissProt indicates that
15 SEQ ID NO:46, showed the most homology, i.e., about 34% identity, between SEQ ID NO:46 and *mus musculus* antithrombin III precursor protein.

Translation of SEQ ID NO:48 suggests that nucleic acid molecule nfSPI8₅₄₉ encodes a serine protease inhibitor variable domain protein of about 149 amino acids, referred to herein as PfSPI8₁₄₉, having amino acid sequence SEQ ID NO:49, assuming
20 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:48 and the last codon spans from nucleotide 447 through nucleotide 449 of SEQ ID NO:48. The complement of SEQ ID NO:48 is represented herein by SEQ ID NO:50.

Comparison of amino acid sequence SEQ ID NO:49 (i.e., the amino acid sequence of PfSPI8₁₄₉) with amino acid sequences reported in SwissProt indicates that
25 SEQ ID NO:49, showed the most homology, i.e., about 36% identity, between SEQ ID NO:49 and human bomapin protein.

Translation of SEQ ID NO:51 suggests that nucleic acid molecule nfSPI9₅₈₁ encodes a serine protease inhibitor variable domain protein of about 136 amino acids, referred to herein as PfSPI9₁₃₆, having amino acid sequence SEQ ID NO:52, assuming
30 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:51 and the

last codon spans from nucleotide 408 through nucleotide 410 of SEQ ID NO:51. The complement of SEQ ID NO:51 is represented herein by SEQ ID NO:53.

Comparison of amino acid sequence SEQ ID NO:52 (i.e., the amino acid sequence of PfSPI9₁₃₆) with amino acid sequences reported in SwissProt indicates that
5 SEQ ID NO:52, showed the most homology, i.e., about 45% identity, between SEQ ID NO:52 and *Bombyx mori* anti-trypsin precursor protein.

Translation of SEQ ID NO:54 suggests that nucleic acid molecule nfSPI10₆₅₄ encodes a serine protease inhibitor variable domain protein of about 118 amino acids, referred to herein as PfSPI10₁₁₈, having amino acid sequence SEQ ID NO:55, assuming
10 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:54 and the last codon spans from nucleotide 354 through nucleotide 356 of SEQ ID NO:54. The complement of SEQ ID NO:54 is represented herein by SEQ ID NO:56.

Comparison of amino acid sequence SEQ ID NO:55 (i.e., the amino acid sequence of PfSPI10₁₁₈) with amino acid sequences reported in SwissProt indicates that
15 SEQ ID NO:55, showed the most homology, i.e., about 38% identity, between SEQ ID NO:55 and *Manduca sexta* alaserpin precursor protein.

Translation of SEQ ID NO:57 suggests that nucleic acid molecule nfSPI11₆₇₀ encodes a serine protease inhibitor variable domain protein of about 125 amino acids, referred to herein as PfSPI11₁₂₅, having amino acid sequence SEQ ID NO:58, assuming
20 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:57 and the last codon spans from nucleotide 375 through nucleotide 377 of SEQ ID NO:57. The complement of SEQ ID NO:57 is represented herein by SEQ ID NO:59.

Comparison of amino acid sequence SEQ ID NO:58 (i.e., the amino acid sequence of PfSPI11₁₂₅) with amino acid sequences reported in SwissProt indicates that
25 SEQ ID NO:58, showed the most homology, i.e., about 43% identity, between SEQ ID NO:58 and *Manduca sexta* alaserpin precursor protein.

Translation of SEQ ID NO:60 suggests that nucleic acid molecule nfSPI12₇₀₆ encodes a serine protease inhibitor variable domain protein of about 136 amino acids, referred to herein as PfSPI12₁₃₆, having amino acid sequence SEQ ID NO:61, assuming
30 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:60 and the

last codon spans from nucleotide 408 through nucleotide 410 of SEQ ID NO:60. The complement of SEQ ID NO:60 is represented herein by SEQ ID NO:62.

Comparison of amino acid sequence SEQ ID NO:61 (i.e., the amino acid sequence of PfSPI12₁₃₆) with amino acid sequences reported in SwissProt indicates that
5 SEQ ID NO:61, showed the most homology, i.e., about 45% identity, between SEQ ID NO:61 and *Manduca sexta* alaserpin precursor protein protein.

Translation of SEQ ID NO:63 suggests that nucleic acid molecule nfSPI13₆₂₃ encodes a serine protease inhibitor variable domain protein of about 122 amino acids, referred to herein as PfSPI13₁₂₂, having amino acid sequence SEQ ID NO:64, assuming
10 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:63 and the last codon spans from nucleotide 366 through nucleotide 368 of SEQ ID NO:63. The complement of SEQ ID NO:63 is represented herein by SEQ ID NO:65.

Comparison of amino acid sequence SEQ ID NO:64 (i.e., the amino acid sequence of PfSPI13₁₂₂) with amino acid sequences reported in SwissProt indicates that
15 SEQ ID NO:64, showed the most homology, i.e., about 39% identity, between SEQ ID NO:64 and human leukocyte esterase inhibitor protein.

Translation of SEQ ID NO:66 suggests that nucleic acid molecule nfSPI14₇₃₁ encodes a serine protease inhibitor variable domain protein of about 137 amino acids, referred to herein as PfSPI14₁₃₇, having amino acid sequence SEQ ID NO:67, assuming
20 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:66 and the last codon spans from nucleotide 411 through nucleotide 413 of SEQ ID NO:66. The complement of SEQ ID NO:66 is represented herein by SEQ ID NO:68.

Comparison of amino acid sequence SEQ ID NO:67 (i.e., the amino acid sequence of PfSPI14₁₃₇) with amino acid sequences reported in SwissProt indicates that
25 SEQ ID NO:67, showed the most homology, i.e., about 40% identity, between SEQ ID NO:67 and *Equus caballus* esterase inhibitor protein.

Translation of SEQ ID NO:69 suggests that nucleic acid molecule nfSPI15₆₈₅ encodes a serine protease inhibitor variable domain protein of about 135 amino acids, referred to herein as PfSPI15₁₃₅, having amino acid sequence SEQ ID NO:70, assuming
30 the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:69 and the

last codon spans from nucleotide 405 through nucleotide 407 of SEQ ID NO:69. The complement of SEQ ID NO:69 is represented herein by SEQ ID NO:71.

Comparison of amino acid sequence SEQ ID NO:70 (i.e., the amino acid sequence of PfSPI15₁₃₅) with amino acid sequences reported in SwissProt indicates that
 5 SEQ ID NO:70, showed the most homology, i.e., about 48% identity, between SEQ ID NO:70 and *Bombyx mori* antichymotrypsin II protein.

More preferred flea SPI proteins of the present invention include proteins comprising amino acid sequences that are at least about 40%, preferably at least about 50%, more preferably at least about 60%, more preferably at least about 70%, more
 10 preferably at least about 80%, and even more preferably at least about 90%, identical to amino acid sequence SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID
 15 NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89 and/or SEQ ID NO:90.

More preferred flea SPI proteins of the present invention include proteins encoded by a nucleic acid molecule comprising at least a portion of nfSPI1₁₅₈₄, nfSPI1₁₁₉₁, nfSPI1₃₇₆, nfSPI2₁₃₅₈, nfSPI2₁₁₉₇, nfSPI2₃₇₆, nfSPI3₁₈₃₈, nfSPI3₁₂₆₀, nfSPI3₃₉₁, nfSPI4₁₄₁₄, nfSPI4₁₁₇₉, nfSPI4₃₇₆, nfSPI5₁₄₉₂, nfSPI5₁₁₉₄, nfSPI5₃₇₆, nfSPI6₁₄₅₄, nfSPI6₁₁₉₁,
 20 nfSPI6₃₇₆, nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI11₆₇₀, nfSPI12₇₀₆, nfSPI13₆₂₃, nfSPI14₇₃₁, nfSPI15₆₈₅, nfSPI3₁₂₂₂, nfSPI6₁₁₅₅, nfSPI2₁₀₆₅, nfSPI4₁₀₇₀, nfSPIC4:V7₁₁₆₈, nfSPIC4:V8₁₂₂₂, nfSPIC4:V9₁₁₇₄, nfSPIC4:V10₁₁₅₉, nfSPIC4:V12₁₁₇₁, nfSPIC4:V13₁₁₇₁, and nfSPIC4:V15₁₁₇₉, or by an allelic variant of such nucleic acid molecules.

Particularly preferred flea SPI proteins are PfSPI1₃₉₇, PfSPI1₃₇₆, PfSPI2₃₉₉, PfSPI2₃₇₆,
 25 PfSPI2₃₅₄, PfSPI3₄₀₆, PfSPI3₄₂₀, PfSPI3₃₉₁, PfSPI4₃₉₃, PfSPI4₃₇₆, PfSPI4₃₅₆, PfSPI5₃₉₈, PfSPI5₃₇₆, PfSPI6₃₉₇, PfSPI6₃₇₆, PfSPI6₃₈₅, PfSPI2₃₅₅, PfSPI3₄₀₆, PfSPI4₃₅₆, PfSPI6₃₈₅, PfSPI7₁₃₄, PfSPI8₁₄₉, PfSPI9₁₃₆, PfSPI10₁₁₈, PfSPI11₁₂₅, PfSPI12₁₃₆, PfSPI13₁₂₂, PfSPI14₁₃₇, PfSPI15₁₃₅, PHis-PfSPIC4:V7, PHis-PfSPIC4:V8, PHis-PfSPIC4:V9, PHis-PfSPIC4:V10, PHis-PfSPIC4:V12, PHis-PfSPIC4:V13, PHis-PfSPIC4:V15.

30 In one embodiment, a preferred SPI protein of the present invention is encoded by at least a portion of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10,

SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:57, SEQ ID NO:60, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81 and/or a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90, and, as such, has an amino acid sequence that includes at least a portion of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90, respectively.

Also preferred is a protein encoded by an allelic variant of a nucleic acid molecule comprising at least a portion of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:57, SEQ ID NO:60, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, and/or a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90. Particularly preferred SPI proteins of the present invention include SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97, and/or SEQ ID NO:98 (including, but not limited to, the proteins consisting of such sequences, fusion proteins and multivalent proteins) and proteins encoded by allelic variants of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:51, SEQ ID

NO:54, SEQ ID NO:57, SEQ ID NO:60, SEQ ID NO:63, SEQ ID NO:66, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, and/or a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90.

5 Another embodiment of the present invention is an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a *C. felis* SPI gene. The identifying characteristics of such a gene are heretofore described. A nucleic acid molecule of the present invention can include an isolated natural flea SPI gene or a homolog thereof, the latter of which is described in more detail below. A nucleic acid
10 molecule of the present invention can include one or more regulatory regions, full-length or partial coding regions, or combinations thereof. The minimal size of a nucleic acid molecule of the present invention is the minimal size that can form a stable hybrid with a *C. felis* SPI gene under stringent hybridization conditions.

 In accordance with the present invention, an isolated nucleic acid molecule is a
15 nucleic acid molecule that has been removed from its natural milieu (i.e., that has been subject to human manipulation) and can include DNA, RNA, or derivatives of either DNA or RNA. As such, "isolated" does not reflect the extent to which the nucleic acid molecule has been purified. An isolated flea SPI nucleic acid molecule of the present invention can be isolated from its natural source or can be produced using recombinant
20 DNA technology (e.g., polymerase chain reaction (PCR) amplification, cloning) or chemical synthesis. Isolated SPI nucleic acid molecules can include, for example, natural allelic variants and nucleic acid molecules modified by nucleotide insertions, deletions, substitutions, and/or inversions in a manner such that the modifications do not substantially interfere with the nucleic acid molecule's ability to encode a SPI protein of
25 the present invention or to form stable hybrids under stringent conditions with natural gene isolates.

 A flea SPI nucleic acid molecule homolog can be produced using a number of methods known to those skilled in the art (see, for example, Sambrook et al., *ibid.*). For example, nucleic acid molecules can be modified using a variety of techniques
30 including, but not limited to, classic mutagenesis and recombinant DNA techniques (e.g., site-directed mutagenesis, chemical treatment, restriction enzyme cleavage,

ligation of nucleic acid fragments and/or PCR amplification), synthesis of oligonucleotide mixtures and ligation of mixture groups to "build" a mixture of nucleic acid molecules and combinations thereof. Nucleic acid molecule homologs can be selected by hybridization with a *C. felis* SPI gene or by screening for function of a
5 protein encoded by the nucleic acid molecule (e.g., ability to elicit an immune response against at least one epitope of a flea SPI protein or has at least some serine protease inhibitor activity).

An isolated nucleic acid molecule of the present invention can include a nucleic acid sequence that encodes at least one flea SPI protein of the present invention,
10 examples of such proteins being disclosed herein. Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides on the nucleic acid molecule, the two phrases can be used interchangeably, especially with respect to a nucleic acid molecule, or a nucleic acid sequence, being capable of encoding a flea SPI
15 protein.

A preferred nucleic acid molecule of the present invention, when administered to an animal, is capable of protecting that animal from infestation by a hematophagous ectoparasite. As will be disclosed in more detail below, such a nucleic acid molecule can be, or can encode, an antisense RNA, a molecule capable of triple helix formation, a
20 ribozyme, or other nucleic acid-based drug compound. In additional embodiments, a nucleic acid molecule of the present invention can encode a protective protein (e.g., a SPI protein of the present invention), the nucleic acid molecule being delivered to the animal, for example, by direct injection (i.e., as a naked nucleic acid) or in a vehicle such as a recombinant virus vaccine or a recombinant cell vaccine.

25 One embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI1₁₅₈₄ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:1 and/or SEQ ID NO:3.

Another embodiment of the present invention is a SPI nucleic acid molecule that
30 hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI2₁₃₅₈

and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:7 and/or SEQ ID NO:9.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI3₁₈₃₈ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:13 and/or SEQ ID NO:15.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI4₁₄₁₄ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:19 and/or SEQ ID NO:21.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI5₁₄₉₂ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:25 and/or SEQ ID NO:27.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI6₁₄₅₄ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:31 and/or SEQ ID NO:33.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI7₅₄₉ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:45 and/or SEQ ID NO:47.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI8₅₄₉ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:48 and/or SEQ ID NO:50.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI9₅₈₁ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:51 and/or SEQ ID NO:53.

Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI10₆₅₄ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:54 and/or SEQ ID NO:56.

- 5 Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI11₆₇₀ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:57 and/or SEQ ID NO:59.

- 10 Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI12₇₀₆ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:60 and/or SEQ ID NO:62.

- 15 Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI13₆₂₃ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:63 and/or SEQ ID NO:65.

- 20 Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI14₇₃₁ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:66 and/or SEQ ID NO:68.

- 25 Another embodiment of the present invention is a SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with nucleic acid molecule nfSPI15₆₈₅ and preferably with a nucleic acid molecule having nucleic acid sequence SEQ ID NO:69 and/or SEQ ID NO:71.

- 30 Comparison of nucleic acid sequence SEQ ID NO:4 (i.e., the nucleic acid sequence of the coding strand of nfSPI1₁₁₉₁) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:4 showed the most homology, i.e., about 55% identity, with accession number L20792, a putative serine proteinase inhibitor (serpin 1, exon 9 copy 2) gene of *Manduca sexta*.

- 30 Comparison of nucleic acid sequence SEQ ID NO:10 (i.e., the nucleic acid sequence of the coding strand of nfSPI2₁₁₉₇) with nucleic acid sequences reported in

GenBank indicates that SEQ ID NO:10 showed the most homology, i.e., about 43% identity, with accession number L20790, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 1) of *Manduca sexta*.

5 Comparison of nucleic acid sequence SEQ ID NO:16 (i.e., the nucleic acid sequence of the coding strand of nfSPI3₁₂₆₀) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:16 showed the most homology, i.e., about 52% identity, with accession number L20792, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 2) of *Manduca sexta*.

10 Comparison of nucleic acid sequence SEQ ID NO:22 (i.e., the nucleic acid sequence of the coding strand of nfSPI4₁₁₇₉) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:22 showed the most homology, i.e., about 55% identity, with accession number L20793, a putative serine proteinase inhibitor gene (serpin 1, exon 9 unknown copy number) of *Manduca sexta*.

15 Comparison of nucleic acid sequence SEQ ID NO:28 (i.e., the nucleic acid sequence of the coding strand of nfSPI5₁₁₉₄) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:28 showed the most homology, i.e., about 45% identity, with accession number L20790, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 1) of *Manduca sexta*.

20 Comparison of nucleic acid sequence SEQ ID NO:34 (i.e., the nucleic acid sequence of the coding strand of nfSPI6₁₁₉₁) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:34 showed the most homology, i.e., about 55% identity, with accession number L20792, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 2) of *Manduca sexta*.

25 Comparison of nucleic acid sequence SEQ ID NO:45 (i.e., the nucleic acid sequence of nfSPI7₅₄₉) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:45, showed the most homology, i.e., about 38% identity, between SEQ ID NO:45 and human bomapin gene.

30 Comparison of nucleic acid sequence SEQ ID NO:48 (i.e., the nucleic acid sequence of nfSPI8₅₄₉) with nucleic acid sequences reported in GeEmbl indicates that SEQ ID NO:48, showed the most homology, i.e., about 41% identity, between SEQ ID NO:48 and human bomapin gene.

Comparison of nucleic acid sequence SEQ ID NO:51 (i.e., the nucleic acid sequence of nfSPI9₅₈₁) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:51, showed the most homology, i.e., about 52% identity, between SEQ ID NO:51 and *Bombyx mori* anti-trypsin gene.

- 5 Comparison of nucleic acid sequence SEQ ID NO:54 (i.e., the nucleic acid sequence of nfSPI10₆₅₄) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:54, showed the most homology, i.e., about 41% identity, between SEQ ID NO:54 and human bomapin gene.

- 10 Comparison of nucleic acid sequence SEQ ID NO:57 (i.e., the nucleic acid sequence of nfSPI11₆₇₀) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:57, showed the most homology, i.e., about 40% identity, between SEQ ID NO:57 and human bomapin gene.

- 15 Comparison of nucleic acid sequence SEQ ID NO:60 (i.e., the nucleic acid sequence of nfSPI12₇₀₆) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:60, showed the most homology, i.e., about 38% identity, between SEQ ID NO:60 and human bomapin gene.

- 20 Comparison of nucleic acid sequence SEQ ID NO:63 (i.e., the nucleic acid sequence of nfSPI13₆₂₃) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:63, showed the most homology, i.e., about 37% identity, between SEQ ID NO:63 and human bomapin gene.

Comparison of nucleic acid sequence SEQ ID NO:66 (i.e., the nucleic acid sequence of nfSPI14₇₃₁) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:66, showed the most homology, i.e., about 38% identity, between SEQ ID NO:66 and human bomapin gene.

- 25 Comparison of nucleic acid sequence SEQ ID NO:69 (i.e., the nucleic acid sequence of nfSPI15₆₈₅) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:69, showed the most homology, i.e., about 38% identity, between SEQ ID NO:69 and human antithrombin III variant gene.

- 30 Preferred flea SPI nucleic acid molecules include nucleic acid molecules having a nucleic acid sequence that is at least about 60%, preferably at least about 70%, more preferably at least about 80%, even more preferably at least about 90% and even more

preferably at least about 95% identical to nucleic acid sequence SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90.

Another preferred nucleic acid molecule of the present invention includes at least a portion of nucleic acid sequence SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90, that is capable of hybridizing to a *C. felis* SPI gene of the present invention, as well as allelic variants thereof. A more preferred nucleic acid molecule includes the nucleic acid sequence SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID

NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid
 5 sequence that encodes an amino acid sequence including SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90, as well as allelic variants thereof. Such nucleic acid molecules can include nucleotides in addition to those included in the SEQ ID NOs, such as, but not limited to, a full-length gene, a full-length coding region, a nucleic acid molecule encoding a fusion protein, or a nucleic acid molecule encoding a multivalent
 10 protective compound. Particularly preferred nucleic acid molecules include nfSPI1₁₅₈₄, nfSPI1₁₁₉₁, nfSPI1₃₇₆, nfSPI2₁₃₅₈, nfSPI2₁₁₉₇, nfSPI2₃₇₆, nfSPI3₁₈₃₈, nfSPI3₁₂₆₀, nfSPI3₃₉₁, nfSPI4₁₄₁₄, nfSPI4₁₁₇₉, nfSPI4₃₇₆, nfSPI5₁₄₉₂, nfSPI5₁₁₉₄, nfSPI5₃₇₆, nfSPI6₁₄₅₄, nfSPI6₁₁₉₁, nfSPI6₃₇₆, nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI11₆₇₀, nfSPI12₇₀₆, nfSPI13₆₂₃, nfSPI14₇₃₁, nfSPI15₆₈₅, nfSPI3₁₂₂₂, nfSPI6₁₁₅₅, nfSPI2₁₀₆₅, nfSPI4₁₀₇₀, nfSPIC4:V7₁₁₆₈,
 15 nfSPIC4:V8₁₂₂₂, nfSPIC4:V9₁₁₇₄, nfSPIC4:V10₁₁₅₉, nfSPIC4:V12₁₁₇₁, nfSPIC4:V13₁₁₇₁, and nfSPIC4:V15₁₁₇₉.

The present invention also includes a nucleic acid molecule encoding a protein having at least a portion of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID
 20 NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO: 97, and SEQ ID NO:98, including nucleic acid molecules that have been modified to accommodate codon usage properties
 25 of the cells in which such nucleic acid molecules are to be expressed.

Knowing the nucleic acid sequences of certain flea SPI nucleic acid molecules of the present invention allows one skilled in the art to, for example, (a) make copies of those nucleic acid molecules, (b) obtain nucleic acid molecules including at least a portion of such nucleic acid molecules (e.g., nucleic acid molecules including full-length
 30 genes, full-length coding regions, regulatory control sequences, truncated coding regions), and (c) obtain SPI nucleic acid molecules from other hematophagous

ectoparasites. Such nucleic acid molecules can be obtained in a variety of ways including screening appropriate expression libraries with antibodies of the present invention; traditional cloning techniques using oligonucleotide probes of the present invention to screen appropriate libraries or DNA; and PCR amplification of appropriate
5 libraries or DNA using oligonucleotide primers of the present invention. Preferred libraries to screen or from which to amplify nucleic acid molecule include flea hemocyte (i.e., cells found in flea hemolymph), pre-pupal, mixed instar (i.e., a combination of 1st instar larval, 2nd instar larval, 3rd instar larval tissue), or fed or unfed adult cDNA libraries as well as genomic DNA libraries. Similarly, preferred DNA sources to screen
10 or from which to amplify nucleic acid molecules include flea hemocyte, pre-pupal, mixed instar, or fed or unfed adult cDNA and genomic DNA. Techniques to clone and amplify genes are disclosed, for example, in Sambrook et al., *ibid*.

The present invention also includes nucleic acid molecules that are oligonucleotides capable of hybridizing, under stringent hybridization conditions, with
15 complementary regions of other, preferably longer, nucleic acid molecules of the present invention such as those comprising flea SPI genes or other flea SPI nucleic acid molecules. Oligonucleotides of the present invention can be RNA, DNA, or derivatives of either. The minimum size of such oligonucleotides is the size required for formation of a stable hybrid between an oligonucleotide and a complementary sequence on a
20 nucleic acid molecule of the present invention. Minimal size characteristics are disclosed herein. The present invention includes oligonucleotides that can be used as, for example, probes to identify nucleic acid molecules, primers to produce nucleic acid molecules or therapeutic reagents to inhibit SPI protein production or activity (e.g., as antisense-, triplex formation-, ribozyme- and/or RNA drug-based reagents). The present
25 invention also includes the use of such oligonucleotides to protect animals from disease using one or more of such technologies. Appropriate oligonucleotide-containing therapeutic compositions can be administered to an animal using techniques known to those skilled in the art.

One embodiment of the present invention includes a recombinant vector, which
30 includes at least one isolated nucleic acid molecule of the present invention, inserted into any vector capable of delivering the nucleic acid molecule into a host cell. Such a vector

contains heterologous nucleic acid sequences, that is nucleic acid sequences that are not naturally found adjacent to nucleic acid molecules of the present invention and that preferably are derived from a species other than the species from which the nucleic acid molecule(s) are derived. The vector can be either RNA or DNA, either prokaryotic or eukaryotic, and typically is a virus or a plasmid. Recombinant vectors can be used in the cloning, sequencing, and/or otherwise manipulation of flea SPI nucleic acid molecules of the present invention.

One type of recombinant vector, referred to herein as a recombinant molecule, comprises a nucleic acid molecule of the present invention operatively linked to an expression vector. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be expressed when transformed into a host cell. As used herein, an expression vector is a DNA or RNA vector that is capable of transforming a host cell and of effecting expression of a specified nucleic acid molecule. Preferably, the expression vector is also capable of replicating within the host cell. Expression vectors can be either prokaryotic or eukaryotic, and are typically viruses or plasmids. Expression vectors of the present invention include any vectors that function (i.e., direct gene expression) in recombinant cells of the present invention, including in bacterial, fungal, endoparasite, insect, other animal, and plant cells. Preferred expression vectors of the present invention can direct gene expression in bacterial, yeast, insect and mammalian cells and more preferably in the cell types disclosed herein.

In particular, expression vectors of the present invention contain regulatory sequences such as transcription control sequences, translation control sequences, origins of replication, and other regulatory sequences that are compatible with the recombinant cell and that control the expression of nucleic acid molecules of the present invention. In particular, recombinant molecules of the present invention include transcription control sequences. Transcription control sequences are sequences which control the initiation, elongation, and termination of transcription. Particularly important transcription control sequences are those which control transcription initiation, such as promoter, enhancer, operator and repressor sequences. Suitable transcription control sequences include any transcription control sequence that can function in at least one of

the recombinant cells of the present invention. A variety of such transcription control sequences are known to those skilled in the art. Preferred transcription control sequences include those which function in bacterial, yeast, insect and mammalian cells, such as, but not limited to, *tac*, *lac*, *trp*, *trc*, oxy-pro, omp/lpp, *rnnB*, bacteriophage lambda (such as lambda p_L and lambda p_R and fusions that include such promoters), bacteriophage T7, T7*lac*, bacteriophage T3, bacteriophage SP6, bacteriophage SP01, metallothionein, alpha-mating factor, *Pichia* alcohol oxidase, alphavirus subgenomic promoters (such as Sindbis virus subgenomic promoters), antibiotic resistance gene, baculovirus, *Heliothis zea* insect virus, vaccinia virus, herpesvirus, raccoon poxvirus, other poxvirus, adenovirus, cytomegalovirus (such as intermediate early promoters), simian virus 40, retrovirus, actin, retroviral long terminal repeat, Rous sarcoma virus, heat shock, phosphate and nitrate transcription control sequences as well as other sequences capable of controlling gene expression in prokaryotic or eukaryotic cells. Additional suitable transcription control sequences include tissue-specific promoters and enhancers as well as lymphokine-inducible promoters (e.g., promoters inducible by interferons or interleukins). Transcription control sequences of the present invention can also include naturally occurring transcription control sequences naturally associated with fleas, such as, *C. felis*.

Suitable and preferred nucleic acid molecules to include in recombinant vectors of the present invention are as disclosed herein. Preferred nucleic acid molecules to include in recombinant vectors, and particularly in recombinant molecules, include nfSPI1₁₅₈₄, nfSPI1₁₁₉₁, nfSPI1₃₇₆, nfSPI2₁₃₅₈, nfSPI2₁₁₉₇, nfSPI2₃₇₆, nfSPI3₁₈₃₈, nfSPI3₁₂₆₀, nfSPI3₃₉₁, nfSPI4₁₄₁₄, nfSPI4₁₁₇₉, nfSPI4₃₇₆, nfSPI5₁₄₉₂, nfSPI5₁₁₉₄, nfSPI5₃₇₆, nfSPI6₁₄₅₄, nfSPI6₁₁₉₁, nfSPI6₃₇₆, nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI11₆₇₀, nfSPI12₇₀₆, nfSPI13₆₂₃, nfSPI14₇₃₁, nfSPI15₆₈₅, nfSPI3₁₂₂₂, nfSPI6₁₁₅₅, nfSPI2₁₀₆₅, nfSPI4₁₀₇₀, nfSPIC4:V7₁₁₆₈, nfSPIC4:V8₁₂₂₂, nfSPIC4:V9₁₁₇₄, nfSPIC4:V10₁₁₅₉, nfSPIC4:V12₁₁₇₁, nfSPIC4:V13₁₁₇₁, and nfSPIC4:V15₁₁₇₉. Particularly preferred recombinant molecules of the present invention include p λP_R -nfSPI2₁₁₃₉, p λP_R -nfSPI3₁₁₇₉, p λP_R -nfSPI4₁₁₄₀, p λP_R -nfSPI5₁₄₉₂, p λP_R -nfSPI6₁₁₃₆, p λP_R -nfSPIC4:V7₁₁₆₈, p λP_R -nfSPIC4:V8₁₂₂₂, p λP_R -nfSPIC4:V9₁₁₇₄, p λP_R -nfSPIC4:V10₁₁₅₉, p λP_R -nfSPIC4:V12₁₁₇₁, p λP_R -nfSPIC4:V13₁₁₇₁,

p λ P_R-nfSPIC4:V15₁₁₇₉, pVL-nfSPI3₁₂₂₂, pVL-nfSPI6₁₁₅₅, pAcG-nfSPI2₁₀₆₅ and pAcG-nfSPI4₁₀₇₀, the production of which are described in the Examples section.

Recombinant molecules of the present invention may also (a) contain secretory signals (i.e., signal segment nucleic acid sequences) to enable an expressed flea protein of the present invention to be secreted from the cell that produces the protein and/or (b) contain fusion sequences which lead to the expression of nucleic acid molecules of the present invention as fusion proteins. Examples of suitable signal segments include any signal segment capable of directing the secretion of a protein of the present invention. Preferred signal segments include, but are not limited to, tissue plasminogen activator (t-PA), interferon, interleukin, growth hormone, histocompatibility and viral envelope glycoprotein signal segments, as well as natural signal segments. Suitable fusion segments encoded by fusion segment nucleic acids are disclosed herein. In addition, a nucleic acid molecule of the present invention can be joined to a fusion segment that directs the encoded protein to the proteosome, such as a ubiquitin fusion segment.

Recombinant molecules may also include intervening and/or untranslated sequences surrounding and/or within the nucleic acid sequences of nucleic acid molecules of the present invention.

Another embodiment of the present invention includes a recombinant cell comprising a host cell transformed with one or more recombinant molecules of the present invention. Transformation of a nucleic acid molecule into a cell can be accomplished by any method by which a nucleic acid molecule can be inserted into the cell. Transformation techniques include, but are not limited to, transfection, electroporation, microinjection, lipofection, adsorption, and protoplast fusion. A recombinant cell may remain unicellular or may grow into a tissue, organ or a multicellular organism. Transformed nucleic acid molecules of the present invention can remain extrachromosomal or can integrate into one or more sites within a chromosome of the transformed (i.e., recombinant) cell in such a manner that their ability to be expressed is retained. Preferred nucleic acid molecules with which to transform a cell include flea SPI nucleic acid molecules disclosed herein. Particularly preferred nucleic acid molecules with which to transform a cell include nfSPI1₁₅₈₄, nfSPI1₁₁₉₁, nfSPI1₃₇₆, nfSPI2₁₃₅₈, nfSPI2₁₁₉₇, nfSPI2₃₇₆, nfSPI3₁₈₃₈, nfSPI3₁₂₆₀, nfSPI3₃₉₁,

nfSPI4₁₄₁₄, nfSPI4₁₁₇₉, nfSPI4₃₇₆, nfSPI5₁₄₉₂, nfSPI5₁₁₉₄, nfSPI5₃₇₆, nfSPI6₁₄₅₄, nfSPI6₁₁₉₁,
 nfSPI6₃₇₆, nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI11₆₇₀, nfSPI12₇₀₆, nfSPI13₆₂₃,
 nfSPI14₇₃₁, nfSPI15₆₈₅, nfSPI13₁₂₂₂, nfSPI6₁₁₅₅, nfSPI2₁₀₆₅, nfSPI4₁₀₇₀, nfSPIC4:V7₁₁₆₈,
 nfSPIC4:V8₁₂₂₂, nfSPIC4:V9₁₁₇₄, nfSPIC4:V10₁₁₅₉, nfSPIC4:V12₁₁₇₁, nfSPIC4:V13₁₁₇₁,
 5 and nfSPIC4:V15₁₁₇₉.

Suitable host cells to transform include any cell that can be transformed with a nucleic acid molecule of the present invention. Host cells can be either untransformed cells or cells that are already transformed with at least one nucleic acid molecule (e.g., nucleic acid molecules encoding one or more proteins of the present invention and/or
 10 other proteins useful in the production of multivalent vaccines). Host cells of the present invention either can be endogenously (i.e., naturally) capable of producing flea SPI proteins of the present invention or can be capable of producing such proteins after being transformed with at least one nucleic acid molecule of the present invention. Host cells of the present invention can be any cell capable of producing at least one protein of the
 15 present invention, and include bacterial, fungal (including yeast), other insect, other animal and plant cells. Preferred host cells include bacterial, mycobacterial, yeast, parasite, insect and mammalian cells. More preferred host cells include *Salmonella*, *Escherichia*, *Bacillus*, *Listeria*, *Saccharomyces*, *Spodoptera*, *Mycobacteria*, *Trichoplusia*, BHK (baby hamster kidney) cells, MDCK cells (normal dog kidney cell
 20 line for canine herpesvirus cultivation), CRFK cells (normal cat kidney cell line for feline herpesvirus cultivation), CV-1 cells (African monkey kidney cell line used, for example, to culture raccoon poxvirus), COS (e.g., COS-7) cells, and Vero cells. Particularly preferred host cells are *Escherichia coli*, including *E. coli* K-12 derivatives; *Salmonella typhi*; *Salmonella typhimurium*, including attenuated strains such as UK-1
 25 x3987 and SR-11 x4072; *Spodoptera frugiperda*; *Trichoplusia ni*; BHK cells; MDCK cells; CRFK cells; CV-1 cells; COS cells; Vero cells; and non-tumorigenic mouse myoblast G8 cells (e.g., ATCC CRL 1246). Additional appropriate mammalian cell hosts include other kidney cell lines, other fibroblast cell lines (e.g., human, murine or chicken embryo fibroblast cell lines), myeloma cell lines, Chinese hamster ovary cells,
 30 mouse NIH/3T3 cells, LMTK³¹ cells and/or HeLa cells. In one embodiment, the proteins

may be expressed as heterologous proteins in myeloma cell lines employing immunoglobulin promoters.

A recombinant cell is preferably produced by transforming a host cell with one or more recombinant molecules, each comprising one or more nucleic acid molecules of the present invention operatively linked to an expression vector containing one or more transcription control sequences. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be expressed when transformed into a host cell.

A recombinant molecule of the present invention is a molecule that can include at least one of any nucleic acid molecule heretofore described operatively linked to at least one of any transcription control sequence capable of effectively regulating expression of the nucleic acid molecule(s) in the cell to be transformed, examples of which are disclosed herein. Particularly preferred recombinant molecules include p λ P_R-nfSPI2₁₁₃₉, p λ P_R-nfSPI3₁₁₇₉, p λ P_R-nfSPI4₁₁₄₀, p λ P_R-nfSPI5₁₄₉₂, p λ P_R-nfSPI6₁₁₃₆, p λ P_R-nfSPIC4:V7₁₁₆₈, p λ P_R-nfSPIC4:V8₁₂₂₂, p λ P_R-nfSPIC4:V9₁₁₇₄, p λ P_R-nfSPIC4:V10₁₁₅₉, p λ P_R-nfSPIC4:V12₁₁₇₁, p λ P_R-nfSPIC4:V13₁₁₇₁, p λ P_R-nfSPIC4:V15₁₁₇₉, pVL-nfSPI3₁₂₂₂, pVL-nfSPI6₁₁₅₅, pAcG-nfSPI2₁₀₆₅ and pAcG-nfSPI4₁₀₇₀.

A recombinant cell of the present invention includes any cell transformed with at least one of any nucleic acid molecule of the present invention. Suitable and preferred nucleic acid molecules as well as suitable and preferred recombinant molecules with which to transform cells are disclosed herein. Particularly preferred recombinant cells include *E.coli*HB:p λ P_R-nfSPI2₁₁₃₉, *E.coli*HB:p λ P_R-nfSPI3₁₁₇₉, *E.coli*HB:p λ P_R-nfSPI4₁₁₄₀, *E.coli*HB:p λ P_R-nfSPI5₁₄₉₂, *E.coli*HB:p λ P_R-nfSPI6₁₁₃₆, *E.coli*:p λ P_R-nfSPIC4:V7₁₁₆₈, *E.coli*:p λ P_R-nfSPIC4:V8₁₂₂₂, *E.coli*:p λ P_R-nfSPIC4:V9₁₁₇₄, *E.coli*:p λ P_R-nfSPIC4:V10₁₁₅₉, *E.coli*:p λ P_R-nfSPIC4:V12₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V13₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V15₁₁₇₉, *S. frugiperda*:pVL-nfSPI3₁₂₂₂, *S. frugiperda*:pVL-nfSPI6₁₁₅₅, *S. frugiperda*:pAcG-nfSPI2₁₀₆₅ and *S. frugiperda*:pAcG-nfSPI4₁₀₇₀. Details regarding the production of these recombinant cells are disclosed herein.

Recombinant cells of the present invention can also be co-transformed with one or more recombinant molecules including flea SPI nucleic acid molecules encoding one or more proteins of the present invention and one or more other nucleic acid molecules

encoding other protective compounds, as disclosed herein (e.g., to produce multivalent vaccines).

Recombinant DNA technologies can be used to improve expression of transformed nucleic acid molecules by manipulating, for example, the number of copies
5 of the nucleic acid molecules within a host cell, the efficiency with which those nucleic acid molecules are transcribed, the efficiency with which the resultant transcripts are translated, and the efficiency of post-translational modifications. Recombinant techniques useful for increasing the expression of nucleic acid molecules of the present invention include, but are not limited to, operatively linking nucleic acid molecules to
10 high-copy number plasmids, integration of the nucleic acid molecules into one or more host cell chromosomes, addition of vector stability sequences to plasmids, substitutions or modifications of transcription control signals (e.g., promoters, operators, enhancers), substitutions or modifications of translational control signals (e.g., ribosome binding sites, Shine-Dalgarno sequences), modification of nucleic acid molecules of the present
15 invention to correspond to the codon usage of the host cell, deletion of sequences that destabilize transcripts, and use of control signals that temporally separate recombinant cell growth from recombinant enzyme production during fermentation. The activity of an expressed recombinant protein of the present invention may be improved by fragmenting, modifying, or derivatizing nucleic acid molecules encoding such a protein.

20 Isolated SPI proteins of the present invention can be produced in a variety of ways, including production and recovery of natural proteins, production and recovery of recombinant proteins, and chemical synthesis of the proteins. In one embodiment, an isolated protein of the present invention is produced by culturing a cell capable of expressing the protein under conditions effective to produce the protein, and recovering
25 the protein. A preferred cell to culture is a recombinant cell of the present invention. Effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. An effective medium refers to any medium in which a cell is cultured to produce a flea SPI protein of the present invention. Such medium typically comprises an aqueous medium having
30 assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present invention can be

cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. Such culturing conditions are within the expertise of one of ordinary skill in the art. Examples of suitable conditions are included
5 in the Examples section.

Depending on the vector and host system used for production, resultant proteins of the present invention may either remain within the recombinant cell; be secreted into the fermentation medium; be secreted into a space between two cellular membranes, such as the periplasmic space in *E. coli*; or be retained on the outer surface of a cell or
10 viral membrane. The phrase "recovering the protein", as well as similar phrases, refers to collecting the whole fermentation medium containing the protein and need not imply additional steps of separation or purification. Proteins of the present invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration,
15 electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization. Proteins of the present invention are preferably retrieved in "substantially pure" form. As used herein, "substantially pure" refers to a purity that allows for the effective use of the protein as a therapeutic composition or diagnostic. A
20 therapeutic composition for animals, for example, should exhibit no substantial toxicity and preferably should be capable of stimulating the production of antibodies in a treated animal.

The present invention also includes isolated (i.e., removed from their natural milieu) antibodies that selectively bind to a flea SPI protein of the present invention or a
25 mimotope thereof (i.e., anti-flea SPI antibodies). As used herein, the term "selectively binds to" a SPI protein refers to the ability of antibodies of the present invention to preferentially bind to specified proteins and mimetopes thereof of the present invention. Binding can be measured using a variety of methods standard in the art including enzyme immunoassays (e.g., ELISA), immunoblot assays, etc.; see, for example,
30 Sambrook et al., *ibid.* An anti-flea SPI antibody preferably selectively binds to a flea SPI protein in such a way as to reduce the activity of that protein.

Isolated antibodies of the present invention can include antibodies in a bodily fluid (such as, but not limited to, serum), or antibodies that have been purified to varying degrees. Antibodies of the present invention can be polyclonal or monoclonal. Functional equivalents of such antibodies, such as antibody fragments and genetically-
5 engineered antibodies (including single chain antibodies or chimeric antibodies that can bind to more than one epitope) are also included in the present invention.

A preferred method to produce antibodies of the present invention includes (a) administering to an animal an effective amount of a protein, peptide or mimetope thereof of the present invention to produce the antibodies and (b) recovering the antibodies. In
10 another method, antibodies of the present invention are produced recombinantly using techniques as heretofore disclosed to produce flea SPI proteins of the present invention. Antibodies raised against defined proteins or mimetopes can be advantageous because such antibodies are not substantially contaminated with antibodies against other substances that might otherwise cause interference in a diagnostic assay or side effects if
15 used in a therapeutic composition.

Antibodies of the present invention have a variety of potential uses that are within the scope of the present invention. For example, such antibodies can be used (a) as therapeutic compounds to passively immunize an animal in order to protect the animal from hematophagous ectoparasites susceptible to treatment by such antibodies
20 and/or (b) as tools to screen expression libraries and/or to recover desired proteins of the present invention from a mixture of proteins and other contaminants. Furthermore, antibodies of the present invention can be used to target cytotoxic agents to hematophagous ectoparasite such as those disclosed herein in order to directly kill such hematophagous ectoparasites. Targeting can be accomplished by conjugating (i.e.,
25 stably joining) such antibodies to the cytotoxic agents using techniques known to those skilled in the art. Suitable cytotoxic agents are known to those skilled in the art.

One embodiment of the present invention is a therapeutic composition that, when administered to an animal in an effective manner, is capable of protecting that animal from infestation by hematophagous ectoparasites. Therapeutic compositions of the
30 present invention include at least one of the following protective compounds: an isolated flea SPI protein (including a peptide of a flea SPI protein capable of inhibiting serine

protease activity), a mimetope of a flea SPI protein, an isolated SPI nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* SPI gene, an isolated antibody that selectively binds to a flea SPI protein, and inhibitors of flea SPI activity (including flea SPI protein substrate analogs, such as serine proteases or serine protease analogs). Preferred hematophagous ectoparasites to target are heretofore disclosed. Examples of protective compounds (e.g., proteins, mimetopes, nucleic acid molecules, antibodies, and inhibitors) are disclosed herein.

Suitable inhibitors of SPI activity are compounds that interact directly with a SPI protein active site, thereby inhibiting that SPI's activity, usually by binding to or otherwise interacting with or otherwise modifying the SPI's active site. SPI inhibitors can also interact with other regions of the SPI protein to inhibit SPI activity, for example, by allosteric interaction. Inhibitors of SPIs are usually relatively small compounds and as such differ from anti-SPI antibodies. Preferably, a SPI inhibitor of the present invention is identified by its ability to bind to, or otherwise interact with, a flea SPI protein, thereby inhibiting the activity of the flea SPI.

Inhibitors of a SPI can be used directly as compounds in compositions of the present invention to treat animals as long as such compounds are not harmful to host animals being treated. Inhibitors of a SPI protein can also be used to identify preferred types of flea SPI proteins to target using compositions of the present invention, for example by affinity chromatography. Preferred inhibitors of a SPI of the present invention include, but are not limited to, flea SPI substrate analogs, and other molecules that bind to a flea SPI (e.g., to an allosteric site) in such a manner that SPI activity of the flea SPI is inhibited. A SPI substrate analog refers to a compound that interacts with (e.g., binds to, associates with, modifies) the active site of a SPI protein. A preferred SPI substrate analog inhibits SPI activity. SPI substrate analogs can be of any inorganic or organic composition, and, as such, can be, but are not limited to, peptides, nucleic acids, and peptidomimetic compounds. SPI substrate analogs can be, but need not be, structurally similar to a SPI protein's natural substrate as long as they can interact with the active site of that SPI protein. SPI substrate analogs can be designed using computer-generated structures of SPI proteins of the present invention or computer

structures of SPI proteins' natural substrates. Substrate analogs can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides, peptidomimetic compounds, or other inorganic or organic molecules, and screening such samples by affinity chromatography techniques using the corresponding binding partner, (e.g., a flea SPI or anti-flea serine protease antibody). A preferred SPI substrate analog is a peptidomimetic compound (i.e., a compound that is structurally and/or functionally similar to a natural substrate of a SPI of the present invention, particularly to the region of the substrate that interacts with the SPI active site, but that inhibits SPI activity upon interacting with the SPI active site).

10 SPI peptides, mimetopes and substrate analogs, as well as other protective compounds, can be used directly as compounds in compositions of the present invention to treat animals as long as such compounds are not harmful to the animals being treated.

The present invention also includes a therapeutic composition comprising at least one flea SPI-based compound of the present invention in combination with at least one additional compound protective against hematophagous ectoparasite infestation.

15 Examples of such compounds are disclosed herein.

In one embodiment, a therapeutic composition of the present invention can be used to protect an animal from hematophagous ectoparasite infestation by administering such composition to a hematophagous ectoparasite, such as to a flea, in order to prevent infestation. Such administration could be orally or by developing transgenic vectors capable of producing at least one therapeutic composition of the present invention. In another embodiment, a hematophagous ectoparasite, such as a flea, can ingest therapeutic compositions, or products thereof, present in the blood of a host animal that has been administered a therapeutic composition of the present invention.

25 Compositions of the present invention can be administered to any animal susceptible to hematophagous ectoparasite infestation (i.e., a host animal), including warm-blooded animals. Preferred animals to treat include mammals and birds, with cats, dogs, humans, cattle, chinchillas, ferrets, goats, mice, minks, rabbits, raccoons, rats, sheep, squirrels, swine, chickens, ostriches, quail and turkeys as well as other furry animals, pets and/or economic food animals, being more preferred. Particularly preferred animals to protect are cats and dogs.

In accordance with the present invention, a host animal (i.e., an animal that is or is capable of being infested with a hematophagous ectoparasite) is treated by administering to the animal a therapeutic composition of the present invention in such a manner that the composition itself (e.g., an inhibitor of a SPI protein, a SPI synthesis suppressor (i.e., a compound that decreases the production of SPI in the hematophagous ectoparasite), an SPI mimetope, or an anti-hematophagous ectoparasite SPI antibody) or a product generated by the animal in response to administration of the composition (e.g., antibodies produced in response to a flea SPI protein or nucleic acid molecule vaccine, or conversion of an inactive inhibitor "prodrug" to an active inhibitor of a SPI protein) ultimately enters the hematophagous ectoparasite. A host animal is preferably treated in such a way that the compound or product thereof enters the blood stream of the animal. Hematophagous ectoparasites are then exposed to the composition or product when they feed from the animal. For example, flea SPI protein inhibitors administered to an animal are administered in such a way that the inhibitors enter the blood stream of the animal, where they can be taken up by feeding fleas. In another embodiment, when a host animal is administered a flea SPI protein or nucleic acid molecule vaccine, the treated animal mounts an immune response resulting in the production of antibodies against the SPI protein (i.e., anti-flea SPI antibodies) which circulate in the animal's blood stream and are taken up by hematophagous ectoparasites upon feeding. Blood taken up by hematophagous ectoparasites enters the hematophagous ectoparasites where compounds of the present invention, or products thereof, such as anti-flea SPI antibodies, flea SPI protein inhibitors, flea mimetopes and/or SPI synthesis suppressors, interact with, and reduce SPI protein activity in the hematophagous ectoparasite.

The present invention also includes the ability to reduce larval hematophagous ectoparasite infestation in that when hematophagous ectoparasites feed from a host animal that has been administered a therapeutic composition of the present invention, at least a portion of compounds of the present invention, or products thereof, in the blood taken up by the hematophagous ectoparasite are excreted by the hematophagous ectoparasite in feces, which is subsequently ingested by hematophagous ectoparasite larvae. In particular, it is of note that flea larvae obtain most, if not all, of their nutrition from flea feces.

In accordance with the present invention, reducing SPI protein activity in a hematophagous ectoparasite can lead to a number of outcomes that reduce hematophagous ectoparasite burden on treated animals and their surrounding environments. Such outcomes include, but are not limited to, (a) reducing the viability
5 of hematophagous ectoparasites that feed from the treated animal, (b) reducing the fecundity of female hematophagous ectoparasites that feed from the treated animal, (c) reducing the reproductive capacity of male hematophagous ectoparasites that feed from the treated animal, (d) reducing the viability of eggs laid by female hematophagous ectoparasites that feed from the treated animal, (e) altering the blood feeding behavior of
10 hematophagous ectoparasites that feed from the treated animal (e.g., hematophagous ectoparasites take up less volume per feeding or feed less frequently), (f) reducing the viability of hematophagous ectoparasite larvae (e.g., by decreasing feeding behavior, inhibiting growth, inhibiting (e.g., slowing or blocking) molting, and/or otherwise inhibiting maturation to adults).

15 Therapeutic compositions of the present invention can be formulated in an excipient that the animal to be treated can tolerate. Examples of such excipients include water, saline, Ringer's solution, dextrose solution, Hank's solution, and other aqueous physiologically balanced salt solutions. Nonaqueous vehicles, such as fixed oils, sesame oil, ethyl oleate, or triglycerides may also be used. Other useful formulations include
20 suspensions containing viscosity enhancing agents, such as sodium carboxymethylcellulose, sorbitol, or dextran. Excipients can also contain minor amounts of additives, such as substances that enhance isotonicity and chemical stability. Examples of buffers include phosphate buffer, bicarbonate buffer and Tris buffer, while examples of preservatives include thimerosal, — or o-cresol, formalin and benzyl
25 alcohol. Standard formulations can either be liquid injectables or solids which can be taken up in a suitable liquid as a suspension or solution for injection. Thus, in a non-liquid formulation, the excipient can comprise dextrose, human serum albumin, preservatives, etc., to which sterile water or saline can be added prior to administration.

In one embodiment of the present invention, a therapeutic composition can
30 include an adjuvant. Adjuvants are agents that are capable of enhancing the immune response of an animal to a specific antigen. Suitable adjuvants include, but are not

limited to, cytokines, chemokines, and compounds that induce the production of cytokines and chemokines (e.g., granulocyte macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), colony stimulating factor (CSF), erythropoietin (EPO), interleukin 2 (IL-2), interleukin-3 (IL-3), interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 6 (IL-6), interleukin 7 (IL-7), interleukin 8 (IL-8), interleukin 10 (IL-10), interleukin 12 (IL-12), interferon gamma, interferon gamma inducing factor I (IGIF), transforming growth factor beta, RANTES (regulated upon activation, normal T cell expressed and presumably secreted), macrophage inflammatory proteins (e.g., MIP-1 alpha and MIP-1 beta), and Leishmania elongation initiating factor (LEIF); bacterial components (e.g., endotoxins, in particular superantigens, exotoxins and cell wall components); aluminum-based salts; calcium-based salts; silica; polynucleotides; toxoids; serum proteins, viral coat proteins; block copolymer adjuvants (e.g., Hunter's Titermax™ adjuvant (Vaxcel™, Inc. Norcross, GA), Ribi adjuvants (Ribi ImmunoChem Research, Inc., Hamilton, MT); and saponins and their derivatives (e.g., Quil A (Superfos Biosector A/S, Denmark). Protein adjuvants of the present invention can be delivered in the form of the protein themselves or of nucleic acid molecules encoding such proteins using the methods described herein.

In one embodiment of the present invention, a therapeutic composition can include a carrier. Carriers include compounds that increase the half-life of a therapeutic composition in the treated animal. Suitable carriers include, but are not limited to, polymeric controlled release vehicles, biodegradable implants, liposomes, bacteria, viruses, other cells, oils, esters, and glycols.

One embodiment of the present invention is a controlled release formulation that is capable of slowly releasing a composition of the present invention into an animal. As used herein, a controlled release formulation comprises a composition of the present invention in a controlled release vehicle. Suitable controlled release vehicles include, but are not limited to, biocompatible polymers, other polymeric matrices, capsules, microcapsules, microparticles, bolus preparations, osmotic pumps, diffusion devices, liposomes, lipospheres, and transdermal delivery systems. Other controlled release formulations of the present invention include liquids that, upon administration to an

animal, form a solid or a gel *in situ*. Preferred controlled release formulations are biodegradable (i.e., bioerodible).

A preferred controlled release formulation of the present invention is capable of releasing a composition of the present invention into the blood of an animal at a constant
5 rate sufficient to attain therapeutic dose levels of the composition to protect an animal from hematophagous ectoparasite infestation. The therapeutic composition is preferably released over a period of time ranging from about 1 to about 12 months. A preferred controlled release formulation of the present invention is capable of effecting a treatment preferably for at least about 1 month, more preferably for at least about 3 months, even
10 more preferably for at least about 6 months, even more preferably for at least about 9 months, and even more preferably for at least about 12 months.

Acceptable protocols to administer therapeutic compositions of the present invention in an effective manner include individual dose size, number of doses, frequency of dose administration, and mode of administration. Determination of such
15 protocols can be accomplished by those skilled in the art. A suitable single dose is a dose that is capable of protecting an animal from disease when administered one or more times over a suitable time period. For example, a preferred single dose of a protein, mimotope or antibody therapeutic composition is from about 1 microgram (μg) to about 10 milligrams (mg) of the therapeutic composition per kilogram body weight of the
20 animal. Booster vaccinations can be administered from about 2 weeks to several years after the original administration. Booster administrations preferably are administered when the immune response of the animal becomes insufficient to protect the animal from disease. A preferred administration schedule is one in which from about 10 μg to about 1 mg of the therapeutic composition per kg body weight of the animal is
25 administered from about one to about two times over a time period of from about 2 weeks to about 12 months. Modes of administration can include, but are not limited to, subcutaneous, intradermal, intravenous, intranasal, oral, transdermal, intraocular and intramuscular routes.

According to one embodiment, a nucleic acid molecule of the present invention
30 can be administered to an animal in a fashion to enable expression of that nucleic acid molecule into a protective protein or protective RNA (e.g., antisense RNA, ribozyme,

triple helix forms or RNA drug) in the animal. Nucleic acid molecules can be delivered to an animal in a variety of methods including, but not limited to, (a) administering a naked (i.e., not packaged in a viral coat or cellular membrane) nucleic acid vaccine (e.g., as naked DNA or RNA molecules, such as is taught, for example in Wolff et al., 1990, *Science* 247, 1465-1468) or (b) administering a nucleic acid molecule packaged as a recombinant virus vaccine or as a recombinant cell vaccine (i.e., the nucleic acid molecule is delivered by a viral or cellular vehicle).

A naked nucleic acid vaccine of the present invention includes a nucleic acid molecule of the present invention and preferably includes a recombinant molecule of the present invention that preferably is replication, or otherwise amplification, competent. A naked nucleic acid vaccine of the present invention can comprise one or more nucleic acid molecules of the present invention in the form of, for example, a bicistronic recombinant molecule having, for example one or more internal ribosome entry sites. Preferred naked nucleic acid vaccines include at least a portion of a viral genome (i.e., a viral vector). Preferred viral vectors include those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, and retroviruses, with those based on alphaviruses (such as Sindbis or Semliki virus), species-specific herpesviruses and species-specific poxviruses being particularly preferred. Any suitable transcription control sequence can be used, including those disclosed as suitable for protein production. Particularly preferred transcription control sequence include cytomegalovirus intermediate early (preferably in conjunction with Intron-A), Rous Sarcoma Virus long terminal repeat, and tissue-specific transcription control sequences, as well as transcription control sequences endogenous to viral vectors if viral vectors are used. The incorporation of "strong" poly(A) sequences are also preferred.

Naked nucleic acid vaccines of the present invention can be administered in a variety of ways, with intramuscular, subcutaneous, intradermal, transdermal, intranasal and oral routes of administration being preferred. A preferred single dose of a naked nucleic acid vaccines ranges from about 1 nanogram (ng) to about 100 µg, depending on the route of administration and/or method of delivery, as can be determined by those skilled in the art. Suitable delivery methods include, for example, by injection, as drops, aerosolized and/or topically. Naked DNA of the present invention can be contained in

an aqueous excipient (e.g., phosphate buffered saline) alone or a carrier (e.g., lipid-based vehicles).

A recombinant virus vaccine of the present invention includes a recombinant molecule of the present invention that is packaged in a viral coat and that can be expressed in an animal after administration. Preferably, the recombinant molecule is packaging-deficient and/or encodes an attenuated virus. A number of recombinant viruses can be used, including, but not limited to, those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, and retroviruses. Preferred recombinant virus vaccines are those based on alphaviruses (such as Sindbis virus), raccoon poxviruses, species-specific herpesviruses and species-specific poxviruses. An example of methods to produce and use alphavirus recombinant virus vaccines is disclosed in PCT Publication No. WO 94/17813, by Xiong et al., published August 18, 1994, which is incorporated by reference herein in its entirety.

When administered to an animal, a recombinant virus vaccine of the present invention infects cells within the immunized animal and directs the production of a protective protein or RNA nucleic acid molecule that is capable of protecting the animal from hematophagous ectoparasite infestation. For example, a recombinant virus vaccine comprising a flea SPI nucleic acid molecule of the present invention is administered according to a protocol that results in the animal producing a sufficient immune response to protect itself from hematophagous ectoparasite infestation. A preferred single dose of a recombinant virus vaccine of the present invention is from about 1×10^4 to about 1×10^7 virus plaque forming units (pfu) per kilogram body weight of the animal. Administration protocols are similar to those described herein for protein-based vaccines, with subcutaneous, intramuscular, intranasal and oral administration routes being preferred.

A recombinant cell vaccine of the present invention includes recombinant cells of the present invention that express at least one protein of the present invention. Preferred recombinant cells for this embodiment include *Salmonella*, *E. coli*, *Listeria*, *Mycobacterium*, *S. frugiperda*, yeast, (including *Saccharomyces cerevisiae*), BHK, CV-1, myoblast G8, COS (e.g., COS-7), Vero, MDCK and CRFK recombinant cells. Recombinant cell vaccines of the present invention can be administered in a variety of

ways but have the advantage that they can be administered orally, preferably at doses ranging from about 10^8 to about 10^{12} cells per kilogram body weight. Administration protocols are similar to those described herein for protein-based vaccines. Recombinant cell vaccines can comprise whole cells, cells stripped of cell walls or cell lysates.

5 The efficacy of a therapeutic composition of the present invention to protect an animal from hematophagous ectoparasite infestation can be tested in a variety of ways including, but not limited to, detection of anti-flea SPI antibodies (using, for example, proteins or mimetopes of the present invention), detection of cellular immunity within the treated animal, or challenge of the treated animal with hematophagous ectoparasites
10 to determine whether, for example, the feeding, fecundity or viability of the hematophagous ectoparasites feeding from the treated animal is disrupted. Challenge studies can include attachment of chambers containing fleas onto the skin of the treated animal. In one embodiment, therapeutic compositions can be tested in animal models such as mice. Such techniques are known to those skilled in the art.

15 One preferred embodiment of the present invention is the use of flea SPI proteins, mimetopes, nucleic acid molecules, antibodies and inhibitory compounds of the present invention, to protect an animal from hematophagous ectoparasite infestation. Preferred protective compounds of the present invention include, but are not limited to, an isolated flea SPI protein or a mimetope thereof, an isolated SPI nucleic acid molecule
20 that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* SPI gene, an isolated antibody that selectively binds to a flea SPI and/or an inhibitor of flea SPI activity (such as, but not limited to, an SPI substrate analog). Additional protection may be obtained by administering additional protective compounds, including other proteins, nucleic acid molecules, antibodies and inhibitory compounds, as disclosed
25 herein.

 An inhibitor of SPI activity can be identified using flea SPI proteins of the present invention. One embodiment of the present invention is a method to identify a compound capable of inhibiting SPI activity of a flea. Such a method includes the steps of (a) contacting (e.g., combining, mixing) an isolated flea SPI protein, preferably a *C.
30 felis* SPI protein, with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has SPI activity, and (b) determining if the

putative inhibitory compound inhibits the SPI activity. Putative inhibitory compounds to screen include small organic molecules, antibodies (including mimetopes thereof) and substrate analogs. Methods to determine SPI activity are known to those skilled in the art.

- 5 The present invention also includes a test kit to identify a compound capable of inhibiting SPI activity of a flea. Such a test kit includes an isolated flea SPI protein, preferably a *C. felis* SPI protein, having SPI activity and a means for determining the extent of inhibition of SPI activity in the presence of (i.e., effected by) a putative inhibitory compound. Such compounds are also screened to identify those that are
- 10 substantially not toxic in host animals.

SPI inhibitors isolated by such a method, and/or test kit, can be used to inhibit any SPI protein that is susceptible to such an inhibitor. Preferred SPI enzymes proteins to inhibit are those produced by fleas. A particularly preferred inhibitor of a SPI protein of the present invention is capable of protecting an animal from flea infestation.

- 15 Effective amounts and dosing regimens can be determined using techniques known to those skilled in the art.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention.

EXAMPLES

- 20 It is to be noted that the Examples include a number of molecular biology, microbiology, immunology and biochemistry techniques considered to be known to those skilled in the art. Disclosure of such techniques can be found, for example, in Sambrook et al., *ibid.*, and related references.

Example 1

- 25 This example describes the isolation of a protein fraction from flea prepupal larvae that was obtained by monitoring for carboxylesterase activity, which surprisingly, also contained flea serine protease inhibitor molecule epitopes of the present invention, discovered as described in Examples 2, 3 and 4 below.

- A prepupal larval protein pool enriched for carboxylesterase activity was isolated
- 30 as follows. About 17,000 bovine blood-fed prepupal larvae were collected and the larvae were homogenized in gut dissection buffer (50 mM Tris pH 8.0, 100 mM CaCl₂)

by sonication in a disposable 50 ml conical centrifuge tube. Sonication entailed 4 bursts of 20 seconds each at a setting of 4 with a probe sonicator using, for example, a model W-380 Sonicator (available from Heat Systems-Ultrasonics, Inc., Farmingdale, NY).

The sonicate was clarified by centrifugation at 4000 rpm for 30 min. in a swinging
5 bucket centrifuge; the supernatant was collected and centrifuged at 18,000 rpm for 30 min in a Sorvall SS-34 rotor (available from DuPont, Wilmington, DE). The supernatant was recovered, and NaCl was added to a final concentration of 400 mM.

Serine proteases were removed from the supernatant using the following method. The supernatant was loaded onto a 5-ml column comprising p-aminobenzamidine cross-
10 linked to Sepharose beads (available from Sigma Chemical Company, St. Louis, MO), previously equilibrated in benzamidine column buffer (50 mM Tris 8.0, 100 mM CaCl₂, 400 mM NaCl) and incubated overnight at 4°C. Unbound protein was slowly washed off and collected from the column with benzamidine column buffer until no protein was detectable by a Bradford Assay (available from Bio-Rad Laboratories, Hercules, CA). A
15 total of about 43 ml was collected. The proteins in this pool were fractionated by precipitation in increasing percent saturation levels of ammonium sulfate.

The ammonium sulfate-precipitated protein fractions, as well as all subsequent protein fractions described in this example, were assayed for carboxylesterase activity by the following method. Samples of about 5 µl of each fraction were added to separate
20 wells of a flat-bottomed microtiter plate (available from Becton Dickinson, Lincoln Park, NJ). A control well was prepared by adding about 5 µl of Tris buffer to an empty well of the plate. About 95 µl of 25 mM Tris-HCl (pH 8.0) was then added to each sample to increase the volume in each well to about 100 µl. About 100 µl of 0.25 mM α-naphthyl acetate (available from Sigma) dissolved in 25 mM Tris-HCl (pH 8.0) was
25 then added to each well. The plate was then incubated for about 15 min. at 37°C. Following the incubation, about 40 µl of 0.3% Fast Blue salt BN (tetrazotized o-dianisidine; available from Sigma), dissolved in 3.3% SDS in water was added to each well, giving a colorimetric reaction. Absorbance levels were measured using a model 7500 Microplate Reader (available from Cambridge Technology, Inc., Watertown, MA)
30 set to 590 nm. Following subtraction of background absorbance, the resulting values gave a relative measure of carboxylesterase activity. Carboxylesterase activity was found

in two of the ammonium sulfate-precipitated fractions. The first, which precipitated between about 0 and 60% ammonium sulfate saturation, was kept as a pool, and the second, which precipitated between about 60 and 80% ammonium sulfate saturation, was kept separately as a pool. Since the latter pool appeared to have higher activity at
5 this point, the pools were treated separately until just prior to the final HPLC step described below, but at that point they were combined.

The two ammonium sulfate-precipitated protein pools were then subjected to cation exchange chromatography, performed as follows. Each protein pool was dialyzed two times against about 500 ml of 20 mM 2-(N-morpholino) ethanesulfonic acid (MES)
10 buffer, pH 6, containing 10 mM NaCl and was then applied to a 40-ml chromatography column containing 10 ml of S-Sepharose Fast Flow cation exchange resin (available from Pharmacia Biochemicals, Piscataway, NJ), previously equilibrated with MES buffer. Each column was rocked overnight at 4°C to facilitate protein binding, and was then drained and washed with more MES buffer to remove all unbound protein in about
15 40 ml total volume. Following elution of the bound proteins, the bound and unbound protein fractions were tested for carboxylesterase activity as described above. Activity was found to reside in the unbound protein fractions from each column, which were then concentrated to about 5 ml using Centriprep® 30 centrifugal concentrators (available from Amicon, Beverly, MA).

20 The two concentrated protein pools were then subjected to anion exchange chromatography, performed as follows. Each pool was adjusted to about pH 7 by the addition of a small amount of 500 mM Tris buffer, pH 8, and was then applied, in about 1 to 1.5 ml aliquots, to a 4.5 mm x 50 mm Poros 10 HQ anion exchange chromatography column (available from PerSeptive Biosystems, Cambridge, MA) equilibrated in 25 mM
25 Tris, pH 6.8 (loading buffer). For each aliquot, the column was washed with the loading buffer, and bound proteins were eluted with a linear gradient of 0 to 1 M NaCl in 25 mM Tris buffer, pH 6.8. All column fractions were tested for carboxylesterase activity as described above. For each aliquot run on the column, the activity peak eluted in fractions 31-34, and at this point in the isolation, the activity levels appeared to be
30 equivalent in both of the original ammonium sulfate-fractionated pools. Therefore, all

column fractions containing carboxylesterase activity were combined into one pool. This pool was concentrated and diafiltered into about 1 ml of Tris-buffered saline (TBS).

The pooled protein preparation was then loaded onto a C1 reverse phase HPLC column (available from TosoHaas, Montgomeryville, PA), previously equilibrated with
5 19% acetonitrile containing 0.05% trifluoroacetic acid (TFA). The column was washed with the equilibration buffer to remove unbound proteins, and bound proteins were eluted from the column by a linear gradient from 19% acetonitrile containing 0.05% TFA to 95% acetonitrile containing 0.05% TFA. The column fractions were tested for carboxylesterase activity as described above, and the activity peak eluted in fractions 27-
10 32. These fractions were combined, concentrated to near dryness using a Speed-Vac™ concentrator (available from Savant Instruments, Molbrook, NY), and resuspended in phosphate-buffered saline (PBS) to a concentration of about 0.2mg/ml. This isolated protein fraction is referred to herein as flea prepupal carboxylesterase fraction-1. Upon analysis by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and silver staining,
15 flea prepupal carboxylesterase fraction-1 appeared to contain, in addition to the recognized carboxylesterase bands migrating at about 60 kD, a strong protein band migrating at about 40 kD.

Example 2

This example describes the generation of polyclonal rabbit antiserum to flea
20 prepupal carboxylesterase fraction-1.

Antibodies against flea prepupal carboxylesterase fraction-1 (the preparation of which is described in Example 1) were generated as follows. A rabbit was initially immunized subcutaneously and intradermally at multiple sites with a total of approximately 50 µg of flea prepupal carboxylesterase fraction-1 emulsified in Complete
25 Freund's Adjuvant. On days 16 and 37 after the initial immunization, the rabbit was boosted intramuscularly with a total of approximately 50 µg of flea prepupal carboxylesterase fraction-1 emulsified in Incomplete Freund's Adjuvant. The rabbit was bled on days 9, 29 and 50 after the initial immunization. Sera from the latter two bleeds, putatively containing antibodies to flea prepupal carboxylesterases, were used separately
30 for immunoscreening experiments, as described in Example 3 below.

Example 3

This example describes the isolation, by immunoscreening, of nucleic acid molecules encoding flea serine protease inhibitor proteins of the present invention.

Surprisingly, six flea serine protease inhibitor nucleic acid molecules were

5 isolated by their ability to encode proteins that selectively bound to at least one component of the immune serum collected from a rabbit immunized with flea prepupal carboxylesterase fraction-1, using the following method. A flea prepupal cDNA library was produced as follows. Total RNA was extracted from approximately 3,653 prepupal larvae using an acid-guanidinium-phenol-chloroform method similar to that described by

10 Chomczynski et al., 1987, *Anal. Biochem.* 162, 156-159. Poly A+ selected RNA was separated from the total RNA preparation by oligo-dT cellulose chromatography using Poly(A)Quick® mRNA isolation kits (available from Stratagene Cloning Systems, La Jolla, CA), according to the method recommended by the manufacturer. A prepupal cDNA expression library was constructed in lambda Uni-ZAP™XR vector (available

15 from Stratagene), using Stratagene's ZAP-cDNA Synthesis Kit® protocol. About 6.72 µg of prepupal poly A+ RNA was used to produce the prepupal library. The resultant prepupal library was amplified to a titer of about 3.5×10^{10} pfu/ml with about 97% recombinants.

Using a modification of the protocol described in the picoBlue immunoscreening

20 kit (available from Stratagene), the pre-pupal cDNA expression library was screened with the flea prepupal carboxylesterase fraction-1 immune rabbit serum, generated as described in Example 2. The protocol was modified in that the secondary peroxidase-conjugated antibody was detected with a chromogen substrate consisting of DAB (3,3' diaminobenzidine) plus cobalt (Sigma Fast, available from Sigma) following the

25 manufacturer's instructions, except that tablets were dissolved in water at one half the recommended final concentration. Plaque lift membranes were placed in the substrate solution for about 2 minutes, rinsed in water, and then dried at room temperature. Immunoscreening of duplicate plaque lifts of the cDNA library with the same immune rabbit serum identified six clones containing flea nucleic acid molecules nfSPI1₁₅₈₄,

30 nfSPI2₁₃₅₈, nfSPI3₁₈₃₈, nfSPI4₁₄₁₄, nfSPI5₁₄₉₂, and nfSPI6₁₄₅₄, respectively. Plaque purified clones including the flea nucleic acid molecules were converted into double

stranded recombinant molecules, herein denoted as p β gal-nfSPI1₁₅₈₄, p β gal-nfSPI2₁₃₅₈, p β gal-nfSPI3₁₈₃₈, p β gal-nfSPI4₁₄₁₄, p β gal-nfSPI5₁₄₉₂, and p β gal-nfSPI6₁₄₅₄, using ExAssist[™] helper phage and SOLR[™] *E. coli* according to the *in vivo* excision protocol described in the Zap-cDNA Synthesis Kit (available from Stratagene). Double-stranded
5 plasmid DNA was prepared using an alkaline lysis protocol, such as that described in Sambrook et al., *ibid*.

Example 4

This example describes the sequencing of several flea serine protease inhibitor nucleic acid molecules of the present invention.

10 The plasmids containing flea nfSPI1₁₅₈₄, nfSPI2₁₃₅₈, nfSPI3₁₈₃₈, nfSPI4₁₄₁₄, nfSPI5₁₄₉₂, and nfSPI6₁₄₅₄ were sequenced by the Sanger dideoxy chain termination method, using the PRISM[™] Ready Dye Terminator Cycle Sequencing Kit with AmpliTaq[®] DNA Polymerase, FS (available from the Perkin-Elmer Corporation, Norwalk, CT). PCR extensions were done in the GeneAmp[™] PCR System 9600
15 (available from Perkin-Elmer). Excess dye terminators were removed from extension products using the Centriflex[™] Gel Filtration Cartridge (available from Advanced Genetics Technologies Corporation, Gaithersburg, MD) following their standard protocol. Samples were resuspended according to ABI protocols and were and run on a Perkin-Elmer ABI PRISM[™] 377 Automated DNA Sequencer. DNA sequence analyses,
20 including the compilation of sequences and the determination of open reading frames, were performed using either the DNAsis[™] program (available from Hitachi Software, San Bruno, CA) or the MacVector[™] program (available from the Eastman Kodak Company, New Haven, CT). Protein sequence analyses, including the determination of molecular weights and isoelectric points (pI) were performed using the MacVector[™]
25 program.

A. An about 1584-nucleotide consensus sequence of the entire flea nfSPI1₁₅₈₄ DNA fragment was determined; the sequences of the two complementary strands are presented as SEQ ID NO:1 (the coding strand) and SEQ ID NO:3 (the complementary strand). The flea nfSPI1₁₅₈₄ sequence contains a full length coding region. The apparent
30 start and stop codons span nucleotides from about 136 through about 138 and from about 1327 through about 1329, respectively, of SEQ ID NO:1. A putative

polyadenylation signal (5' AATAAA 3') is located in a region spanning from about nucleotide 1533 through about 1538 of SEQ ID NO:1.

Translation of SEQ ID NO:1 yields a protein of about 397 amino acids, denoted PfSPII₃₉₇, the amino acid sequence of which is presented in SEQ ID NO:2. The nucleic acid molecule consisting of the coding region encoding PfSPII₃₉₇ is referred to herein as nfSPII₁₁₉₁, the nucleic acid sequence of which is represented in SEQ ID NO:4 (the coding strand) and SEQ ID NO:5 (the complementary strand). The amino acid sequence of flea PfSPII₃₉₇ (i.e., SEQ ID NO:2) predicts that PfSPII₃₉₇ has an estimated molecular weight of about 44.4 kD and an estimated pI of about 4.97. Analysis of SEQ ID NO:2 suggests the presence of a signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 21. The proposed mature protein, denoted herein as PfSPII₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:6. The amino acid sequence of flea PfSPII₃₇₆ (i.e. SEQ ID NO:6) predicts that PfSPII₃₇₆ has an estimated molecular weight of about 42.1 kD, an estimated pI of about 4.90, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

Homology searches of the non-redundant protein and nucleotide sequence databases were performed through the National Center for Biotechnology Information using the BLAST network. The protein database includes SwissProt +PIR + SPUpdate + Genpept + GPUUpdate. The nucleotide database includes GenBank + EMBL + DDBJ + PDB. The protein search was performed using SEQ ID NO:2, which showed significant homology to certain serine protease inhibitor proteins. The highest scoring match of the homology search at the amino acid level was GenBank accession number 1378131: *Manduca sexta*, which was about 36% identical with SEQ ID NO:2. At the nucleotide level, the search was performed using SEQ ID NO:4, which was most similar to accession number L20792, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 2) of *Manduca sexta*, being about 55% identical.

B. An about 1358-nucleotide consensus sequence of the entire flea nfSPII₁₃₅₈ DNA fragment was determined; the sequences of the two complementary strands are presented as SEQ ID NO:7 (the coding strand) and SEQ ID NO:9 (the complementary strand). The flea nfSPII₁₃₅₈ sequence contains a partial coding region, which is truncated

at the 5' end. The first in-frame codon spans nucleotides from 2 through 4 and the stop codon spans nucleotides from 1199 through 1201 of SEQ ID NO:7.

Translation of SEQ ID NO:7 yields a protein of about 399 amino acids, denoted PfSPI2₃₉₉, the amino acid sequence of which is presented in SEQ ID NO:8. The nucleic acid molecule consisting of the coding region encoding PfSPI2₃₉₉ is referred to herein as nfSPI2₁₁₉₇, the nucleic acid sequence of which is represented in SEQ ID NO:10 (the coding strand) and SEQ ID NO:11 (the complementary strand). Analysis of SEQ ID NO:8 suggests the presence of a partial signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 23. The proposed mature protein, denoted herein as PfSPI2₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:12. The amino acid sequence of flea PfSPI1₃₇₆ (i.e. SEQ ID NO:12) predicts that PfSPI2₃₇₆ has an estimated molecular weight of about 42.1 kD, an estimated pI of about 4.87, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

BLAST searches were performed as described in Section A. The protein search was performed using SEQ ID NO:8, which showed significant homology to certain serine protease inhibitor proteins. The highest scoring match of the homology search at the amino acid level was GenBank accession number 1345616: *Homo sapiens*, which was about 36% identical with SEQ ID NO:8. At the nucleotide level, the search was performed using SEQ ID NO:10, which was most similar to accession number L20790, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 1) of *Manduca sexta*, being about 43% identical.

C. An about 1838-nucleotide consensus sequence of the entire flea nfSPI3₁₈₃₈ DNA fragment was determined; the sequences of the two complementary strands are presented as SEQ ID NO:13 (the coding strand) and SEQ ID NO:15 (the complementary strand). The flea nfSPI3₁₈₃₈ sequence contains a full-length coding region. The apparent start and stop codons span nucleotides from about 306 through about 308 and from about 1566 through about 1568, respectively, of SEQ ID NO:13. A putative polyadenylation signal (5' AATAAA 3') is located in a region spanning from about nucleotide 1803 through about 1808 of SEQ ID NO:13.

Translation of SEQ ID NO:13 yields a protein of about 420 amino acids, denoted PfSPI3₄₂₀, the amino acid sequence of which is presented in SEQ ID NO:14. The nucleic acid molecule consisting of the coding region encoding PfSPI3₄₂₀ is referred to herein as nfSPI3₁₂₆₀, the nucleic acid sequence of which is represented in SEQ ID NO:16 (the coding strand) and SEQ ID NO:17 (the complementary strand). The amino acid sequence of flea PfSPI3₄₂₀ (i.e., SEQ ID NO:14) predicts that PfSPI3₄₂₀ has an estimated molecular weight of about 47.1 kD and an estimated pI of about 4.72. Analysis of SEQ ID NO:14 suggests the presence of a signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 30. The proposed mature protein, denoted herein as PfSPI3₃₉₀, contains about 390 amino acids which is represented herein as SEQ ID NO:18. The amino acid sequence of flea PfSPI3₃₉₀ (i.e., SEQ ID NO:18) predicts that PfSPI3₃₉₀ has an estimated molecular weight of about 43.7 kD, an estimated pI of about 4.63, and two predicted asparagine-linked glycosylation sites extending from about amino acid 252 to about amino acid 254 and from about amino acid 369 to about amino acid 371.

BLAST searches were performed as described in Section A. The protein search was performed using SEQ ID NO:14, which showed significant homology to certain serine protease inhibitor proteins. The highest scoring match of the homology search at the amino acid level was GenBank accession number 1345616: *Homo sapiens*, which was about 35% identical with SEQ ID NO:14. At the nucleotide level, the search was performed using SEQ ID NO:16, which was most similar to accession number L20792, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 2) of *Manduca sexta*, being about 52% identical.

D. An about 1414-nucleotide consensus sequence of the entire flea nfSPI4₁₄₁₄ DNA fragment was determined; the sequences of the two complementary strands are presented as SEQ ID NO:19 (the coding strand) and SEQ ID NO:21 (the complementary strand). The flea nfSPI4₁₄₁₄ sequence contains a partial coding region, truncated at the 5' end. The first in-frame codon spans nucleotides from 2 through 4 and the stop codon spans nucleotides from 1181 through 1183 of SEQ ID NO:19. A putative polyadenylation signal (5' AATAAA 3') is located in a region spanning from nucleotide 1179 through 1184 of SEQ ID NO:19.

Translation of SEQ ID NO:19 yields a protein of about 393 amino acids, denoted PfSPI4₃₉₃, the amino acid sequence of which is presented in SEQ ID NO:20. The nucleic acid molecule consisting of the coding region encoding PfSPI4₃₉₃ is referred to herein as nfSPI4₁₁₇₉, the nucleic acid sequence of which is represented in SEQ ID NO:22 (the coding strand) and SEQ ID NO:23 (the complementary strand). Analysis of SEQ ID NO:20 suggests the presence of a partial signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 17. The proposed mature protein, denoted herein as PfSPI4₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:24. The amino acid sequence of flea PfSPI4₃₇₆ (i.e. SEQ ID NO:24) predicts that PfSPI4₃₇₆ has an estimated molecular weight of about 42.2 kD, an estimated pI of about 5.31, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

BLAST searches were performed as described in Section A. The protein search was performed using SEQ ID NO:20, which showed significant homology to certain serine protease inhibitor proteins. The highest scoring match of the homology search at the amino acid level was GenBank accession number 1345616: *Homo sapiens*, which was about 38% identical with SEQ ID NO:20. At the nucleotide level, the search was performed using SEQ ID NO:22, which was most similar to accession number L20793, a putative serine proteinase inhibitor gene (serpin 1, exon 9 unknown copy number) of *Manduca sexta*, being about 55% identical.

E. An about 1492-nucleotide consensus sequence of the entire flea nfSPI5₁₄₉₂ DNA fragment was determined; the sequences of the two complementary strands are presented as SEQ ID NO:25 (the coding strand) and SEQ ID NO:27 (the complementary strand). The flea nfSPI5₁₄₉₂ sequence contains a partial coding region, truncated at the 5' end. The first in-frame codon spans nucleotides from 3 through 5 and the stop codon spans nucleotides from 1197 through 1199 of SEQ ID NO:25. A putative polyadenylation signal (5' AATAAA 3') is located in a region spanning from nucleotide 1416 through 1421 of SEQ ID NO:25.

Translation of SEQ ID NO:25 yields a protein of about 398 amino acids, denoted PfSPI5₃₉₈, the amino acid sequence of which is presented in SEQ ID NO:26. The nucleic acid molecule consisting of the coding region encoding PfSPI5₃₉₈ is referred to

herein as nfSPI5₁₁₉₄, the nucleic acid sequence of which is represented in SEQ ID NO:28 (the coding strand) and SEQ ID NO:29 (the complementary strand). Analysis of SEQ ID NO:26 suggests the presence of a partial signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 22. The proposed
5 mature protein, denoted herein as PfSPI5₃₇₆, contains about 376 amino acids which is represented herein as SEQ ID NO:30. The amino acid sequence of flea PfSPI5₃₇₆ (i.e. SEQ ID NO:30) predicts that PfSPI5₃₇₆ has an estimated molecular weight of about 42.3 kD, an estimated pI of about 5.31 and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

10 BLAST searches were performed as described in Section A. The protein search was performed using SEQ ID NO:26, which showed significant homology to certain serine protease inhibitor proteins. The highest scoring match of the homology search at the amino acid level was GenBank accession number 1345616: *Homo sapiens*, which was about 38% identical with SEQ ID NO:26. At the nucleotide level, the search was
15 performed using SEQ ID NO:28, which was most similar to accession number L20790, a putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 1) of *Manduca sexta*, being about 45% identical.

F. An about 1454-nucleotide consensus sequence of the entire flea nfSPI6₁₄₅₄ DNA fragment was determined; the sequences of the two complementary strands are
20 presented as SEQ ID NO:31 (the coding strand) and SEQ ID NO:33 (the complementary strand). The flea nfSPI6₁₄₅₄ sequence contains a full length coding region. The apparent start and stop codons span nucleotides from about 20 through about 22 and from about 1211 through about 1213, respectively, of SEQ ID NO:31. A putative polyadenylation signal (5' AATAAA 3') is located in a region spanning from about nucleotide 1419
25 through about 1424 of SEQ ID NO:31.

Translation of SEQ ID NO:31 yields a protein of about 397 amino acids, denoted PfSPI6₃₉₇, the amino acid sequence of which is presented in SEQ ID NO:32. The nucleic acid molecule consisting of the coding region encoding PfSPI6₃₉₇ is referred to herein as nfSPI6₁₁₉₁, the nucleic acid sequence of which is represented in SEQ ID NO:34
30 (the coding strand) and SEQ ID NO:35 (the complementary strand). The amino acid sequence of flea PfSPI6₃₉₇ (i.e., SEQ ID NO:32) predicts that PfSPI6₃₉₇ has an estimated

molecular weight of about 44.4 kD and an estimated pI of about 4.90. Analysis of SEQ ID NO:32 suggests the presence of a signal peptide encoded by a stretch of amino acids spanning from about amino acid 1 through about amino acid 21. The proposed mature protein, denoted herein as PfSPI6₃₇₆, contains about 376 amino acids which is

5 represented herein as SEQ ID NO:36. The amino acid sequence of flea PfSPI6₃₇₆ (i.e. SEQ ID NO:36) predicts that PfSPI6₃₇₆ has an estimated molecular weight of about 42.1 kD, an estimated pI of about 4.84, and a predicted asparagine-linked glycosylation site extending from about amino acid 252 to about amino acid 254.

BLAST searches were performed as described in Section A. The protein search
10 was performed using SEQ ID NO:32, which showed significant homology to certain serine protease inhibitor proteins. The highest scoring match of the homology search at the amino acid level was GenBank accession number 1378131: *Manduca sexta*, which was about 36% identical with SEQ ID NO:32. At the nucleotide level, the search was performed using SEQ ID NO:34, which was most similar to accession number L20792, a
15 putative serine proteinase inhibitor gene (serpin 1, exon 9 copy 2) of *Manduca sexta*, being about 55% identical.

Example 5

This example discloses the production of a several recombinant cells of the present invention.

20 A. Recombinant molecule p λ P_R-nfSPI2₁₁₃₉, containing a portion of a flea serine protease inhibitor molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1185-nucleotide DNA fragment containing nucleotides spanning from about 26 through about 1202 of SEQ ID
25 NO:7, denoted herein as nfSPI2₁₁₈₅, was PCR amplified from nucleic acid molecule nfSPI2₁₃₅₈, produced as described in Example 3, using sense primer JPI5, having the nucleic acid sequence 5' GTG TTT CTT TTT GTA TCA GTG 3', denoted as SEQ ID NO:37, and antisense primer, JPI18, having the nucleic acid sequence 5' CGG AAT TCT TTA AAG GGA TTT AAC AC 3' (*Eco*RI site in bold), denoted SEQ ID NO:38.
30 The amplified gene sequence contained a natural *Bam*HI site about 24 bp downstream of the 3' end of JPI5 that was used for subcloning into the expression vector. Recombinant

molecule $p\lambda P_R$ -nfSPI2₁₁₃₉ was produced by digesting nfSPI2₁₁₈₅-containing PCR product with *Bam*HI and *Eco*RI restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector $P_R/T^2ori/S10HIS$ -RSET-A9, the production of which is described in PCT Publication No. US95/02941, by Tripp et al., published 9/14/95, Example 7, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel purified.

Recombinant molecule $p\lambda P_R$ -nfSPI2₁₁₃₉ was transformed into *E. coli* strain HB101 competent cells (available from Gibco/BRL, Gaithersburg, MD) to form recombinant cell *E.coli*: $p\lambda P_R$ -nfSPI2₁₁₃₉ using standard techniques as disclosed in Sambrook, et al., *ibid*.

The recombinant cells were cultured in enriched bacterial growth medium containing 0.1 mg/ml ampicillin and 0.1% glucose at about 32°C. When the cells reached an OD₆₀₀ of about 0.4-0.5, expression of recombinant protein was induced under heat shift conditions in which the cells were grown at 32°C for about 2 hours, and then grown at 42°C. Immunoblot analysis of recombinant cell *E.coli*: $p\lambda P_R$ -nfSPI2₁₁₃₉ lysates using the T7 tag monoclonal antibody (available from Novagen, Inc., Madison, WI) directed against the fusion portion of the recombinant PHis-PfSPI2₃₇₆ fusion protein identified proteins of appropriate size, namely an about 41 kD protein for each fusion protein.

Expression of the recombinant PHis-PfSPI2₃₇₆ fusion protein was improved by transforming supercoiled plasmid $p\lambda P_R$ -nfSPI2₁₁₃₉ DNA harvested from *E.coli*: $p\lambda P_R$ -nfSPI2₁₁₃₉ cells into the BL-21 strain of *E. coli* (available from Novagen). The amount of expression of PHis-PfSPI2₃₇₆ was confirmed by immunoblot using the method described immediately above.

E. coli cells expressing recombinant protein PHis-PfSPI2₃₇₆ were harvested from about 1 liter of media and suspended in about 40 ml of 50 mM Tris, pH 8, 50 mM NaCl, and 1 mg lysozyme (Lysis Buffer). The cells incubated in an ice bath for about 30 minutes (min) and then were centrifuged at about 30,000 x g for 30 min at 4°C. The supernatant (S1) was recovered and the pellet resuspended in about 40 ml Lysis Buffer containing 0.1% Triton X-100 and centrifuged at about 30,000 x g for 30 min at 4°C. The supernatant (S2) was recovered and the pellet resuspended in about 20 ml of

phosphate buffered saline (PBS) containing 8 M urea (S3). Aliquots of each supernatant were analyzed by SDS-PAGE and immunoblot using a T7 tag monoclonal antibody (available from Novagen, Inc., Madison, WI). The results indicated that the PHis-PfSPI2₃₇₆ protein was located in the final supernatant (S3). The PHis-PfSPI2₃₇₆ was loaded onto a 5 ml, metal chelating HiTrap™ column charged with NiCl₂ (available from Pharmacia Biotech Inc., Piscataway, NJ), previously equilibrated with PBS containing 8 M urea. The column was washed with PBS containing 8 M urea until all unbound protein was removed. Bound PHis-PfSPI2₃₇₆ protein was eluted with linear gradient from 0 to 1 M imidazole in PBS containing 8 M urea. Column fractions were analyzed for the presence of PHis-PfSPI2₃₇₆ by SDS-PAGE and immunoblot using a T7 tag monoclonal antibody. The results indicated that PHis-PfSPI2₃₇₆ was eluted at about 300 mM imidazole. The column fractions containing PHis-PfSPI2₃₇₆ protein were combined and diluted in 20 mM Tris, pH 8 containing 8 M urea in preparation for anion exchange chromatography. The sample was then loaded onto a 4.5 mm x 50 mm Poros 10 HQ anion exchange chromatography column (available from PerSeptive Biosystems, Framingham, MA), previously equilibrated with 20 mM Tris, pH 8 containing 8 M urea. Unbound proteins were washed from the column using the same buffer. Bound proteins were eluted with a linear gradient of from 0 to 1 M NaCl in 20 mM Tris, pH 8 containing 8 M urea. Column fractions were analyzed for the presence of PHis-PfSPI2₃₇₆ by SDS-PAGE. The results indicated that PHis-PfSPI2₃₇₆ was eluted at about 500 mM NaCl.

The purified PHis-PfSPI2₃₇₆ protein was used to produce an anti-SPI2 polyclonal antiserum as follows. Fractions containing PHis-PfSPI2₃₇₆ protein were combined and diluted to a concentration of about 0.1 mg/ml in PBS. A rabbit was immunized and boosted with about 1 mL of a 1:1 mix of antigen and adjuvant. The primary immunization was performed using antigen combined with Complete Freund's Adjuvant. About 500 µl of the mixture was injected subcutaneously into 5 different sites (0.1 ml/site) and 500 µl was injected intradermally into 5 different sites (0.1 ml/site) of the rabbit. Boosts were administered using antigen combined with Incomplete Freund's Adjuvant and were given on days 14 and 36 after the primary immunization, in 250 µl/site doses, intramuscularly, in 4 different sites. Blood samples were obtained prior to

immunization (pre-bleed), and approximately every two weeks after the primary immunization. Serum samples from the pre-immunization and days 27, 41, and 55 after the primary immunization were used for subsequent immunoblot experiments.

B. Recombinant molecule p λ P_R-nfSPI3₁₁₇₉, containing a portion of a flea serine protease inhibitor molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1225-nucleotide DNA fragment containing nucleotides spanning from about 351 through about 1570 of SEQ ID NO:13, denoted herein as nfSPI3₁₂₂₅, was PCR amplified from nucleic acid molecule nfSPI3₁₈₃₈, produced as described in Example 3, using sense primer JPI5 (SEQ ID NO:37), and antisense primer was JPI15, having the nucleic acid sequence 5' CGG AAT TCT AAT TGG TAA ATC TC 3' (*Eco*RI site in bold), denoted SEQ ID NO:39. The amplified gene sequence contained a natural *Bam*HI site about 24 bp downstream of the 3' end of JPI5 that was used for subcloning into the expression vector. Recombinant molecule p λ P_R-nfSPI3₁₁₇₉ was produced by digesting nfSPI3₁₂₂₅-containing PCR product with *Bam*HI and *Eco*RI restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²*ori*/S10HIS-RSET-A9, as described in Section A above, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel purified.

Recombinant molecule p λ P_R-nfSPI3₁₁₇₉ was transformed into *E. coli* strain HB101 competent cells (available from Gibco/BRL) to form recombinant cell *E. coli*:p λ P_R-nfSPI3₁₁₇₉ using standard techniques as disclosed in Sambrook, et al., *ibid*.

C. Recombinant molecule p λ P_R-nfSPI4₁₁₄₀, containing a portion of a flea serine protease inhibitor molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1186-nucleotide DNA fragment containing nucleotides spanning from about 8 through about 1186 of SEQ ID NO:19, denoted herein as nfSPI4₁₁₈₆, was PCR amplified from nucleic acid molecule nfSPI4₁₄₁₄, produced as described in Example 3, using sense primer JPI5 (SEQ ID NO:37), and antisense primer was JPI17, having the nucleic acid sequence 5' CGG AAT TCT TTT ATT CAG TTG TTG G 3' (*Eco*RI site in bold), denoted SEQ ID NO:40. The

amplified gene sequence contained a natural *Bam*HI site about 24 bp downstream of the 3' end of JPI5 that was used for subcloning into the expression vector. Recombinant molecule p λ P_R-nfSPI4₁₁₄₀ was produced by digesting nfSPI4₁₁₈₆-containing PCR product with *Bam*HI and *Eco*RI restriction endonucleases, column purifying the resulting
5 fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, as described in Section A above, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel purified.

Recombinant molecule p λ P_R-nfSPI4₁₁₄₀ was transformed into *E. coli* strain HB101 competent cells (available from Gibco/BRL) to form recombinant cell
10 *E.coli*:p λ P_R-nfSPI4₁₁₄₀ using standard techniques as disclosed in Sambrook, et al., *ibid*.

D. Recombinant molecule p λ P_R-nfSPI5₁₁₄₀, containing a portion of a flea serine protease inhibitor molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1186-nucleotide DNA
15 fragment containing nucleotides spanning from about 24 through about 1202 of SEQ ID NO:25, denoted herein as nfSPI5₁₁₈₆, was PCR amplified from nucleic acid molecule nfSPI5₁₄₉₂, produced as described in Example 3, using sense primer JPI5 (SEQ ID NO:37), and antisense primer was JPI17 (SEQ ID NO:40). The amplified gene sequence
20 used for subcloning into the expression vector. Recombinant molecule p λ P_R-nfSPI5₁₁₄₀ was produced by digesting nfSPI5₁₁₈₆-containing PCR product with *Bam*HI and *Eco*RI restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, as described in Section A above, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel
25 purified.

Recombinant molecule p λ P_R-nfSPI5₁₁₄₀ was transformed into *E. coli* strain HB101 competent cells (available from Gibco/BRL) to form recombinant cell
E.coli:p λ P_R-nfSPI5₁₁₄₀ using standard techniques as disclosed in Sambrook, et al., *ibid*.

E. Recombinant molecule p λ P_R-nfSPI6₁₁₃₆, containing a portion of a flea serine
30 protease inhibitor molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment

comprising 6 histidines was produced as follows. An about 1182-nucleotide DNA fragment containing nucleotides spanning from about 38 through about 1214 of SEQ ID NO:31, denoted herein as nfSPI6₁₁₈₂, was PCR amplified from nucleic acid molecule nfSPI6₁₄₅₄, produced as described in Example 3, using sense primer JPI5 (SEQ ID NO:37), and antisense primer was JPI16, having the nucleic acid sequence 5' CGG AAT TCA TAG AGT TTG AAC TC 3' (*EcoRI* site in bold), denoted SEQ ID NO:41. The amplified gene sequence contained a natural *Bam*HI site about 24 bp downstream of the 3' end of JPI5 that was used for subcloning into the expression vector. Recombinant molecule pλP_R-nfSPI6₁₁₃₆ was produced by digesting nfSPI6₁₁₈₂-containing PCR product with *Bam*HI and *Eco*RI restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, as described in Section A above, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel purified.

Recombinant molecule pλP_R-nfSPI6₁₁₃₆ was transformed into *E. coli* strain HB101 competent cells (available from BRL) to form recombinant cell *E.coli*:pλP_R-nfSPI6₁₁₃₆ using standard techniques as disclosed in Sambrook, et al., *ibid*.

Example 6

This Example describes the production in bacteria of several flea serine protease inhibitor proteins of the present invention.

Recombinant cells *E.coli*:pλP_R-nfSPI2₁₁₃₉, *E.coli*:pλP_R-nfSPI3₁₁₇₉, *E.coli*:pλP_R-nfSPI4₁₁₄₀, and *E.coli*:pλP_R-nfSPI6₁₁₃₆, produced as described in Example 5, were cultured in shake flasks containing an enriched bacterial growth medium containing 0.1 mg/ml ampicillin and 0.1% glucose at about 32°C. When the cells reached an OD₆₀₀ of about 0.4 to about 0.5, expression of flea pλP_R-nfSPI2₁₁₃₉, pλP_R-nfSPI3₁₁₇₉, pλP_R-nfSPI4₁₁₄₀, and pλP_R-nfSPI6₁₁₃₆, was induced by elevating the temperature to 42°C, and culturing the cells for about 3 hours. Protein production was monitored by SDS-PAGE of recombinant cell lysates, followed by Coomassie Blue staining and immunoblot analyses using a T7 Tag monoclonal antibody (available from Novagen, Inc.).

Recombinant cells *E.coli*:pλP_R-nfSPI2₁₁₃₉, *E.coli*:pλP_R-nfSPI3₁₁₇₉, *E.coli*:pλP_R-nfSPI4₁₁₄₀, and *E.coli*:pλP_R-nfSPI6₁₁₃₆ produced fusion proteins, denoted herein as PHis-

PfSPI2₃₇₆, PHis-PfSPI3₃₉₀, PHis-PfSPI4₃₇₆, and PHis-PfSPI6₃₇₆, that migrated with an apparent molecular weights of about 45 to 50 kD as predicted.

Example 7

This example describes analysis of the variable and constant domains of the
5 nucleic acid molecules of the present invention.

The sequences of each of the flea serine protease inhibitor cDNA molecules nfSPI1₁₅₈₄, nfSPI2₁₃₅₈, nfSPI3₁₈₃₈, nfSPI4₁₄₁₄, nfSPI5₁₄₉₂, and nfSPI6₁₄₅₄, presented in Example 4, were subdivided into three domains based on comparisons between the six sequences. The observed versions of the three domains are summarized in Table 1.

10 Domain I, spanning from about nucleotide 1 to about nucleotide 142 in nfSPI1₁₅₈₄, from about nucleotide 1 to about nucleotide 14 in nfSPI2₁₃₅₈, from about nucleotide 1 to about nucleotide 339 in nfSPI3₁₈₃₈, not present in nfSPI4₁₄₁₄, from about nucleotide 1 to about nucleotide 12 in nfSPI5₁₄₉₂, and from about nucleotide 1 to about nucleotide 26 in
15 nfSPI6₁₄₅₄, contains upstream untranslated sequences and the coding regions for the amino termini of the serine protease inhibitor proteins. Domain II, spanning from about nucleotide 143 to about nucleotide 1195 in nfSPI1₁₅₈₄, from about nucleotide 15 to about nucleotide 1067 in nfSPI2₁₃₅₈, from about nucleotide 340 to about nucleotide 1392 in
20 nfSPI3₁₈₃₈, from about nucleotide 1 to about nucleotide 1049 in nfSPI4₁₄₁₄, from about nucleotide 13 to about nucleotide 1065 in nfSPI5₁₄₉₂, and from about nucleotide 27 to about nucleotide 1079 in nfSPI6₁₄₅₄, consists of the central core of the coding sequence and encodes 350 amino acids that are extremely highly conserved (i.e. less than
approximately 2% variation) between the six serine protease inhibitor clones. The predicted mature N-terminus of the serine protease inhibitors is within Domain II; thus, the variability of Domain I should have no effect on the sequence of mature serine
25 protease inhibitor polypeptides. Domain III sequences are highly variable, yet still related to one another; Domain III, spanning from about nucleotide 1196 to about nucleotide 1584 in nfSPI1₁₅₈₄, from about nucleotide 1068 to about nucleotide 1358 in
nfSPI2₁₃₅₈, from about nucleotide 1393 to about nucleotide 1838 in nfSPI3₁₈₃₈, from about nucleotide 1050 to about nucleotide 1414 in nfSPI4₁₄₁₄, from about nucleotide
30 1066 to about nucleotide 1492 in nfSPI5₁₄₉₂, and from about nucleotide 1080 to about

nucleotide 1454 in nfSPI6₁₄₅₄, encodes the C-termini of the serine protease inhibitor proteins.

While not being bound by theory, the most probable explanation for the mixing of the domain versions within the six clones sequenced is a mechanism of alternative mRNA splicing. Such a pattern was described previously by Jiang et al., 1994, *J. Biol. Chem.* 269, 55-58 for serpins in *Manduca sexta*. For this family of serpins, eight exons encode a 336-amino acid constant region, followed by a 40-45-amino acid variable region that is encoded by the ninth exon. At least twelve alternative forms of the ninth exon are tandemly arranged in the genome between exons 8 and 10. Thus, mutually exclusive exon use can account for the variability the authors observed in cDNA clones.

Based on analogy to the *Manduca* system, flea serine protease inhibitors probably exhibit a similar gene structure in that the C-terminal variable region (Domain III) is encoded by multiple exons that are used in a mutually exclusive splicing mechanism. The flea serine protease inhibitor molecules appear to differ from *Manduca* in that for the flea molecules there are at least two alternative exons at the 5' end of the gene (Domain I) as well, and there does not appear to be final constant exon (exon 10 in *Manduca*) at the 3' end. It is probable that other versions of Domain III are present in the flea genome that were not observed in the six cDNA sequences presented herein.

Table 1. Summary of sequence variations of the three domains of flea serine protease inhibitor cDNA clones. Letters represent widely divergent sequences (e.g., A vs. B); numbers denote minor variations (i.e., less than 2%) between lettered sequences (e.g., K1 vs. K2).

<u>Clone</u>	<u>Domain I</u>	<u>Domain II</u>	<u>Domain III</u>
nfSe1 ₁₅₈₄	A	K1	W1
25 nfSe2 ₁₃₅₈	B	K2	X
nfSe3 ₁₈₃₈	B	K2	Y
nfSe4 ₁₄₁₄	missing	K2	Z
nfSe5 ₁₄₉₂	B	K3	Z
nfSe6 ₁₄₅₄	A	K2	W2

Example 8

This example describes the sequencing of several flea serine protease inhibitor variable domain nucleic acid molecules.

Nucleic acid molecules encoding serine protease inhibitor variable domains were identified as follows. Two primers were designed based on the 3' end of the constant domain sequence of nfSPI₄₁₄₁₄, referred to herein as primer 5' new BsaI or primer 5' new HincII. Each primer was designed so that, when used in conjunction with an antisense vector primer, a properly amplified fragment of a flea serine protease inhibitor genes. Primer 5' new BsaI has nucleic acid sequence 5' CAA AAC TGG TCT CCC CGC TC 3' (*BsaI* site in bold), represented herein as SEQ ID NO:42; and primer 5' new HincII has nucleic acid sequence 5' ATT ACA AAA TGT TGA CTT GC 3' (*HincII* site in bold), represented herein as SEQ ID NO:43. Primer 5' new *BsaI* and primer 5' new *HincII* were each used separately in combination with the vector specific primer T7 having nucleic acid sequence 5' TAA TAC GAC TCA CTA TAG GG 3', represented herein as SEQ ID NO:44.

The two primer pairs were used to amplify nucleic acid molecules using standard PCR amplification conditions (e.g., Sambrook et al., *ibid.*) from a variety of cDNA libraries representing different *C. felis* developmental stages. The cDNA libraries were produced as follows. The pre-pupal cDNA library was produced as described above in Example 3. A flea mixed instar cDNA library was produced using unfed 1st instar, bovine blood-fed 1st instar, bovine blood-fed 2nd instar and bovine blood-fed 3rd instar flea larvae (this combination of tissues is referred to herein as mixed instar larval tissues for purposes of this example). Total RNA was extracted from mixed instar using the method described above using about 5,164 mixed instar larvae. Poly A⁺ selected RNA was isolated as described above and about 6.34 µg of mixed instar poly A⁺ RNA was used to construct a mixed instar cDNA expression library in lambda Uni-ZAPTMXR vector (available from Stratagene), using Stratagene's ZAP-cDNA Synthesis Kit® protocol. The resultant mixed instar library was amplified to a titer of about 2.17 x 10¹⁰ pfu/ml with about 97% recombinants. An unfed whole adult flea cDNA library was

produced by the standard method generally described in Example 8 of related PCT Publication No. WO 96/11706.

A bovine blood-fed flea gut cDNA library was produced as follows. Total RNA was extracted from approximately 3500 guts from bovine blood-fed fleas using a standard guanidinium thiocyanate procedure for lysis and denaturation of the gut tissue, followed by centrifugation in cesium chloride to pellet the RNA. Messenger RNA was isolated from the total RNA using a Fast Track™ Kit (available from InVitrogen, San Diego, CA). A bovine blood-fed flea gut cDNA expression library was constructed in lambda Uni-ZAP™XR vector (available from Stratagene), using Stratagene's ZAP-cDNA Synthesis Kit® protocol

PCR products using the different cDNA libraries were each gel purified and cloned into the TA Vector™ (available from InVitrogen). The nucleic acid molecule was subjected to nucleic acid sequencing using the Sanger dideoxy chain termination method, as described in Sambrook et al., *ibid*.

A. A first flea serine protease inhibitor variable domain nucleic acid molecule isolated from the mixed instar cDNA library was determined to comprise nucleic acid molecule nfSPI7₅₄₉, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:45. Translation of SEQ ID NO:45 suggests that nucleic acid molecule nfSPI7₅₄₉ encodes a portion of a serine protease inhibitor protein of about 134 amino acids, referred to herein as PfSPI7₁₃₄, having amino acid sequence SEQ ID NO:46, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:45 and the last codon spans from nucleotide 402 through nucleotide 404 of SEQ ID NO:45. The complement of SEQ ID NO:45 is represented herein by SEQ ID NO:47. Comparison of amino acid sequence SEQ ID NO:46 (i.e., the amino acid sequence of PfSPI7₁₃₄) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:46, showed the most homology, i.e., about 34% identity, between SEQ ID NO:46 and *Mus musculus* antithrombin III precursor protein. Comparison of nucleic acid sequence SEQ ID NO:45 (i.e., the nucleic acid sequence of nfSPI7₅₄₉) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:45, showed the most homology, i.e., about 38% identity, between SEQ ID NO:45 and human bomapin gene.

B. A second flea serine protease inhibitor variable domain nucleic acid molecule isolated from the mixed instar cDNA library was determined to comprise nucleic acid molecule nfSPI8₅₄₉, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:48. Translation of SEQ ID NO:48 suggests that nucleic acid molecule nfSPI8₅₄₉ encodes a serine protease inhibitor variable domain protein of about 149 amino acids, referred to herein as PfSPI8₁₄₉, having amino acid sequence SEQ ID NO:49, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:48 and the last codon spans from nucleotide 447 through nucleotide 449 of SEQ ID NO:48. The complement of SEQ ID NO:48 is represented herein by SEQ ID NO:50. Comparison of amino acid sequence SEQ ID NO:49 (i.e., the amino acid sequence of PfSPI8₁₄₉) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:49, showed the most homology, i.e., about 36% identity, between SEQ ID NO:49 and human bomapin precursor protein. Comparison of nucleic acid sequence SEQ ID NO:48 (i.e., the nucleic acid sequence of nfSPI8₅₄₉) with nucleic acid sequences reported in GeEmbl indicates that SEQ ID NO:48, showed the most homology, i.e., about 41% identity, between SEQ ID NO:48 and human bomapin gene.

C. A third flea serine protease inhibitor variable domain nucleic acid molecule isolated from the bovine blood-fed gut cDNA library was determined to comprise nucleic acid molecule nfSPI9₅₈₁, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:51. Translation of SEQ ID NO:51 suggests that nucleic acid molecule nfSPI9₅₈₁ encodes a serine protease inhibitor variable domain protein of about 136 amino acids, referred to herein as PfSPI9₁₃₆, having amino acid sequence SEQ ID NO:52, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:51 and the last codon spans from nucleotide 408 through nucleotide 410 of SEQ ID NO:51. The complement of SEQ ID NO:51 is represented herein by SEQ ID NO:53. Comparison of amino acid sequence SEQ ID NO:52 (i.e., the amino acid sequence of PfSPI9₁₃₆) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:52, showed the most homology, i.e., about 45% identity, between SEQ ID NO:52 and *Bombyx mori* anti-trypsin precursor protein. Comparison of nucleic acid sequence SEQ ID NO:51 (i.e., the nucleic acid sequence of nfSPI9₅₈₁) with nucleic acid sequences reported in GenBank indicates that SEQ ID NO:51, showed the

most homology, i.e., about 52% identity, between SEQ ID NO:51 and *Bombyx mori* anti-trypsin gene.

D. A fourth flea serine protease inhibitor variable domain nucleic acid molecule isolated from the flea pre-pupal cDNA library was determined to comprise nucleic acid molecule nfSPI10₆₅₄, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:54. Translation of SEQ ID NO:54 suggests that nucleic acid molecule nfSPI10₆₅₄ encodes a serine protease inhibitor variable domain protein of about 118 amino acids, referred to herein as PfSPI10₁₁₈, having amino acid sequence SEQ ID NO:55, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:54 and the last codon spans from nucleotide 354 through nucleotide 356 of SEQ ID NO:54. The complement of SEQ ID NO:54 is represented herein by SEQ ID NO:56. Comparison of amino acid sequence SEQ ID NO:55 (i.e., the amino acid sequence of PfSPI10₁₁₈) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:55, showed the most homology, i.e., about 38% identity, between SEQ ID NO:55 and *Manduca sexta* alaserpin precursor protein. Comparison of nucleic acid sequence SEQ ID NO:54 (i.e., the nucleic acid sequence of nfSPI10₆₅₄) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:54, showed the most homology, i.e., about 41% identity, between SEQ ID NO:54 and human bomapin gene.

E. A fifth flea serine protease inhibitor variable domain nucleic acid molecule isolated from the flea pre-pupal cDNA library was determined to comprise nucleic acid molecule nfSPI11₆₇₀, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:57. Translation of SEQ ID NO:57 suggests that nucleic acid molecule nfSPI11₆₇₀ encodes a serine protease inhibitor variable domain protein of about 125 amino acids, referred to herein as PfSPI11₁₂₅, having amino acid sequence SEQ ID NO:58, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:57 and the last codon spans from nucleotide 375 through nucleotide 377 of SEQ ID NO:57. The complement of SEQ ID NO:57 is represented herein by SEQ ID NO:59. Comparison of amino acid sequence SEQ ID NO:58 (i.e., the amino acid sequence of PfSPI11₁₂₅) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:58, showed the most homology, i.e., about 43% identity, between SEQ ID NO:58 and *Manduca sexta* alaserpin precursor protein. Comparison of nucleic acid

sequence SEQ ID NO:57 (i.e., the nucleic acid sequence of nfSPI1₁₆₇₀) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:57, showed the most homology, i.e., about 40% identity, between SEQ ID NO:57 and human bomapin gene.

F. A sixth flea serine protease inhibitor variable domain nucleic acid molecule isolated from the unfed whole adult flea cDNA library was determined to comprise nucleic acid molecule nfSPI12₇₀₆, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:60. Translation of SEQ ID NO:60 suggests that nucleic acid molecule nfSPI12₇₀₆ encodes a serine protease inhibitor variable domain protein of about 136 amino acids, referred to herein as PfSPI12₁₃₆, having amino acid sequence SEQ ID NO:61, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:60 and the last codon spans from nucleotide 408 through nucleotide 410 of SEQ ID NO:60. The complement of SEQ ID NO:60 is represented herein by SEQ ID NO:62. Comparison of amino acid sequence SEQ ID NO:61 (i.e., the amino acid sequence of PfSPI12₁₃₆) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:61, showed the most homology, i.e., about 45% identity, between SEQ ID NO:61 and *Manduca sexta* alaserpin precursor protein protein. Comparison of nucleic acid sequence SEQ ID NO:60 (i.e., the nucleic acid sequence of nfSPI12₇₀₆) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:60, showed the most homology, i.e., about 38% identity, between SEQ ID NO:60 and human bomapin gene.

G. A seventh flea serine protease inhibitor variable domain nucleic acid molecule isolated from the flea pre-pupal cDNA library was determined to comprise nucleic acid molecule nfSPI13₆₂₃, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:63. Translation of SEQ ID NO:63 suggests that nucleic acid molecule nfSPI13₆₂₃ encodes a serine protease inhibitor variable domain protein of about 122 amino acids, referred to herein as PfSPI13₁₂₂, having amino acid sequence SEQ ID NO:64, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:63 and the last codon spans from nucleotide 366 through nucleotide 368 of SEQ ID NO:63. The complement of SEQ ID NO:63 is represented herein by SEQ ID NO:65. Comparison of amino acid sequence SEQ ID NO:64 (i.e., the amino acid sequence of PfSPI13₁₂₂) with amino acid sequences reported in SwissProt indicates that

SEQ ID NO:64, showed the most homology, i.e., about 39% identity, between SEQ ID NO:64 and human leukocyte esterase inhibitor protein. Comparison of nucleic acid sequence SEQ ID NO:63 (i.e., the nucleic acid sequence of nfSPI13₆₂₃) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:63, showed the most
5 homology, i.e., about 37% identity, between SEQ ID NO:63 and human bomapin gene.

H. A eighth flea serine protease inhibitor variable domain nucleic acid molecule isolated from the bovine blood-fed flea gut cDNA library was determined to comprise nucleic acid molecule nfSPI14₇₃₁, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:66. Translation of SEQ ID NO:66
10 suggests that nucleic acid molecule nfSPI14₇₃₁ encodes a serine protease inhibitor variable domain protein of about 137 amino acids, referred to herein as PfSPI14₁₃₇, having amino acid sequence SEQ ID NO:67, assuming the first codon spans from nucleotide 3 through nucleotide 5 of SEQ ID NO:66 and the last codon spans from nucleotide 411 through nucleotide 413 of SEQ ID NO:66. The complement of SEQ ID
15 NO:66 is represented herein by SEQ ID NO:68. Comparison of amino acid sequence SEQ ID NO:67 (i.e., the amino acid sequence of PfSPI14₁₃₇) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:67, showed the most homology, i.e., about 40% identity, between SEQ ID NO:67 and *Equus caballus* esterase inhibitor protein. Comparison of nucleic acid sequence SEQ ID NO:66 (i.e., the nucleic acid
20 sequence of nfSPI14₇₃₁) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:66, showed the most homology, i.e., about 38% identity, between SEQ ID NO:66 and human bomapin gene.

I. A ninth flea serine protease inhibitor variable domain nucleic acid molecule isolated from the unfed whole adult flea cDNA library was determined to
25 comprise nucleic acid molecule nfSPI15₆₈₅, the nucleic acid sequence of the coding strand which is denoted herein as SEQ ID NO:69. Translation of SEQ ID NO:69 suggests that nucleic acid molecule nfSPI15₆₈₅ encodes a serine protease inhibitor variable domain protein of about 135 amino acids, referred to herein as PfSPI15₁₃₅, having amino acid sequence SEQ ID NO:70, assuming the first codon spans from
30 nucleotide 3 through nucleotide 5 of SEQ ID NO:69 and the last codon spans from nucleotide 405 through nucleotide 407 of SEQ ID NO:69. The complement of SEQ ID

NO:69 is represented herein by SEQ ID NO:71. Comparison of amino acid sequence SEQ ID NO:70 (i.e., the amino acid sequence of PfSPI15₁₃₅) with amino acid sequences reported in SwissProt indicates that SEQ ID NO:70, showed the most homology, i.e., about 48% identity, between SEQ ID NO:70 and *Bombyx mori* antichymotrypsin II protein. Comparison of nucleic acid sequence SEQ ID NO:69 (i.e., the nucleic acid sequence of nfSPI15₆₈₅) with nucleic acid sequences reported in GenEmbl indicates that SEQ ID NO:69, showed the most homology, i.e., about 38% identity, between SEQ ID NO:69 and human antithrombin III variant gene.

Example 9

10 This example discloses the production of a several recombinant cells of the present invention using serine protease inhibitor variable domain nucleic acid molecules of the present invention.

Each of nucleic acid molecules nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI12₇₀₆, nfSPI13₆₂₃ and nfSPI15₆₈₅, were digested with either the restriction enzymes *HincII* and *XhoI*, or *BsaI* and *XhoI*. The resulting *HincII* and *XhoI*, or *BsaI* and *XhoI* digested fragments were ligated to a portion of DNA that had been isolated from nfSPI4₁₄₁₄ digested with *BamHI* and *HincII*, or *BamHI* and *BsaI*. The nfSPI4₁₄₁₄ *BamHI* and *HincII* fragment, or nfSPI4₁₄₁₄ *BamHI* and *BsaI* fragment, encoded the majority of the constant domain of nfSPI4₁₄₁₄. The resulting ligation products that include chimeric serine protease inhibitor open reading frames, are referred to herein as nfSPIC4:V7, nfSPIC4:V8, nfSPIC4:V9, nfSPIC4:V10, nfSPIC4:V12, nfSPIC4:V13 and nfSPIC4:V15, respectively. The nfSPIC4:V7, nfSPIC4:V9, nfSPIC4:V10 or nfSPIC4:V12 ligation products were then digested with the restriction enzymes *BamHI* and *XhoI* and separately ligated into pBluescript vector which had been digested with the same restriction enzymes. The resulting ligation products are referred to herein as pBluSPI:C4:V7, pBluSPI:C4:V9, pBluSPI:C4:V10 and pBluSPI:C4:V12, respectively.

A. Recombinant molecule pλP_R-nfSPIC4:V7₁₁₆₈, containing a chimeric serine protease inhibitor open reading frame molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1168-nucleotide DNA fragment denoted herein as nfSPIC4:V7₁₁₆₈ containing nucleotides

spanning from 1 through 761 of nfSPI4₁₄₁₄ ligated to nucleotides spanning from 1 through 407 of nfSPI7₅₄₉, was PCR amplified from nucleic acid molecule pBluSPI:C4:V7, using sense primer T-3pBS, having the nucleic acid sequence 5' ATT AAC CCT CAC TAA AG 3' (SEQ ID NO:83), and antisense primer, Srp73'end, having
 5 nucleic acid sequence 5' GCG **GAA TTC** TTA AGG ATT AAC GTG TTG AAC 3' and denoted herein as SEQ ID NO:93 (*Eco*RI site shown in bold). The amplified gene sequence contained a natural *Bam*HI site about 100 bp downstream of the T-3pBS primer that was used for subcloning into the expression vector. Recombinant molecule pλP_R-nfSPIC4:V7₁₁₆₈ was produced by digesting nfSPIC4:V7₁₁₆₈ with *Bam*HI and *Eco*RI
 10 restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²*ori*/S10HIS-RSET-A9, the production of which is described in PCT Publication No. US95/02941, by Tripp et al., published 9/14/95, Example 7, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel purified.

15 Recombinant molecule pλP_R-nfSPIC4:V7₁₁₆₈ was transformed into *E. coli* strain HB101 competent cells (available from Gibco/BRL, Gaithersburg, MD) to form recombinant cell *E.coli*:pλP_R-nfSPIC4:V7₁₁₆₈ using standard techniques as disclosed in Sambrook, et al., *ibid*.

B. Recombinant molecule pλP_R-nfSPIC4:V9₁₁₇₄, was produced using the
 20 methods described above in section 9(A) except the antisense primer used to produce a PCR product from pBluSPI:C4:V9 was Srp93'end, having nucleic acid sequence 5' **GGA ATT CTT ATT GCA CAA ATC ATC C** 3' and denoted herein as SEQ ID NO:94 (*Eco*RI site shown in bold). An about 1174-nucleotide DNA fragment denoted herein as nfSPIC4:V9₁₁₇₄ containing nucleotides spanning from 1 through 794 of nfSPI4₁₄₁₄ and
 25 nucleotides spanning from 22 through 413 of SEQ ID NO:51, was PCR amplified from nucleic acid molecule pBluSPI:C4:V9, produced as described in section 9. Recombinant molecule pλP_R-nfSPIC4:V9₁₁₇₄ was produced by digesting nfSPIC4:V9₁₁₇₄ with *Bam*HI and *Eco*RI restriction endonucleases, gel purifying the resulting fragment and subcloning the fragment into the expression vector P_R/T²*ori*/S10HIS-RSET-A9, which had been
 30 similarly cleaved with *Bam*HI and *Eco*RI and gel purified, to produce the recombinant molecule pλP_R-nfSPIC4:V9₁₁₇₄.

Recombinant molecule p λ P_R-nfSPIC4:V9₁₁₇₄ was transformed into *E. coli* strain HB101 competent cells to form recombinant cell *E. coli*:p λ P_R-nfSPIC4:V9₁₁₇₄ using methods described in Section 9(A).

C. Recombinant molecule p λ P_R-nfSPIC4:V10₁₁₅₉, was produced using the methods described above in section 9(A) except the antisense primer used to produce a PCR product from pBluSPI:C4:V10 was Srp103'end, having nucleic acid sequence 5' **GCG GAA TTC AAC AAA AGT GTG TTC** 3' and denoted herein as SEQ ID NO:87 (*Eco*RI site shown in bold) and the sense primer used was the T-3pBS primer (SEQ ID NO:83). An about 1159-nucleotide DNA fragment denoted herein as nfSPIC4:V10₁₁₅₉ containing nucleotides spanning from 1 through 803 of nfSPI4₁₄₁₄ and nucleotides spanning from 1 through 356 of SEQ ID NO:54, was PCR amplified from nucleic acid molecule pBluSPI:C4:V10, produced as described in section 9. Recombinant molecule p λ P_R-nfSPIC4:V10₁₁₅₉ was produced by digesting nfSPIC4:V10₁₁₅₉ with *Bam*HI and *Eco*RI restriction endonucleases, gel purifying the resulting fragment and subcloning the fragment into the expression vector P_R/T²ori/S10HIS-RSET-A9, which had been similarly cleaved with *Bam*HI and *Eco*RI and gel purified, to produce the recombinant molecule p λ P_R-nfSPIC4:V10₁₁₅₉.

Recombinant molecule p λ P_R-nfSPIC4:V10₁₁₅₉ was transformed into *E. coli* strain HB101 competent cells to form recombinant cell *E. coli*:p λ P_R-nfSPIC4:V10₁₁₅₉ using methods described in Section 9(A).

D. Recombinant molecule p λ P_R-nfSPIC4:V8₁₂₂₂, containing a chimeric serine protease inhibitor open reading frame molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1222 nucleotide DNA fragment denoted herein as nfSPIC4:V8₁₂₂₂ containing nucleotides spanning from 1 to 794 of nfSPI4₁₄₁₄ ligated to nucleotides spanning from 22 through 449 of nfSPI8₅₄₉ was PCR amplified from nucleic acid molecule nfSPIC4:V8 using sense primer serpin5' end having nucleic acid sequence 5' **ATA GGA TCC CCA GGA ATT GTC** 3' (SEQ ID NO 84; *Bam*HI site in bold), and antisense primer, Srp8 3'end, having nucleic acid sequence 5' **GCG AGA TCT CTA GTT ATT AAT ATT GGT TAA** 3' and denoted herein as SEQ ID NO:85 (*Bgl*III site shown in bold). Recombinant

molecule p λ P_R-nfSPIC4:V8 was produced by digesting nfSPIC4:V8₁₂₂₂ with *Bam*HI and *Bgl*II restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, which had been similarly cleaved with *Bam*HI and *Bgl*II and gel purified, to produce
5 the recombinant molecule p λ P_R-nfSPIC4:V8₁₂₂₂.

Recombinant molecule p λ P_R-nfSPIC4:V8₁₂₂₂ was transformed into *E. coli* strain HB101 competent cells to form recombinant cell *E.coli*:p λ P_R-nfSPIC4:V8₁₂₂₂ using methods described in Section 9(A).

E. Recombinant molecule p λ P_R-nfSPIC4:V15₁₁₇₉, containing a chimeric
10 serine protease inhibitor open reading frame molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1179 nucleotide DNA fragment denoted herein as nfSPIC4:V15₁₁₇₉ containing nucleotides spanning from 1 to 794 of nfSPI4₁₄₁₄ ligated to nucleotides spanning from 22
15 through 449 of nfSPI15₆₈₅ was PCR amplified from nucleic acid molecule nfSPIC4:V15 using the sense primer serpin5' end (SEQ ID NO:84) and the antisense primer, Srp15 3', having nucleic acid sequence 5' GCGGAATTCTCATGGTGACTGAACGCG 3' (denoted herein as SEQ ID NO:86; *Eco*R1 site shown in bold). Recombinant molecule p λ P_R-nfSPIC4:V15₁₁₇₉ was produced by digesting nfSPIC4:V15₁₁₇₉ with *Bam*HI and
20 *Eco*R1 restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, which had been similarly cleaved with *Bam*HI and *Eco*R1 and gel purified, to produce the recombinant molecule p λ P_R-nfSPIC4:V15₁₁₇₉.

Recombinant molecule p λ P_R-nfSPIC4:V15₁₁₇₉ was transformed into *E. coli* strain
25 HB101 competent cells to form recombinant cell *E.coli*:p λ P_R-nfSPIC4:V15₁₁₇₉ using methods described in Section 9(A).

F. Recombinant molecule p λ P_R-nfSPIC4:V12₁₁₇₁, containing a chimeric serine protease inhibitor open reading frame molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding
30 a poly-histidine segment comprising 6 histidines was produced as follows. An about 1171 nucleotide DNA fragment denoted herein as nfSPIC4:V12₁₁₇₁ containing

nucleotides spanning from 1 to 761 of nfSPI4₁₄₁₄ ligated to nucleotides spanning from 1 through 410 of nfSPI12₇₀₆ was PCR amplified from nucleic acid molecule pBluSPIC4:V12 using sense primer T-3pBS (SEQ ID NO:83), and antisense primer, Srp123' end, having nucleic acid sequence 5' GCG **GAA TTC** TTA TTT GGG AGA TAT AAC TCG 3' and denoted herein as SEQ ID NO:91 (*Eco*R1 site shown in bold).
 5 Recombinant molecule pλP_R-nfSPIC4:V12₁₁₇₁ was produced by digesting nfSPIC4:V12₁₁₇₁ with *Bam*HI and *Eco*R1 restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, which had been similarly cleaved with *Bam*HI and *Eco*R1
 10 and gel purified, to produce the recombinant molecule pλP_R-nfSPIC4:V12₁₁₇₁.

Recombinant molecule pλP_R-nfSPIC4:V12₁₁₇₁ was transformed into *E. coli* strain HB101 competent cells to form recombinant cell *E.coli*:pλP_R-nfSPIC4:V12₁₁₇₁ using methods described in Section 9(A).

G. Recombinant molecule pλP_R-nfSPIC4:V13₁₁₇₁, containing a chimeric
 15 serine protease inhibitor open reading frame molecule operatively linked to bacteriophage lambda transcription control sequences and to a fusion sequence encoding a poly-histidine segment comprising 6 histidines was produced as follows. An about 1171 nucleotide DNA fragment denoted herein as nfSPIC4:V13₁₁₇₁ containing nucleotides spanning from 1 to 803 of nfSPI4₁₄₁₄ ligated to nucleotides spanning from 1 through 368 of nfSPI13₆₂₃ was PCR amplified from nucleic acid molecule nfSPIC4:V13
 20 using the sense primer serpin5' end (SEQ ID NO:84), and antisense primer Srp13 3', having nucleic acid sequence 5' CGC **GAA TTC** TCA TTC GAC AAA ATG ACC 3' and denoted herein as SEQ ID NO:92 (*Eco*RI site shown in bold). Recombinant molecule pλP_R-nfSPIC4:V13₁₁₇₁ was produced by digesting nfSPIC4:V13₁₁₇₁ with
 25 *Bam*HI and *Eco*RI restriction endonucleases, column purifying the resulting fragment, and directionally subcloning the fragment into expression vector P_R/T²ori/S10HIS-RSET-A9, which had been similarly cleaved with *Bam*HI and *Eco*R1 and gel purified, to produce the recombinant molecule pλP_R-nfSPIC4:V13₁₁₇₁.

Recombinant molecule pλP_R-nfSPIC4:V13₁₁₇₁ was transformed into *E. coli* strain
 30 HB101 competent cells to form recombinant cell *E.coli*:pλP_R-nfSPIC4:V13₁₁₇₁ using methods described in Section 9(A).

Example 10

This Example describes the production in bacteria of several flea serine protease inhibitor proteins of the present invention.

Recombinant cells *E.coli*:p λ P_R-nfSPIC4:V7₁₁₆₈, *E.coli*:p λ P_R-nfSPIC4:V8₁₂₂₂,
 5 *E.coli*:p λ P_R-nfSPIC4:V9₁₁₇₄, *E.coli*:p λ P_R-nfSPIC4:V10₁₁₅₉, *E.coli*:p λ P_R-
 nfSPIC4:V12₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V13₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V15₁₁₇₉, produced
 as described in Example 9, were cultured in shake flasks containing an enriched bacterial
 growth medium containing 0.1 mg/ml ampicillin and 0.1% glucose at about 32°C. When
 the cells reached an OD₆₀₀ of about 0.4 to about 0.5, expression of flea *E.coli*:p λ P_R-
 10 nfSPIC4:V7₁₁₆₈, *E.coli*:p λ P_R-nfSPIC4:V9₁₁₇₄, *E.coli*:p λ P_R-nfSPIC4:V10₁₁₅₉, *E.coli*:p λ P_R-
 nfSPIC4:V12₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V13₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V15₁₁₇₉, were each
 induced by elevating the temperature to 42°C, and culturing the cells for about 3 hours.
 Expression of flea *E.coli*:p λ P_R-nfSPIC4:V8₁₂₂₂ was induced by the addition of 0.5 mM
 isopropyl-B-D-thiogalactoside (IPTG) to the culture medium, and the cells were cultured
 15 for about 2 hours at about 32°C.

Protein production was monitored by SDS-PAGE of recombinant cell lysates and
 immunoblot analyses using a T7 Tag monoclonal antibody (available from Novagen,
 Inc.) and the anti-SPI2 polyclonal antiserum (described in detail in Example 5).
 Recombinant cells *E.coli*:p λ P_R-nfSPIC4:V7₁₁₆₈, *E.coli*:p λ P_R-nfSPIC4:V9₁₁₇₄ and
 20 *E.coli*:p λ P_R-nfSPIC4:V15₁₁₇₉ produced fusion proteins, denoted herein as PHis-
 PfSPIC4:V7, PHis-PfSPIC4:V9 and PHis-PfSPIC4:V15 that migrated with an apparent
 molecular weight of about 45 kD as predicted. Recombinant cells *E.coli*:p λ P_R-
 nfSPIC4:V10₁₁₅₉ produced the fusion protein denoted herein as PHis-PfSPIC4:V10 that
 migrated with an apparent molecular weight of about 44 kD as predicted. Recombinant
 25 cells *E.coli*:p λ P_R-nfSPIC4:V8₁₂₂₂ produced the fusion protein denoted herein as PHis-
 PfSPIC4:V8 that migrated with an apparent molecular weight of about 51 kD as
 predicted. Recombinant cells *E.coli*:p λ P_R-nfSPIC4:V12₁₁₇₁ and *E.coli*:p λ P_R-
 nfSPIC4:V13₁₁₇₁ produced the fusion protein denoted herein as PHis-PfSPIC4:V12 and
 PHis-PfSPIC4:V13, respectively, each of which migrated with an apparent molecular
 30 weight of about 49 kD as predicted.

Example 11

This example demonstrates the production of a serine protease inhibitor protein of the present invention in eukaryotic cells.

A. Recombinant molecule pBv-nfSPI3₁₂₂₂, containing a flea serine protease inhibitor nucleic acid molecule spanning nucleotides from about 325 through about 1546 of SEQ ID NO:13, operatively linked to baculovirus polyhedron transcription control sequences were produced in the following manner. A PCR fragment of 1222 nucleotides, herein denoted nfSPI3₁₂₂₂, having SEQ ID NO:72 was amplified from nfSPI3₁₈₃₈ using the sense primer Serpin3For, having the nucleic acid sequence 5' - GGA **AGA TCT ATA AAT ATG CCG CGT CCT CAG TTT G** -3' (SEQ ID NO:73; *Bgl*III site shown in bold) and the antisense primer Serpin3Rev, having the nucleic acid sequence 5' - **CGG AAT TCT AAT TGG TAA ATC TCC CAG AG** -3' (SEQ ID NO:74; *Eco*RI site shown in bold). A portion of the sense primer was designed from the pol h sequence of baculovirus with modifications to enhance expression in the baculovirus system.

The resulting 1222-bp PCR product (referred to as Bv-nfSPI3₁₂₂₂) was digested with *Bgl*III and *Eco*RI restriction endonucleases and subcloned into unique *Bgl*III and *Eco*RI sites of pVL1392 baculovirus shuttle plasmid (available from Pharmingen, San Diego, CA) to produce the recombinant molecule referred to herein as pVL-nfSPI3₁₂₂₂.

The resultant recombinant molecule pVL-nfSPI3₁₂₂₂, was verified for proper insert orientation by restriction mapping. Such a recombinant molecule can be co-transfected with a linear Baculogold baculovirus DNA (available from Pharmingen) into *S. frugiperda* Sf9 cells (available from InVitrogen) to form the recombinant cells denoted *S. frugiperda*:pVL-nfSPI3₁₂₂₂. *S. frugiperda*:pVL-nfSPI3₁₂₂₂ was cultured in order to produce a flea serine protease inhibitor protein PfSPI3₄₀₆ (referred to herein as SEQ ID NO:95).

An immunoblot of supernatant from cultures of *S. frugiperda*:pVL-nfSPI3₁₂₂₂ cells producing the flea serine protease inhibitor protein PfSPI3₄₀₆ was performed using the anti-SPI2 polyclonal antiserum described in detail in Example 5. Blots were incubated using serum samples from the pre-bleed or from serum collected 14 days after

the first boost of the rabbit. Analysis of the supernatant from cultures of *S.*

frugiperda:pVL-nfSPI3₁₂₂₂ cells identified an about 41 kD and about 46 kD proteins.

B. Recombinant molecule pBv-nfSPI6₁₁₅₅, containing a flea serine protease inhibitor nucleic acid molecule spanning nucleotides from about 154 through about 1308
 5 of SEQ ID NO:31, operatively linked to baculovirus polyhedron transcription control sequences were produced in the following manner. A PCR fragment of 1155 nucleotides, herein denoted nfSPI6₁₁₅₅, having SEQ ID NO:75 was amplified from nfSPI6₁₄₅₄ using the sense primer Serpin6For, having the nucleic acid sequence 5'- GGA
 AGA **TCT** ATA AAT ATG ATT AAC GCA CGA CTT -3' (SEQ ID NO:76; *Bgl*III site
 10 shown in bold) and the antisense primer Serpin6Rev, having the nucleic acid sequence 5'-CCG **GAA TTC** ATA GAG TTT GAA CTC GCC C -3' (SEQ ID NO:77; *Eco*RI site shown in bold). A portion of the sense primer was designed from the pol h sequence of baculovirus with modifications to enhance expression in the baculovirus system.

The resulting 1155-bp PCR product (referred to as Bv-nfSPI6₁₁₅₅) was digested
 15 with *Bgl*III and *Eco*RI restriction endonucleases and subcloned into unique *Bgl*III and *Eco*RI sites of pVL1392 baculovirus shuttle plasmid to produce the recombinant molecule referred to herein as pVL-nfSPI6₁₁₅₅.

The resultant recombinant molecule pVL-nfSPI6₁₁₅₅, was verified for proper insert orientation by restriction mapping. Such a recombinant molecule can be co-
 20 transfected with a linear Baculogold baculovirus DNA into *S. frugiperda* Sf9 cells to form the recombinant cells denoted *S. frugiperda*:pVL-nfSPI6₁₁₅₅. *S. frugiperda*:pVL-nfSPI6₁₁₅₅ was cultured in order to produce a flea serine protease inhibitor protein PfSPI6₃₈₅ (referred to herein as SEQ ID NO:96).

An immunoblot of supernatant from cultures of *S. frugiperda*:pVL-nfSPI6₁₁₅₅
 25 cells producing the flea serine protease inhibitor protein PfSPI6₃₈₅ was performed using the anti-SPI2 polyclonal antiserum described in detail in Example 5. Blots were incubated using serum samples from the pre-bleed or from serum collected 14 days after the first boost of the rabbit. Analysis of the supernatant from cultures of *S.*
frugiperda:pVL-nfSPI6₁₁₅₅ cells identified an about 41 kD and about 45 kD proteins.

30 C. Recombinant molecule pBv-nfSPI2₁₀₆₅, containing a flea serine protease inhibitor nucleic acid molecule spanning nucleotides from about 102 through about 1066

of SEQ ID NO:7, operatively linked to baculovirus polyhedron transcription control sequences were produced in the following manner. A PCR fragment of 1066 nucleotides, herein denoted nfSPI2₁₀₆₅, having SEQ ID NO:78 was amplified from nfSPI2₁₃₅₈ using the sense primer Serpin2For, having the nucleic acid sequence 5' - GCG
 5 **GAA TTC** GAT CCC CAG GAA TTG TCT ACA AGT ATT AAC C -3' (SEQ ID NO:79; *Eco*RI site shown in bold) and the antisense primer Serpin2Rev, having the nucleic acid sequence 5' - GCG **AGA TCT** TTA AAG GGA TTT AAC ACA TCC ACT GAA CAA AAC AG -3' (SEQ ID NO:80; *Bgl*II site shown in bold).

The resulting 1065-bp PCR product (referred to as Bv-nfSPI2₁₀₆₅) was digested
 10 with *Bgl*II and *Eco*RI restriction endonucleases and subcloned into unique *Bgl*II and *Eco*RI sites of pAcGP67 (available from Pharmingen)s baculovirus shuttle plasmid to produce the recombinant molecule referred to herein as pAcG-nfSPI2₁₀₆₅.

The resultant recombinant molecule pAcG-nfSPI2₁₀₆₅, was verified for proper insert orientation by restriction mapping. Such a recombinant molecule can be co-
 15 transfected with a linear Baculogold baculovirus DNA into *S. frugiperda* Sf9 cells to form the recombinant cells denoted *S. frugiperda*:pAcG-nfSPI2₁₀₆₅. *S. frugiperda*:pAcG-nfSPI2₁₀₆₅ was cultured in order to produce a flea serine protease inhibitor protein PfSPI2₃₅₄ (referred to herein as SEQ ID NO:97).

An immunoblot of supernatant from cultures of *S. frugiperda*:pAcG-nfSPI2₁₀₆₅
 20 cells producing the flea serine protease inhibitor protein PfSPI2₃₅₅ was performed using the anti-SPI2 polyclonal antiserum described in detail in Example 5. Blots were incubated using serum samples from the pre-bleed or from serum collected 14 days after the first boost of the rabbit. Analysis of the supernatant from cultures of *S. frugiperda*:pAcG-nfSPI2₁₀₆₅ cells identified an about 45 kD protein.

25 D. Recombinant molecule pBv-nfSPI4₁₀₇₀, containing a flea serine protease inhibitor nucleic acid molecule spanning nucleotides from about 84 through about 1153 of SEQ ID NO:19, operatively linked to baculovirus polyhedron transcription control sequences were produced in the following manner. A PCR fragment of 1070 nucleotides, herein denoted nfSPI4₁₀₇₀, having SEQ ID NO:81 was amplified from
 30 nfSPI4₁₄₁₄ using the sense primer Serpin2For described above and the antisense primer

Serpin4Rev, having the nucleic acid sequence 5' - CGC **AGA** TCT TTA TTC AGT TGT TGG TTT AAC AAG ACG ACC -3' (SEQ ID NO:82; *Bgl*III site shown in bold).

The resulting 1070-bp PCR product (referred to as Bv-nfSPI4₁₀₇₀) was digested with *Bgl*III and *Eco*RI restriction endonucleases and subcloned into unique *Bgl*III and
5 *Eco*RI sites of pAcGP67 baculovirus shuttle plasmid to produce the recombinant molecule referred to herein as pAcG-nfSPI4₁₀₇₀.

The resultant recombinant molecule pAcG-nfSPI4₁₀₇₀ was verified for proper insert orientation by restriction mapping. Such a recombinant molecule can be co-transfected with a linear Baculogold baculovirus DNA into *S. frugiperda* Sf9 cells to
10 form the recombinant cells denoted *S. frugiperda*:pAcG-nfSPI4₁₀₇₀. *S. frugiperda*:pAcG-nfSPI4₁₀₇₀ was cultured in order to produce a flea serine protease inhibitor protein PfSPI4₃₅₆ (referred to herein as SEQ ID NO:98).

An immunoblot of supernatant from cultures of *S. frugiperda*:pAcG-nfSPI4₁₀₇₀ cells producing the flea serine protease inhibitor protein PfSPI4₃₅₆ was performed using
15 the anti-SPI2 polyclonal antiserum described in detail in Example 5. Blots were incubated using serum samples from the pre-bleed or from serum collected 14 days after the first boost of the rabbit. Analysis of the supernatant from cultures of *S. frugiperda*:pAcG-nfSPI4₁₀₇₀ cells identified an about 41 kD protein.

Example 12

20 This example describes the purification of serine protease inhibitor proteins from wandering larvae.

About 15,000 bovine blood-fed wandering larvae were homogenized in Tris buffered saline (TBS), pH 8 by sonication in 50 ml Oak Ridge centrifuge tubes (available from Nalgene Co., Rochester, NY) by sonicating 4 times 30 seconds each at a
25 setting of 5 of a model W-380 Sonicator (available from Heat Systems-Ultrasonics, Inc.). The sonicates were clarified by centrifugation at 27,000 x g for 30 minutes to produce an extract. Soluble protein in the extract was removed by aspiration and diluted to a volume of about 15 ml in TBS. Sodium chloride (NaCl) was then added to the extract to bring the final concentration of NaCl to about 400 mM. The extract was then
30 applied to a column containing about 2 ml of *p*-aminobenzamidine cross-linked to Sepharose® beads (available from Sigma, St. Louis, MO), previously equilibrated in 50

- mM Tris, pH 8, 400 mM NaCl, and incubated overnight. The unbound serine protease inhibitor proteins were then drained from the column and dialyzed against 2 changes of about 1 liter of 10 mM phosphate buffer, pH 7.2, 10 mM NaCl. Two aliquots of about 9 ml each were applied to a chromatography column containing about 10 ml of Macro-
- 5 Prep Ceramic Hydroxyapatite, Type I, 20 μ m beads (available from Bio-Rad Laboratories, Hercules, CA), previously equilibrated with 10 mM phosphate buffer, pH 7.2 containing 10 mM NaCl. The column was washed with 10 mM phosphate buffer, pH 7.2 containing 10 mM NaCl until all unbound protein was removed. Protein bound to the column was then eluted with a linear gradient from 10 mM phosphate buffer, pH
- 10 7.2 containing 10 mM NaCl to 0.5 M phosphate buffer, pH 6.5 containing 10 mM NaCl. Fractions were assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the rabbit anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that serine protease inhibitor proteins were eluted at about 120 mM phosphate.
- 15 The fractions that contained the most serine protease inhibitor proteins were combined and diafiltered into about 25 ml of 25 mM Tris (pH 8), 10 mM NaCl, in preparation for anion exchange chromatography. The sample was then applied to a Uno Q6 anion exchange column (available from Bio-Rad). The column was washed with 25 mM Tris (pH 8), 10 mM NaCl until all unbound protein was removed. Protein bound to
- 20 the column was then eluted with a linear gradient from 10 mM to 1 M NaCl in 25 mM Tris, pH 8. Fractions were assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that the serine protease inhibitor proteins were eluted at about 260 mM NaCl.
- 25 Fractions containing the most serine protease inhibitor proteins were pooled and diafiltered into a total volume of about 6 ml of 20 mM MES buffer (2-(N-morpholino)ethanesulfonic acid), pH 6, containing 10 mM NaCl, in preparation for cation exchange chromatography. The sample was then applied to an Uno S1 cation exchange column (available from Bio-Rad) equilibrated in MES buffer containing 10
- 30 mM NaCl. The column was washed with MES buffer containing 10 mM NaCl until all unbound protein was removed. Protein bound to the column was then eluted with a

linear gradient from 10 mM to 1 M NaCl in 20 mM MES buffer, pH 6 and fractions were collected. The fractions were assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that serine protease inhibitor proteins were not
5 retained on the cation exchange column using the above conditions, and most of the serine protease inhibitor proteins were found in the flow-through fractions.

The cation exchange fractions containing the most serine protease inhibitor proteins were combined and concentrated to about 400 μ l using an Ultrafree-20 15 ml centrifugal concentrator (available from Millipore Corp, Bedford, MA) in preparation
10 for size exclusion chromatography. The sample was applied to a Bio-Select SEC 125-5 size exclusion chromatography column (available from Bio-Rad), previously equilibrated in TBS, pH 7.2. The column was eluted with TBS, pH 7.2 at a flow rate of about 0.5 ml/min, and fractions of about 250 μ l were collected. Fractions were assayed
15 for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that serine protease inhibitor proteins were eluted in about 7 ml of buffer, corresponding to a molecular weight of about 30 kD to 66 kD based on the elution volumes of gel filtration molecular weight standard proteins (available from Sigma, St. Louis, MO).

The size exclusion chromatography fractions that contained the most serine
20 protease inhibitor proteins were combined and brought to about 40% saturation with ammonium sulfate in preparation for hydrophobic interaction chromatography. The sample was applied to a 1 ml HighTrap™ Phenyl Sepharose® HP hydrophobic interaction chromatography column (available from Pharmacia) equilibrated with TBS, 40% saturated with ammonium sulfate. The column was washed with TBS, 40%
25 saturated with ammonium sulfate until all unbound protein was removed. Bound protein was eluted from the column with a linear gradient from TBS, 40% saturated with ammonium sulfate to TBS with no ammonium sulfate. Fractions were assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that serine
30 protease inhibitor proteins were eluted when the buffer was about 30% saturated with ammonium sulfate.

The hydrophobic interaction chromatography fractions that contained the most serine protease inhibitor proteins were combined and assayed for protein concentration using Micro BCA Protein Assay Reagent (available from Pierce, Rockford, IL) with bovine serum albumin as a standard. About 10 µg of serine protease inhibitor proteins were concentrated to about 20 µl using a Microcon 3 centrifugal concentrator (available from Amicon, Beverly, MA), resolved on a reducing 14% SDS-PAGE gel (available from Novex, San Diego, CA) and then blotted onto a polyvinylidene difluoride (PVDF) membrane (available from Applied Biosystems, Foster City, CA) for about 60 min in 10 mM CAPS buffer (3-[cyclohexylamino]-1-propanesulfonic acid; available from Sigma, St. Louis, MO), pH 11, with 0.5 mM dithiothreitol (DTT). The membrane was stained for 1 minute in 0.1% Coomassie Blue R-250 dissolved in 40% methanol and 1% acetic acid. The membrane was destained in 50% methanol for about 10 minutes, rinsed with water and air dried. A stained protein band was identified having an apparent molecular weight identical to the proteins identified by the immunoblot method described above, at about 36 kD. A portion of the membrane containing the band was excised, and protein contained in the membrane segment was subjected to N-terminal amino sequencing using a 473A Protein Sequencer (available from Applied Biosystems) and using standard techniques. The results indicated that the N-terminal amino acid sequence of the 36 kD protein was Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met (using standard 3 letter amino acid code), referred to herein as SEQ ID NO:88.

Example 13

This example describes the purification of serine protease inhibitor proteins from cat blood fed adult flea midguts.

About 45,000 cat blood-fed wandering larvae were homogenized by freeze-fracture and sonicated in Tris buffer comprising 50 mM Tris, pH 8 and 100 mM CaCl₂. The sonicates were clarified by centrifugation at about 14,000 x g for 20 min to produce an extract. Soluble protein in the extract was removed by aspiration and diluted to a volume of about 45 ml in Tris buffer. Sodium chloride was then added to the extract to bring the final concentration of NaCl to about 400 mM. The extract was then applied in two aliquots to a column containing about 1 ml of *p*-aminobenzamidine cross-linked to

Sepharose® beads, previously equilibrated in 50 mM Tris, pH 8, 400 mM NaCl. After an overnight incubation, the columns were drained and the flow-through fractions were retained. The flow-through fractions, which contained most of the midgut proteins except serine proteases, were combined and diafiltered into about 16 ml of 25 mM Tris, pH 8, containing 10 mM NaCl in preparation for anion exchange chromatography. Two aliquots of about 8 ml were then applied to an Uno Q6 column and fractions assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that the serine protease inhibitor proteins were eluted at about 160 mM NaCl.

10 The anion exchange column fractions that contained the most serine protease inhibitor proteins were pooled and diafiltered into a total of about 3 ml of 20 mM MES buffer, pH 6, containing 10 mM NaCl in preparation for cation exchange chromatography. The sample was then applied to an Uno S1 column and fractions assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that serine protease inhibitor proteins were not retained on the cation exchange column using the above conditions, and most of the serine protease inhibitor proteins were found in the flow-through fractions.

20 The cation exchange fractions that contained the most serine protease inhibitor proteins were combined and diafiltered into about 3 ml of 25 mM Tris, pH 8, containing 10 mM NaCl in preparation for anion exchange chromatography. The sample was applied to a Bio-Scale Q2 column (available from Bio-Rad), previously equilibrated in 25 mM Tris, pH 8, containing 10 mM NaCl. The column was washed with 25 mM Tris, pH 8, 10 mM NaCl until all unbound protein was removed. Protein bound to the column was then eluted with a linear gradient from 10 mM to 1 M NaCl in 25 mM Tris, pH 8. Fractions were assayed for the presence of serine protease inhibitor proteins by immunoblot analysis using the anti-SPI2 polyclonal antiserum described in Example 5. The results indicated that serine protease inhibitor proteins were eluted at about 140 mM NaCl.

30 About 500 µl of the anion exchange column fraction that contained the most serine protease inhibitor protein was concentrated to about 25 µl using a Microcon 3

centrifugal concentrator (available from Amicon, Beverly, MA), and then separated by SDS-PAGE, electroblotted onto a PVDF membrane, and two stained protein bands, at about 35 kD and 36 kD, were N-terminally sequenced as described in Example 12. The results indicated that the N-terminal amino acid sequence of the 35 kD protein was Ser

5 Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn
Leu Ile Met (using standard 3 letter amino acid code; referred to herein as SEQ ID
NO:89) and the N-term sequence of the 36 kD protein was Ser Thr Ser Ile Asn Gln Phe
Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser Pro
(using standard 3 letter amino acid code; referred to herein as SEQ ID NO:90).

10 Example 14

This example describes the identification of serine protease inhibitor proteins in different flea tissues.

Tissue samples were isolated from unfed or bovine blood-fed 1st instar
Ctenocephalides felis flea larvae; bovine blood-fed 3rd instar *C. felis* flea larvae, bovine
15 blood-fed wandering *C. felis* flea larvae, unfed or cat blood-fed adult *C. felis* flea midgut
tissue, cat blood-fed adult *C. felis* flea tissues that had their midguts and heads removed
(adult partial fleas), and whole unfed or cat blood-fed adult *C. felis* fleas. The 1st instar,
3rd instar, wandering and adult midgut tissues were then homogenized by freeze-fracture
and sonicated in Tris buffered saline (TBS). The adult partial fleas and adult whole fleas
20 were then homogenized by freeze-fracture and ground with a microtube mortar and
pestle. The extracts were centrifuged at about 14,000 x g for 20 min and the soluble
material recovered. The soluble material was then diluted to a final concentration of
about 1 tissue equivalent per 2 μ l. Each soluble extract sample was then assayed for the
presence of serine protease inhibitor proteins by immunoblot analysis using the anti-
25 SPI2 polyclonal antiserum described in Example 5.

The results shown in Figure 1 indicated that all tissue extracts except the unfed
1st instar tissues contained proteins of about 25 kD to 97 kD that were cross reactive with
the rabbit anti-SPI2 polyclonal antiserum, and were therefore comprised at least partially
of serine protease inhibitor proteins.

-94-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Wisnewski, Nancy
Brandt, Kevin S.
Silver, Gary M.
Maddux, Joely D.
- 10 (ii) TITLE OF INVENTION: Novel Serine Protease
Inhibitor Nucleic Acid
Molecules, Proteins and
Uses Thereof
- (iii) NUMBER OF SEQUENCES: 98
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Lahive & Cockfield, LLP
(B) STREET: 28 State Street
(C) CITY: Boston
(D) STATE: Massachusetts
(E) COUNTRY: USA
(F) ZIP: 02109
- 20 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: Windows 95
(D) SOFTWARE: WordPerfect for Windows, Version 7.0
- 25 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:
- 30 (vii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Rothenberger, Scott D.
(B) REGISTRATION NUMBER: 41,277
(C) REFERENCE/DOCKET NUMBER: HKV-011PC
- 35 (viii) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (617) 227-7400
(B) TELEFAX: (617) 742-4214

(2) INFORMATION FOR SEQ ID NO:1:

- 40 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1584 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA

-95-

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 136..1326

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1

5	GCCTGGAAGG TGATAAGTAA ACGGGCACGG TAGTGTTTGT TTTTAGAAAA TAATTTTAAT	60
	TCGTACGACG TACGTTTTTG TGATTTTAAT TTTTGTAGCT TTTTGTAGCT CTGAAAGAGC	120
	CGAAATTTTA GCAAA ATG ATT AAC GCA CGA CTT GTG TTT CTT TTT GTA TCA	171
	Met Ile Asn Ala Arg Leu Val Phe Leu Phe Val Ser	
	1 5 10	
10	GTG TTA TTA CCA ATT TCA ACA ATG GCC GAT CCC CAG GAA TTG TCT ACA	219
	Val Leu Leu Pro Ile Ser Thr Met Ala Asp Pro Gln Glu Leu Ser Thr	
	15 20 25	
	AGT ATT AAC CAG TTT GCT GGA AGC CTG TAC AAT ACA GTT GCT TCT GGC	267
	Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly	
15	30 35 40	
	AAC AAA GAC AAT CTC ATC ATG TCC CCA TTG TCT GTA CAA ACT GTT CTA	315
	Asn Lys Asp Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu	
	45 50 55 60	
20	TCC CTG GTG TCA ATG GGA GCT GGT GGC AAT ACT GCC ACA CAA ATA GCT	363
	Ser Leu Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile Ala	
	65 70 75	
	GCT GGT TTG CGT CAG CCT CAA TCA AAA GAA AAA ATT CAA GAT GAC TAC	411
	Ala Gly Leu Arg Gln Pro Gln Ser Lys Glu Lys Ile Gln Asp Asp Tyr	
	80 85 90	
25	CAC GCA TTG ATG AAC ACT CTT AAT ACA CAA AAA GGT GTA ACT CTG GAA	459
	His Ala Leu Met Asn Thr Leu Asn Thr Gln Lys Gly Val Thr Leu Glu	
	95 100 105	
	ATT GCC AAT AAA GTT TAT GTT ATG GAA GGC TAT ACA TTA AAA CCC ACC	507
	Ile Ala Asn Lys Val Tyr Val Met Glu Gly Tyr Thr Leu Lys Pro Thr	
30	110 115 120	
	TTC AAA GAA GTT GCC ACC AAC AAA TTC TTA GCT GGA GCA GAA AAC TTG	555
	Phe Lys Glu Val Ala Thr Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu	
	125 130 135 140	
	AAC TTT GCC CAA AAT GCT GAA AGC GCT AAA GTT ATC AAC ACT TGG GTT	603
35	Asn Phe Ala Gln Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val	
	145 150 155	
	GAA GAA AAA ACT CAT GAC AAA ATT CAT GAT TTG ATC AAA GCC GGT GAT	651
	Glu Glu Lys Thr His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp	
	160 165 170	
40	CTA GAC CAG GAT TCA AGA ATG GTT CTT GTC AAT GCA TTG TAC TTC AAG	699
	Leu Asp Gln Asp Ser Arg Met Val Leu Val Asn Ala Leu Tyr Phe Lys	
	175 180 185	
	GGT CTT TGG GAG AAA CAA TTC AAA AAG GAA AAT ACC CAA GAC AAA CCT	747
	Gly Leu Trp Glu Lys Gln Phe Lys Lys Glu Asn Thr Gln Asp Lys Pro	
45	190 195 200	

-96-

	TTC	TAT	GTT	ACT	GAA	ACA	GAG	ACA	AAG	AAT	GTA	CGA	ATG	ATG	CAC	ATT	795
	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg	Met	Met	His	Ile	
	205					210					215					220	
5	AAG	GAT	AAA	TTC	CGT	TAT	GGA	GAA	TTT	GAA	GAA	TTA	GAT	GCC	AAG	GCT	843
	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu	Glu	Leu	Asp	Ala	Lys	Ala	
					225					230					235		
	GTA	GAA	TTG	CCC	TAC	AGG	AAC	TCA	GAT	TTG	GCC	ATG	TTA	ATC	ATT	TTG	891
	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	
				240					245					250			
10	CCA	AAC	AGC	AAA	ACT	GGT	CTC	CCC	GCT	CTT	GAA	GAA	AAA	TTA	CAA	AAT	939
	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	
				255				260					265				
	GTT	GAT	TTG	CAA	AAC	TTG	ACT	CAA	CGC	ATG	TAC	TCT	GTT	GAA	GTT	ATT	987
15	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	
		270					275					280					
	TTG	GAT	CTG	CCT	AAA	TTC	AAG	ATT	GAA	TCT	GAA	ATT	AAT	TTG	AAT	GAT	1035
	Leu	Asp	Leu	Pro	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	
		285				290					295					300	
	CCT	CTG	AAA	AAG	TTG	GGT	ATG	TCT	GAT	ATG	TTT	GTT	CCT	GGA	AAA	GCT	1083
20	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	Met	Phe	Val	Pro	Gly	Lys	Ala	
					305					310					315		
	GAT	TTC	AAA	GGA	TTG	CTT	GAA	GGA	TCT	GAT	GAG	ATG	TTA	TAT	ATT	TCT	1131
	Asp	Phe	Lys	Gly	Leu	Leu	Glu	Gly	Ser	Asp	Glu	Met	Leu	Tyr	Ile	Ser	
				320					325					330			
25	AAA	GTA	ATT	CAA	AAA	GCT	TTC	ATT	GAA	GTA	AAT	GAA	GAA	GGT	GCT	GAA	1179
	Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile	Glu	Val	Asn	Glu	Glu	Gly	Ala	Glu	
				335				340					345				
	GCT	GCA	GCT	GCC	ACA	GCT	ACC	TTT	ATG	GTT	ACC	TAT	GAA	CTG	GAG	GTT	1227
30	Ala	Ala	Ala	Ala	Thr	Ala	Thr	Phe	Met	Val	Thr	Tyr	Glu	Leu	Glu	Val	
		350					355					360					
	TCC	CTG	GAT	CTT	CCC	ACT	GTT	TTT	AAA	GTC	GAT	CAT	CCA	TTC	AAT	ATT	1275
	Ser	Leu	Asp	Leu	Pro	Thr	Val	Phe	Lys	Val	Asp	His	Pro	Phe	Asn	Ile	
		365				370					375					380	
	GTT	TTG	AAG	ACA	GGT	GAT	ACT	GTT	ATT	TTT	AAT	GGG	CGA	GTT	CAA	ACT	1323
35	Val	Leu	Lys	Thr	Gly	Asp	Thr	Val	Ile	Phe	Asn	Gly	Arg	Val	Gln	Thr	
					385					390					395		
	TTA	TAA	AATGGATAGT	GTAAAAAGAA	TACAAGATCT	ATCTGAATCT	CTGGATTAAT										1379
	Leu																
40	GAAGTAATTTT	TTCTACAATA	TTTTTTTAATA	GTTATTAGGT	CTAAAAATAAG	TTTATTTTTTTT											1439
	AGTATGTGGT	ATAAATCGTG	TAGACGAAAA	ATGTTTTTGT	TTAGTTTTTCA	CTTTTTTATGA											1499
	ATGTAATCAC	CTATATAATG	TTGTAGTTTA	TGTAATAAAA	ATGTTAAATG	TGAAAAAAA											1559
	AAAAAAAAAA	AAAAAAAAAA	AAAAA														1584

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- 45 (A) LENGTH: 397 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

-97-

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Ile	Asn	Ala	Arg	Leu	Val	Phe	Leu	Phe	Val	Ser	Val	Leu	Leu	Pro	1	5	10	15
5	Ile	Ser	Thr	Met	Ala	Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	20	25	30
	Phe	Ala	Gly	Ser	Leu	Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	35	40	45
10	Leu	Ile	Met	Ser	Pro	Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	50	55	60
	Met	Gly	Ala	Gly	Gly	Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	65	70	75
	Gln	Pro	Gln	Ser	Lys	Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	85	90	95
15	Asn	Thr	Leu	Asn	Thr	Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	100	105	110
	Val	Tyr	Val	Met	Glu	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	115	120	125
20	Ala	Thr	Asn	Lys	Phe	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	130	135	140
	Asn	Ala	Glu	Ser	Ala	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	145	150	155
	His	Asp	Lys	Ile	His	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	165	170	175
25	Ser	Arg	Met	Val	Leu	Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	180	185	190
	Lys	Gln	Phe	Lys	Lys	Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	195	200	205
30	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	210	215	220
	Arg	Tyr	Gly	Glu	Phe	Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	225	230	235
	Tyr	Arg	Asn	Ser	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	245	250	255
35	Thr	Gly	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	260	265	270
	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp	Leu	Pro	275	280	285
40	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	Pro	Leu	Lys	Lys	290	295	300

-98-

Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly
 305 310 315 320
 Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln
 325 330 335
 5 Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu Ala Ala Ala
 340 345 350
 Thr Ala Thr Phe Met Val Thr Tyr Glu Leu Glu Val Ser Leu Asp Leu
 355 360 365
 10 Pro Thr Val Phe Lys Val Asp His Pro Phe Asn Ile Val Leu Lys Thr
 370 375 380
 Gly Asp Thr Val Ile Phe Asn Gly Arg Val Gln Thr Leu
 385 390 395

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 15 (A) LENGTH: 1584 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA
 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	TTTCACATTT	AACATTTTTTA	TTACATAAAC	60
TACAACATTA	TATAGGTGAT	TACATTCATA	AAAAGTGAAA	ACTAAAACAA	AACATTTTTTC	120
GTCTACACGA	TTTATACCAC	ATACTAAAAA	ATGAACCTTAT	TTTAGACCTA	ATAACTATTA	180
AAAAATATTG	TAGAAAAATT	ACTTCATTAA	TCCAGAGATT	CAGATAGATC	TTGTATTCTT	240
25 TTTACACTAT	CCATTTTATA	AAGTTTGAAC	TCGCCCATTAA	AAAATAACAG	TATCACCTGT	300
CTTCAAAACA	ATATTGAATG	GATGATCGAC	TTTAAAAACA	GTGGGAAGAT	CCAGGGAAAC	360
CTCCAGTTCA	TAGGTAACCA	TAAAGGTAGC	TGTGGCAGCT	GCAGCTTCAG	CACCTTCTTC	420
ATTTACTTCA	ATGAAAGCTT	TTTGAATTAC	TTTAGAAATA	TATAACATCT	CATCAGATCC	480
TTCAAGCAAT	CCTTTGAAAT	CAGCTTTTCC	AGGAACAAAC	ATATCAGACA	TACCCAACTT	540
30 TTTCAGAGGA	TCATTCAAAT	TAATTTTCTA	TTCAATCTTG	AATTTAGGCA	GATCCAAAAT	600
AACTTCAACA	GAGTACATGC	GTTGAGTCAA	GTTTTGCAAA	TCAACATTTT	GTAATTTTTC	660
TTCAAGAGCG	GGGAGACCAG	TTTTGCTGTT	TGGCAAAATG	ATTAACATGG	CCAAATCTGA	720
GTTCTGTAG	GGCAATTCTA	CAGCCTTGGC	ATCTAATTCT	TCAAATTCTC	CATAACGGAA	780
TTTATCCTTA	ATGTGCATCA	TTCGTACATT	CTTTGTCTCT	GTTTCAGTAA	CATAGAAAGG	840
35 TTTGTCTTGG	GTATTTTCCT	TTTTGAATTG	TTTCTCCCAA	AGACCCTTGA	AGTACAATGC	900
ATTGACAAGA	ACCATTCTTG	AATCCTGGTC	TAGATCACCG	GCTTTGATCA	AATCATGAAT	960
TTTGTCAATG	GTTTTTTCTT	CAACCCAAGT	GTTGATAACT	TTAGCGCTTT	CAGCATTTTG	1020
GGCAAAGTTC	AAGTTTTCTG	CTCCAGCTAA	GAATTTGTTG	GTGGCAACTT	CTTTGAAGGT	1080
GGGTTTTAAT	GTATAGCCTT	CCATAACATA	AACTTTATTG	GCAATTTCCA	GAGTTACACC	1140
40 TTTTGTGTGA	TTAAGAGTGT	TCATCAATGC	GTGGTAGTCA	TCTTGAATTT	TTTCTTTTGA	1200
TTGAGGTGTA	CGCAAACCAG	CAGCTATTTG	TGTGGCAGTA	TTGCCACCAG	CTCCCATTTG	1260
CACCAGGGAT	AGAACAGTTT	GTACAGACAA	TGGGGACATG	ATGAGATTGT	CTTTGTTGCC	1320
AGAAGCAACT	GTATTGTACA	GGCTTCCAGC	AAACTGGTTA	ATACTTGTAG	ACAATTCCCTG	1380
GGGATCGGCC	ATTGTTGAAA	TTGGTAATAA	CACTGATACA	AAAAGAAACA	CAAGTCGTGC	1440
45 GTTAATCATT	TTGCTAAAAT	TTCGGCTCTT	TCAGAGCTAC	AAAAACACTA	AAAAATTAAA	1500
ATCACAAAAA	CGTACGTCGT	ACGAATTAAA	ATTATTTTCT	AAAACAAAAC	ACTACCGTGC	1560
CCGTTTACTT	ATCACCTTCC	AGGC				1584

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1191 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

	ATGATTAACG	CACGACTTGT	GTTTCTTTTT	GTATCAGTGT	TATTACCAAT	TTCAACAATG	60
10	GCCGATCCCC	AGGAATTGTC	TACAAGTATT	AACCAGTTTG	CTGGAAGCCT	GTACAATACA	120
	GTTGCTTCTG	GCAACAAAGA	CAATCTCATC	ATGTCCCCAT	TGCTCTGTACA	AACTGTTCTA	180
	TCCCTGGTGT	CAATGGGAGC	TGGTGGCAAT	ACTGCCACAC	AAATAGCTGC	TGGTTTGCCT	240
	CAGCCTCAAT	CAAAAGAAAA	AATTCAAGAT	GACTACCACG	CATTGATGAA	CACTCTTAAT	300
	ACACAAAAAG	GTGTAACCTC	GGAAATTGCC	AATAAAGTTT	ATGTTATGGA	AGGCTATACA	360
15	TTAAAACCCA	CCTTCAAAGA	AGTTGCCACC	AACAAATTCT	TAGCTGGAGC	AGAAAACCTG	420
	AACTTTGCCC	AAAATGCTGA	AAGCGCTAAA	GTTATCAACA	CTGGGGTTGA	AGAAAAAACT	480
	CATGACAAAA	TTCATGATTT	GATCAAAGCC	GGTGATCTAG	ACCAGGATTC	AAGAATGGTT	540
	CTTGTCAATG	CATTGTACTT	CAAGGGTCTT	TGGGAGAAAC	AATTCAAAAA	GGAAAATACC	600
	CAAGACAAAC	CTTTCTATGT	TACTGAAACA	GAGACAAAGA	ATGTACGAAT	GATGCACATT	660
20	AAGGATAAAT	TCCGTTATGG	AGAATTTGAA	GAATTAGATG	CCAAGGCTGT	AGAATTGCCC	720
	TACAGGAAC	CAGATTTGGC	CATGTTAATC	ATTTTGCCAA	ACAGCAAAAC	TGGTCTCCCC	780
	GCTCTTGAAG	AAAAATTACA	AAATGTTGAT	TTGCAAAACT	TGACTCAACG	CATGTACTCT	840
	GTTGAAGTTA	TTTTGGATCT	GCCTAAATTC	AAGATTGAAT	CTGAAATTAA	TTTGAATGAT	900
	CCTCTGAAAA	AGTTGGGTAT	GTCTGATATG	TTTGTTCCTG	GAAAAGCTGA	TTTCAAAGGA	960
25	TTGCTTGAAG	GATCTGATGA	GATGTTATAT	ATTTCTAAAG	TAATTCAAAA	AGCTTTCATT	1020
	GAAGTAAATG	AAGAAGGTGC	TGAAGCTGCA	GCTGCCACAG	CTACCTTTAT	GGTTACCTAT	1080
	GAAGTGGAGG	TTTCCCTGGA	TCTTCCCACT	GT'TTTTAAAG	TCGATCATCC	ATTCAATATT	1140
	GTTTTGAAGA	CAGGTGATAC	TGTTATTTTT	AATGGGCGAG	TTCAAACCTT	A	1191

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1191 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

	TAAAGTTTGA	ACTCGCCCAT	TAAAAATAAC	AGTATCACCT	GTCTTCAAAA	CAATATTGAA	60
	TGGATGATCG	ACTTTAAAAA	CAGTGGGAAG	ATCCAGGGAA	ACCTCCAGTT	CATAGGTAAC	120
	CATAAAGGTA	GCTGTGGCAG	CTGCAGCTTC	AGCACCTTCT	TCATTTACTT	CAATGAAAGC	180
40	TTTTTGAATT	ACTTTAGAAA	TATATAACAT	CTCATCAGAT	CCTTCAAGCA	ATCCTTTGAA	240
	ATCAGCTTTT	CCAGGAACAA	ACATATCAGA	CATACCCAAC	TTTTTCAGAG	GATCATTCAA	300
	ATTAATTTCA	GATTCAATCT	TGAATTTAGG	CAGATCCAAA	ATAACTTCAA	CAGAGTACAT	360
	GCGTTGAGTC	AAGTTTTGCA	AATCAACATT	TTGTAATTTT	TCTTCAAGAG	CGGGGAGACC	420
	AGTTTTGCTG	TTTGGCAAAA	TGATTAACAT	GGCCAAATCT	GAGTTCCTGT	AGGGCAATTC	480
45	TACAGCCTTG	GCATCTAATT	CTTCAAATTC	TCCATAACGG	AATTTATCCT	TAATGTGCAT	540
	CATTCGTACA	TTCTTTGTCT	CTGTTTCAGT	AACATAGAAA	GGTTTGTCTT	GGGTATTTTC	600
	CTTTTGAAT	TGTTTCTCCC	AAAGACCCTT	GAAGTACAAT	GCATTGACAA	GAACCATTCT	660
	TGAATCCTGG	TCTAGATCAC	CGGCTTTGAT	CAAATCATGA	ATTTTGTTCAT	GAGTTTTTTC	720
	TTCAACCCAA	GTGTTGATAA	CTTTAGCGCT	TTCAGCATTT	TGGGCAAAGT	TCAAGTTTTTC	780
50	TGCTCCAGCT	AAGAATTTGT	TGGTGGCAAC	TTCTTTGAAG	GTGGGTTTTA	ATGTATAGCC	840
	TTCCATAACA	TAAACTTTAT	TGGCAATTTT	CAGAGTTACA	CCTTTTGTGT	TATTAAGAGT	900
	GTTTCATCAAT	GCGTGGTAGT	CATCTTGAAT	TTTTTCTTTT	GATTGAGGCT	GACGCAAACC	960
	AGCAGCTATT	TGTGTGGCAG	TATTGCCACC	AGCTCCCAT	GACACCAGGG	ATAGAACAGT	1020

-100-

TTGTACAGAC AATGGGGACA TGATGAGATT GTCTTTGTTG CCAGAAGCAA CTGTATTGTA 1080
 CAGGCTTCCA GCAAACCTGGT TAATACTTGT AGACAATTCC TGGGGATCGG CCATTGTTGA 1140
 AATTGGTAAT AACACTGATA CAAAAAGAAA CACAAGTCGT GCGTTAATCA T 1191

(2) INFORMATION FOR SEQ ID NO:6:

5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 376 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu	1	5	10	15
Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro	20	25	30	
Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	35	40	45	
Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	50	55	60	
Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	65	70	75	80
Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	85	90	95	
Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	100	105	110	
Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	115	120	125	
Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	130	135	140	
Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	145	150	155	160
Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	165	170	175	
Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	180	185	190	
Asn	Val	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	195	200	205	
Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	210	215	220	
Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	225	230	235	240
Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	245	250	255	

-101-

	Met	Tyr	Ser	Val 260	Glu	Val	Ile	Leu	Asp 265	Leu	Pro	Lys	Phe	Lys 270	Ile	Glu
	Ser	Glu	Ile 275	Asn	Leu	Asn	Asp	Pro 280	Leu	Lys	Lys	Leu	Gly 285	Met	Ser	Asp
5	Met	Phe 290	Val	Pro	Gly	Lys	Ala 295	Asp	Phe	Lys	Gly	Leu 300	Leu	Glu	Gly	Ser
	Asp 305	Glu	Met	Leu	Tyr	Ile 310	Ser	Lys	Val	Ile	Gln 315	Lys	Ala	Phe	Ile	Glu 320
10	Val	Asn	Glu	Glu	Gly 325	Ala	Glu	Ala	Ala	Ala 330	Ala	Thr	Ala	Thr	Phe 335	Met
	Val	Thr	Tyr	Glu 340	Leu	Glu	Val	Ser	Leu 345	Asp	Leu	Pro	Thr	Val 350	Phe	Lys
	Val	Asp	His 355	Pro	Phe	Asn	Ile	Val 360	Leu	Lys	Thr	Gly	Asp 365	Thr	Val	Ile
15	Phe	Asn 370	Gly	Arg	Val	Gln	Thr 375	Leu								

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1358 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

```

25      (ix)  FEATURE:
          (A)  NAME/KEY:  CDS
          (B)  LOCATION:  2..1198

```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

	C	GCG	ATA	GTT	CAA	CAC	GCA	CGA	CTT	GTG	TTT	CTT	TTT	GTA	TCA	GTG	46
	Ala	Ile	Val	Gln	His	Ala	Arg	Leu	Val	Phe	Leu	Phe	Val	Ser	Val		
30	1					5				10					15		
	TTA	ATA	CCA	ATT	TCA	ACA	ATG	GCG	GAT	CCC	CAG	GAA	TTG	TCT	ACA	AGT	94
	Leu	Ile	Pro	Ile	Ser	Thr	Met	Ala	Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	
					20					25					30		
	ATT	AAC	CAG	TTT	GCT	GGA	AGC	CTG	TAC	AAT	ACG	GTT	GCT	TCT	GGC	AAC	142
35	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu	Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	
				35					40					45			
	AAA	GAC	AAT	CTC	ATC	ATG	TCC	CCA	TTG	TCT	GTA	CAA	ACT	GTT	CTA	TCC	190
	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro	Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	
			50					55					60				
	CTG	GTG	TCA	ATG	GGA	GCT	GGT	GGT	AAT	ACT	GCC	ACA	CAA	ATA	GCT	GCT	238
40	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	
		65					70					75					

-102-

	GGT	TTA	CGT	CAG	CCT	CAA	TCA	AAA	GAA	AAA	ATT	CAA	GAT	GAC	TAC	CAT	286
	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	
	80					85					90					95	
5	GCA	TTG	ATG	AAC	ACT	CTT	AAT	ACA	CAA	AAA	GGT	GTA	ACT	CTG	GAA	ATT	334
	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	
					100					105					110		
	GCC	AAC	AAA	GTT	TAC	GTT	ATG	GAA	GGC	TAT	ACA	TTG	AAA	CCC	ACC	TTC	382
	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	
				115					120					125			
10	AAA	GAA	GTT	GCC	ACC	AAC	AAA	TTC	TTA	GCT	GGA	GCA	GAA	AAC	TTG	AAC	430
	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	
			130					135						140			
	TTT	GCC	CAA	AAT	GCT	GAA	AGC	GCT	AAA	GTT	ATC	AAC	ACT	TGG	GTT	GAA	478
15	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	
		145						150					155				
	GAA	AAA	ACT	CAT	GAC	AAA	ATT	CAT	GAT	TTG	ATC	AAA	GCC	GGT	GAT	CTA	526
	Glu	Lys	Thr	His	Asp	Lys	Ile	His	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	
	160					165					170					175	
20	GAC	CAG	GAT	TCA	AGA	ATG	GTT	CTT	GTC	AAT	GCA	TTG	TAC	TTC	AAG	GGT	574
	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	
					180					185					190		
	CTT	TGG	GAG	AAA	CAA	TTC	AAG	AAG	GAA	AAC	ACT	CAA	GAC	AAA	CCT	TTC	622
	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	
				195					200					205			
25	TAT	GTT	ACT	GAA	ACA	GAG	ACA	AAG	AAT	GTA	CGA	ATG	ATG	CAC	ATT	AAG	670
	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg	Met	Met	His	Ile	Lys	
			210					215						220			
	GAT	AAA	TTC	CGT	TAT	GGA	GAA	TTT	GAA	GAA	TTA	GAT	GCC	AAG	GCT	GTA	718
30	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	
		225					230					235					
	GAA	TTG	CCC	TAC	AGG	AAC	TCA	GAT	TTG	GCC	ATG	TTA	ATC	ATT	TTG	CCA	766
	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	
	240					245					250					255	
35	AAC	AGC	AAA	ACT	GGT	CTC	CCC	GCT	CTT	GAA	GAA	AAA	TTA	CAA	AAT	GTT	814
	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	
					260					265					270		
	GAC	TTG	CAA	AAC	TTG	ACT	CAA	CGC	ATG	TAC	TCT	GTT	GAA	GTT	ATT	TTG	862
	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	
				275					280					285			
40	GAT	CTG	CCT	AAA	TTC	AAG	ATT	GAA	TCT	GAA	ATT	AAT	TTG	AAT	GAT	CCT	910
	Asp	Leu	Pro	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	Pro	
			290					295					300				
45	CTG	AAA	AAG	TTG	GGT	ATG	TCT	GAT	ATG	TTT	GTT	CCT	GGA	AAA	GCT	GAT	958
	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	Met	Phe	Val	Pro	Gly	Lys	Ala	Asp	
		305					310					315					

-103-

```

TTC AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT AAA      1006
Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys
320                               325                               330                               335

GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT GCT GAA GCT      1054
5 Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu Ala
                               340                               345                               350

GCA GCT GCC ACA GGC ATT GTC ATG CTT GGT TGC TGT ATG CCA ATG ATG      1102
Ala Ala Ala Thr Gly Ile Val Met Leu Gly Cys Cys Met Pro Met Met
                               355                               360                               365

10 GAT CTT TCT CCA GTA GTT TTT AAT ATT GAT CAC CCA TTT TAT TAC TCA      1150
Asp Leu Ser Pro Val Val Phe Asn Ile Asp His Pro Phe Tyr Tyr Ser
                               370                               375                               380

TTG ATG ACT TGG GAT ACT GTT TTG TTC AGT GGA TGT GTT AAA TCC CTT      1198
15 Leu Met Thr Trp Asp Thr Val Leu Phe Ser Gly Cys Val Lys Ser Leu
                               385                               390                               395

TAA ATTTCTTCTT AGAATGAAGG TATTTTCAGTG TCTAATGGCA TTGATAGACC      1251
CAAAAATTTT AATTCTGACC ATGCTTTCTA CCTCATGATA ACGGCAGGGA AAACGATTTC      1311
AATTAGAGGT CGTTTCTATA ACTCCTAGTA TATGTTATAT GACTAGT      1358

```

(2) INFORMATION FOR SEQ ID NO:8:

```

20 (i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 399 amino acids
    (B) TYPE: amino acid
    (D) TOPOLOGY: linear

    (ii) MOLECULE TYPE: protein

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Ala Ile Val Gln His Ala Arg Leu Val Phe Leu Phe Val Ser Val Leu
 1                               5                               10                               15

Ile Pro Ile Ser Thr Met Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile
                               20                               25                               30

30 Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys
   35                               40                               45

Asp Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu Ser Leu
   50                               55                               60

35 Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile Ala Ala Gly
   65                               70                               75                               80

Leu Arg Gln Pro Gln Ser Lys Glu Lys Ile Gln Asp Asp Tyr His Ala
                               85                               90                               95

Leu Met Asn Thr Leu Asn Thr Gln Lys Gly Val Thr Leu Glu Ile Ala
                               100                               105                               110

40 Asn Lys Val Tyr Val Met Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys
   115                               120                               125

Glu Val Ala Thr Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe
   130                               135                               140

```

-104-

Ala Gln Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu
 145 150 155 160

Lys Thr His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp Leu Asp
 165 170 175

5 Gln Asp Ser Arg Met Val Leu Val Asn Ala Leu Tyr Phe Lys Gly Leu
 180 185 190

Trp Glu Lys Gln Phe Lys Lys Glu Asn Thr Gln Asp Lys Pro Phe Tyr
 195 200 205

10 Val Thr Glu Thr Glu Thr Lys Asn Val Arg Met Met His Ile Lys Asp
 210 215 220

Lys Phe Arg Tyr Gly Glu Phe Glu Glu Leu Asp Ala Lys Ala Val Glu
 225 230 235 240

Leu Pro Tyr Arg Asn Ser Asp Leu Ala Met Leu Ile Ile Leu Pro Asn
 245 250 255

15 Ser Lys Thr Gly Leu Pro Ala Leu Glu Glu Lys Leu Gln Asn Val Asp
 260 265 270

Leu Gln Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp
 275 280 285

20 Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu
 290 295 300

Lys Lys Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe
 305 310 315 320

Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val
 325 330 335

25 Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu Ala Ala
 340 345 350

Ala Ala Thr Gly Ile Val Met Leu Gly Cys Cys Met Pro Met Met Asp
 355 360 365

30 Leu Ser Pro Val Val Phe Asn Ile Asp His Pro Phe Tyr Tyr Ser Leu
 370 375 380

Met Thr Trp Asp Thr Val Leu Phe Ser Gly Cys Val Lys Ser Leu
 385 390 395

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
 35 (A) LENGTH: 1358 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ACTAGTCATA TAACATATAC TAGGAGTTAT AGAAACGACC TCTAATTGAA ATCGTTTTCC 60
 CTGCCGTTAT CATGAGGTAG AAAGCATGGT CAGAATTGAA ATTTTGGGT CTATCAATGC 120

-105-

	CATTAGACAC	TGAAATACCT	TCATTCTAAG	AAGAAATTTA	AAGGGATTTA	ACACATCCAC	180
	TGAACAAAAC	AGTATCCCAA	GTCATCAATG	AGTAATAAAA	TGGGTGATCA	ATATTAAAAA	240
	CTACTGGAGA	AAGATCCATC	ATTGGCATAC	AGCAACCAAG	CATGACAATG	CCTGTGGCAG	300
	CTGCAGCTTC	AGCACCTTCT	TCATTTACTT	CAATGAAAGC	TTTTTGAATT	ACTTTAGAAA	360
5	TATATAACAT	CTCATCAGAT	CCTTCAAGCA	ATCCTTTGAA	ATCAGCTTTT	CCAGGAACAA	420
	ACATATCAGA	CATACCCAAC	TTTTTCAGAG	GATCATTTCA	ATTAATTTCA	GATTCAATCT	480
	TGAATTTAGG	CAGATCCAAA	ATAACTTCAA	CAGAGTACAT	GCGTTGAGTC	AAGTTTGTGA	540
	AGTCAACATT	TTGTAATTTT	TCTTCAAGAG	CGGGGAGACC	AGTTTTGCTG	TTTGGCAAAA	600
	TGATTAACAT	GGCCAAATCT	GAGTTCCTGT	AGGGCAATTC	TACAGCCTTG	GCATCTAATT	660
10	CTTCAAAATC	TCCATAACGG	AATTTATCCT	TAATGTGCAT	CATTTCGTACA	TTCTTTGTCT	720
	CTGTTTCAGT	AACATAGAAA	GGTTTGTCTT	GAGTGTTTTC	CTTCTTGAAT	TGTTTCTCCC	780
	AAAGACCCCT	GAAGTACAAT	GCATTGACAA	GAACCATTCT	TGAATCCTGG	TCTAGATCAC	840
	CGGCTTTGAT	CAAATCATGA	ATTTTGTTCAT	GAGTTTTTTC	TTCAACCCAA	GTGTGATAAA	900
	CTTTAGCGCT	TTCAGCATTT	TGGGCAAAGT	TCAAGTTTTT	TGCTCCAGCT	AAGAATTTGT	960
15	TGGTGGCAAC	TTCTTTGAAG	GTGGGTTTCA	ATGTATAGCC	TTCCATAACG	TAAACTTTGT	1020
	TGGCAATTTT	CAGAGTTACA	CCTTTTTGTG	TATTAAGAGT	GTTTCATCAAT	GCATGGTAGT	1080
	CATCTTGAAAT	TTTTTCTTTT	GATTGAGGCT	GACGTAAACC	AGCAGCTATT	TGTGTGTCAG	1140
	TATTACCACC	AGCTCCCATT	GACACCAGGG	ATAGAACAGT	TTGTACAGAC	AATGGGGACA	1200
	TGATGAGATT	GTCTTTGTTG	CCAGAAGCAA	CCGTATTGTA	CAGGCTTCCA	GCAAACGGT	1260
20	TAATACTTGT	AGACAATTCC	TGGGGATCCG	CCATTGTTGA	AATTGGTATT	AACACTGATA	1320
	CAAAAAGAAA	CACAAGTCGT	GCGTGTGAA	CTATCGCG			1358

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 1197 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

30	GCGATAGTTC	AACACGCACG	ACTTGTGTTT	CTTTTTGTAT	CAGTGTTAAT	ACCAATTTCA	60
	ACAATGGCGG	ATCCCCAGGA	ATTGTCTACA	AGTATTAACC	AGTTTGCTGG	AAGCCTGTAC	120
	AATACGGTTG	CTTCTGGCAA	CAAAGACAAT	CTCATCATGT	CCCCATTGTC	TGTACAAACT	180
	GTTCTATCCC	TGGTGTCAAT	GGGAGCTGGT	GGTAATACTG	CCACACAAAT	AGCTGCTGGT	240
	TTACGTCAGC	CTCAATCAAA	AGAAAAAATT	CAAGATGACT	ACCATGCATT	GATGAACACT	300
35	CTTAATACAC	AAAAAGGTGT	AACTCTGGAA	ATTGCCAACA	AAGTTTACGT	TATGGAAGGC	360
	TATACATTGA	AACCCACCTT	CAAAGAAGTT	GCCACCAACA	AATTCTTAGC	TGGAGCAGAA	420
	AACTTGAAC	TTGCCCCAAA	TGCTGAAAGC	GCTAAAGTTA	TCAACACTTG	GGTTGAAGAA	480
	AAAACTCATG	ACAAAATTCA	TGATTTGATC	AAAGCCGGTG	ATCTAGACCA	GGATTCAAGA	540
	ATGGTTCCTG	TCAATGCATT	GTACTTCAAG	GGTCTTTGGG	AGAAACAATT	CAAGAAGGAA	600
40	AACACTCAAG	ACAAACCTTT	CTATGTTACT	GAAACAGAGA	CAAAGAATGT	ACGAATGATG	660
	CACATTAAGG	ATAAATTCCG	TTATGGAGAA	TTTGAAGAAT	TAGATGCCAA	GGCTGTAGAA	720
	TTGCCCTACA	GGAACTCAGA	TTTGGCCATG	TTAATCATT	TGCCAAACAG	CAAACTGGT	780
	CTCCCCGCTC	TTGAAGAAAA	ATTACAAAAT	GTTGACTTGC	AAAACCTGAC	TCAACGCATG	840
	TACTCTGTTG	AAGTTATTTT	GGATCTGCCT	AAATTCGAAG	TTGAATCTGA	AATTAATTTG	900
45	AATGATCCTC	TGAAAAAGTT	GGGTATGTCT	GATATGTTTG	TTCTTGAAA	AGCTGATTTT	960
	AAAGGATTCG	TTGAAGGATC	TGATGAGATG	TTATATATTT	CTAAAGTAAT	TCAAAAAGCT	1020
	TTCATTGAAG	TAAATGAAGA	AGGTGCTGAA	GCTGCAGCTG	CCACAGGCAT	TGTCATGCTT	1080
	GGTTGCTGTA	TGCCAATGAT	GGATCTTTCT	CCAGTAGTTT	TTAATATTGA	TCACCCATTT	1140
	TATTACTCAT	TGATGACTTG	GGATACTGTT	TTGTTTCAGT	GATGTGTTAA	ATCCCTT	1197

50 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- 55 (A) LENGTH: 1197 nucleic acid
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-106-

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```

AAGGGATTTA ACACATCCAC TGAACAAAAC AGTATCCCAA GTCATCAATG AGTAATAAAA    60
TGGGTGATCA ATATTAAAAA CTACTGGAGA AAGATCCATC ATTGGCATA CAGCAACCAAG    120
5  CATGACAATG CCTGTGGCAG CTGCAGCTTC AGCACCTTCT TCATTTACTT CAATGAAAGC    180
TTTTTGAATT ACTTTAGAAA TATATAACAT CTCATCAGAT CCTTCAAGCA ATCCTTTGAA    240
ATCAGCTTTT CCAGGAACAA ACATATCAGA CATACCCAA CTTTTCAGAG GATCATTCAA    300
ATTAATTTCA GATTCAATCT TGAATTTAGG CAGATCCAAA ATAACCTCAA CAGAGTACAT    360
GCGTTGAGTC AAGTTTTCGA AGTCAACATT TTGTAATTTT TCTTCAAGAG CGGGGAGACC    420
10 AGTTTGTGCTG TTTGGCAAAA TGATTAACAT GGCCAAATCT GAGTTCCTGT AGGGCAATTC    480
TACAGCCTTG GCATCTAATT CTTCAAATTC TCCATAACGG AATTTATCCT TAATGTGCAT    540
CATTCGTACA TTCTTTGTCT CTGTTTCAGT AACATAGAAA GGTTTGTCTT GAGTGTTTTC    600
CTTCTTGAAT TGTTTCTCCC AAAGACCCTT GAAGTACAAT GCATTGACAA GAACCATTCT    660
TGAATCCTGG TCTAGATCAC CGGCTTTGAT CAAATCATGA ATTTTGTCTT GAGTTTTTTC    720
15 TTCAACCCAA GTGTTGATAA CTTTAGCGCT TTCAGCATTT TGGGCAAAGT TCAAGTTTTT    780
TGCTCCAGCT AAGAAATTTGT TGGTGGCAAC TTCTTTGAAG GTGGGTTTCA ATGTATAGCC    840
TTCCATAACG TAAACTTTGT TGGCAATTTT CAGAGTTACA CCTTTTGTG TATTAAGAGT    900
GTTTCATCAAT GCATGGTAGT CATCTTGAAT TTTTCTTTT GATTGAGGCT GACGTAAACC    960
AGCAGCTATT TGTGTGGCAG TATTACCACC AGCTCCCAT TACACCAGGG ATAGAACAGT   1020
20 TTGTACAGAC AATGGGGACA TGATGAGATT GTCTTTGTTG CCAGAAGCAA CCGTATTGTA   1080
CAGGCTTCCA GCAAACTGGT TAATACTTGT AGACAATTCC TGGGGATCCG CCATTGTTGA   1140
AATTGGTATT AACACTGATA CAAAAAGAAA CACAAGTCGT GCGTGTGAA CTATCGC    1197

```

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 376 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

30 Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu
   1           5           10           15
Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser Pro
   20           25           30
Leu Ser Val Gln Thr Val Leu Ser Leu Val Ser Met Gly Ala Gly Gly
35           35           40           45
Asn Thr Ala Thr Gln Ile Ala Ala Gly Leu Arg Gln Pro Gln Ser Lys
   50           55           60
Glu Lys Ile Gln Asp Asp Tyr His Ala Leu Met Asn Thr Leu Asn Thr
   65           70           75           80
40 Gln Lys Gly Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val Met Glu
   85           90           95
Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr Asn Lys Phe
   100          105          110
Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln Asn Ala Glu Ser Ala
45          115          120          125
Lys Val Ile Asn Thr Trp Val Glu Glu Lys Thr His Asp Lys Ile His
   130          135          140

```


-107-

Asp Leu Ile Lys Ala Gly Asp Leu Asp Gln Asp Ser Arg Met Val Leu
 145 150 155 160
 Val Asn Ala Leu Tyr Phe Lys Gly Leu Trp Glu Lys Gln Phe Lys Lys
 165 170 175
 5 Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr Glu Thr Glu Thr Lys
 180 185 190
 Asn Val Arg Met Met His Ile Lys Asp Lys Phe Arg Tyr Gly Glu Phe
 195 200 205
 10 Glu Glu Leu Asp Ala Lys Ala Val Glu Leu Pro Tyr Arg Asn Ser Asp
 210 215 220
 Leu Ala Met Leu Ile Ile Leu Pro Asn Ser Lys Thr Gly Leu Pro Ala
 225 230 235 240
 Leu Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln Arg
 245 250 255
 15 Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile Glu
 260 265 270
 Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp
 275 280 285
 20 Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser
 290 295 300
 Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu
 305 310 315 320
 Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Ala Thr Gly Ile Val Met
 325 330 335
 25 Leu Gly Cys Cys Met Pro Met Met Asp Leu Ser Pro Val Val Phe Asn
 340 345 350
 Ile Asp His Pro Phe Tyr Tyr Ser Leu Met Thr Trp Asp Thr Val Leu
 355 360 365
 30 Phe Ser Gly Cys Val Lys Ser Leu
 370 375

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1838 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA
 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 306..1565
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ATTGTGCAAA GTCAAATTAC GCATTAGAA TATTAAATC AGTATCTCCA AAAATACATA 60

-108-

	CAAATCAATT CAATAACTAT CATTCAAATG ACATCATGTT CAAAATAAAT TAAACACAAA	120
	TATAAAAATG AAGCTAATTT TTGGAAACTG TGTGATTCCA AGGACGACAG AAATATAAAA	180
	CAGATTCATG TGTGTTGTTT CGCGAAGCCA AATGTTTGAA TGTATATAGT GTGTTATTCA	240
	AACATTCTTA GTATTTCTAT ATTATACAAT ATGACTCACA AACGATTCTA ATATCTAGAG	300
5	TTTTG ATG CCG CGT CCT CAG TTT GAC GCG ATA GTT CAA CAC GCA CGA CTT	350
	Met Pro Arg Pro Gln Phe Asp Ala Ile Val Gln His Ala Arg Leu	
	1 5 10 15	
	GTG TTT CTT TTT GTA TCA GTG TTA ATA CCA ATT TCA ACA ATG GCG GAT	398
	Val Phe Leu Phe Val Ser Val Leu Ile Pro Ile Ser Thr Met Ala Asp	
10	20 25 30	
	CCC CAG GAA TTG TCT ACA AGT ATT AAC CAG TTT GCT GGA AGC CTG TAC	446
	Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr	
	35 40 45	
	AAT ACG GTT GCT TCT GGC AAC AAA GAC AAT CTC ATC ATG TCC CCA TTG	494
	Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser Pro Leu	
15	50 55 60	
	TCT GTA CAA ACT GTT CTA TCC CTG GTG TCA ATG GGA GCT GGT GGT AAT	542
	Ser Val Gln Thr Val Leu Ser Leu Val Ser Met Gly Ala Gly Gly Asn	
	65 70 75	
20	ACT GCC ACA CAA ATA GCT GCT GGT TTA CGT CAG CCT CAA TCA AAA GAA	590
	Thr Ala Thr Gln Ile Ala Ala Gly Leu Arg Gln Pro Gln Ser Lys Glu	
	80 85 90 95	
	AAA ATT CAA GAT GAC TAC CAT GCA TTG ATG AAC ACT CTT AAT ACA CAA	638
	Lys Ile Gln Asp Asp Tyr His Ala Leu Met Asn Thr Leu Asn Thr Gln	
25	100 105 110	
	AAA GGT GTA ACT CTG GAA ATT GCC AAC AAA GTT TAC GTT ATG GAA GGC	686
	Lys Gly Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val Met Glu Gly	
	115 120 125	
	TAT ACA TTG AAA CCC ACC TTC AAA GAA GTT GCC ACC AAC AAA TTC TTA	734
	Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr Asn Lys Phe Leu	
30	130 135 140	
	GCT GGA GCA GAA AAC TTG AAC TTT GCC CAA AAT GCT GAA AGC GCT AAA	782
	Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln Asn Ala Glu Ser Ala Lys	
	145 150 155	
35	GTT ATC AAC ACT TGG GTT GAA GAA AAA ACT CAT GAC AAA ATT CAT GAT	830
	Val Ile Asn Thr Trp Val Glu Glu Lys Thr His Asp Lys Ile His Asp	
	160 165 170 175	
	TTG ATC AAA GCC GGT GAT CTA GAC CAG GAT TCA AGA ATG GTT CTT GTC	878
	Leu Ile Lys Ala Gly Asp Leu Asp Gln Asp Ser Arg Met Val Leu Val	
40	180 185 190	
	AAT GCA TTG TAC TTC AAG GGT CTT TGG GAG AAA CAA TTC AAG AAG GAA	926
	Asn Ala Leu Tyr Phe Lys Gly Leu Trp Glu Lys Gln Phe Lys Lys Glu	
	195 200 205	
45	AAC ACT CAA GAC AAA CCT TTC TAT GTT ACT GAA ACA GAG ACA AAG AAT	974
	Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr Glu Thr Glu Thr Lys Asn	
	210 215 220	

-109-

	GTA CGA ATG ATG CAC ATT AAG GAT AAA TTC CGT TAT GGA GAA TTT GAA	1022
	Val Arg Met Met His Ile Lys Asp Lys Phe Arg Tyr Gly Glu Phe Glu	
	225 230 235	
5	GAA TTA GAT GCC AAG GCT GTA GAA TTG CCC TAC AGG AAC TCA GAT TTG	1070
	Glu Leu Asp Ala Lys Ala Val Glu Leu Pro Tyr Arg Asn Ser Asp Leu	
	240 245 250 255	
	GCC ATG TTA ATC ATT TTG CCA AAC AGC AAA ACT GGT CTC CCC GCT CTT	1118
	Ala Met Leu Ile Ile Leu Pro Asn Ser Lys Thr Gly Leu Pro Ala Leu	
	260 265 270	
10	GAA GAA AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT CAA CGC ATG	1166
	Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln Arg Met	
	275 280 285	
15	TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC AAG ATT GAA TCT	1214
	Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile Glu Ser	
	290 295 300	
	GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG TTG GGT ATG TCT GAT ATG	1262
	Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp Met	
	305 310 315	
20	TTT GTT CCT GGA AAA GCT GAT TTC AAA GGA TTG CTT GAA GGA TCT GAT	1310
	Phe Val Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp	
	320 325 330 335	
	GAG ATG TTA TAT ATT TCT AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA	1358
	Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val	
	340 345 350	
25	AAT GAA GAA GGT GCT GAA GCT GCA GCT GCC ACA GCG GTG CTT TTA GTA	1406
	Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Ala Val Leu Leu Val	
	355 360 365	
30	ACG GAA TCT TAT GTA CCT GAG GAA GTA TTC GAA GCT AAT CAT CCC TTT	1454
	Thr Glu Ser Tyr Val Pro Glu Glu Val Phe Glu Ala Asn His Pro Phe	
	370 375 380	
	TAT TTT GCA CTC TAT AAA TCT GCA CAA AAT CCA GTA GAA TCT GAA AAT	1502
	Tyr Phe Ala Leu Tyr Lys Ser Ala Gln Asn Pro Val Glu Ser Glu Asn	
	385 390 395	
35	GAA AGC TCT GAA AAT GAA AAC CCT GAA AAT GTT GAA GTA CTA TTC TCT	1550
	Glu Ser Ser Glu Asn Glu Asn Pro Glu Asn Val Glu Val Leu Phe Ser	
	400 405 410 415	
	GGG AGA TTT ACC AAT TAG AAAAAATATGT GTTACTAGCC TTGTGATTAT	1598
	Gly Arg Phe Thr Asn	
	420	
40	AAGCAGGACA AATTTCAAAA ATACAAGATC TATCTGAATC TCTGGATTAA TGAAGTAATT	1658
	TTTCTACAAT ATTTTTTAAT AGTTATTAGG TCTAAAATAA GTTCATTTTT TAGTATGTGG	1718
	TATAAATCGT GTAGACGAAA AATGTTTTGT TTTAGTTTTT ACTTTTTATG AATGTAATCA	1778
	CCTATATAAT GTTGTAGTTT ATGTAATAAA AATGTTAAAT GTGAAAAAAA AAAAAAAAAA	1838

-110-

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 420 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

	Met	Pro	Arg	Pro	Gln	Phe	Asp	Ala	Ile	Val	Gln	His	Ala	Arg	Leu	Val	
	1				5					10					15		
10	Phe	Leu	Phe	Val	Ser	Val	Leu	Ile	Pro	Ile	Ser	Thr	Met	Ala	Asp	Pro	
				20					25					30			
	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu	Tyr	Asn	
			35					40					45				
15	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro	Leu	Ser	
		50					55					60					
	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	Asn	Thr	
	65					70				75						80	
	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	Glu	Lys	
					85				90						95		
20	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	Gln	Lys	
				100					105					110			
	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	Gly	Tyr	
			115					120					125				
25	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	Leu	Ala	
		130					135					140					
	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	Lys	Val	
	145				150					155					160		
	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	Asp	Leu	
				165					170						175		
30	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	Val	Asn	
				180				185						190			
	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	Glu	Asn	
			195				200					205					
35	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn	Val	
		210					215					220					
	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu	Glu	
	225					230				235					240		
	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	Leu	Ala	
				245					250						255		
40	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	Leu	Glu	
				260					265					270			

-111-

Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln Arg Met Tyr
275 280 285

Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile Glu Ser Glu
290 295 300

5 Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe
305 310 315 320

Val Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu
325 330 335

10 Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn
340 345 350

Glu Glu Gly Ala Glu Ala Ala Ala Thr Ala Val Leu Leu Val Thr
355 360 365

Glu Ser Tyr Val Pro Glu Glu Val Phe Glu Ala Asn His Pro Phe Tyr
370 375 380

15 Phe Ala Leu Tyr Lys Ser Ala Gln Asn Pro Val Glu Ser Glu Asn Glu
385 390 395 400

Ser Ser Glu Asn Glu Asn Pro Glu Asn Val Glu Val Leu Phe Ser Gly
405 410 415

20 Arg Phe Thr Asn
420

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1838 nucleotides
(B) TYPE: nucleic acid
25 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

	TTTTTTTTTT	TTTTTTTCAC	ATTTAACATT	TTTATTACAT	AAACTACAAC	ATTATATAGG	60
30	TGATTACATT	CATAAAAAGT	GAAAACTAAA	ACAAAACATT	TTTCGTCTAC	ACGATTTATA	120
	CCACATACTA	AAAAATGAAC	TTATTTTAGA	CCTAATAACT	ATTAAAAAAT	ATTGTAGAAA	180
	AATTACTTCA	TTAATCCAGA	GATTCAGATA	GATCTTGAT	TTTTGAAATT	TGTCCTGCTT	240
	ATAATCACAA	GGCTAGTAAC	ACATATTTTT	CTAATTGGTA	AATCTCCCAG	AGAATAGTAC	300
	TTCAACATTT	TCAGGGTTTT	CATTTTCAGA	GCTTTCATTT	TCAGATTCTA	CTGGATTTTG	360
35	TGCAGATTTA	TAGAGTGCAA	AATAAAAGGG	ATGATTAGCT	TCGAATACTT	CCTCAGGTAC	420
	ATAAGATTCC	GTTACTAAAA	GCACCGCTGT	GGCAGCTGCA	GCTTCAGCAC	CTTCTTCATT	480
	TACTTCAATG	AAAGCTTTTT	GAATTACTTT	AGAAATATAT	AACATCTCAT	CAGATCCTTC	540
	AAGCAATCCT	TTGAAATCAG	CTTTTCCAGG	AACAAACATA	TCAGACATAC	CCAACTTTTT	600
	CAGAGGATCA	TTCAAATTAA	TTTCAGATTC	AATCTTGAAT	TTAGGCAGAT	CCAAAAATAC	660
40	TTCAACAGAG	TACATGCGTT	GAGTCAAGTT	TTGCAAGTCA	ACATTTTGTA	ATTTTTCCTC	720
	AAGAGCGGGG	AGACCAGTTT	TGCTGTTTGG	CAAAATGATT	AACATGGCCA	AATCTGAGTT	780
	CCTGTAGGGC	AATTCTACAG	CCTTGGCATC	TAATTCTTCA	AATTCTCCAT	AACGGAATTT	840
	ATCCTTAATG	TGCATCATTC	GTACATTCCT	TGTCTCTGTT	TCAGTAACAT	AGAAAGGTTT	900
	GTCTTGAGTG	TTTTCTTCT	TGAATTGTTT	CTCCCAAAGA	CCCTTGAAGT	ACAATGCATT	960
45	GACAAGAACC	ATTCTTGAAT	CCTGGTCTAG	ATCACCGGCT	TTGATCAAAT	CATGAATTTT	1020
	GTCATGAGTT	TTTTCTTCAA	CCCAAGTGTT	GATAACTTTA	GCGCTTTCAG	CATTTTGGGC	1080
	AAAGTTCAAG	TTTTCTGCTC	CAGCTAAGAA	TTTGTGGTG	GCAACTTCTT	TGAAGGTGGG	1140
	TTTCAATGTA	TAGCCTTCCA	TAACGTAAAC	TTTGTGGCA	ATTTCCAGAG	TTACACCTTT	1200

-112-

	TTGTGTATTA	AGAGTGTTCA	TCAATGCATG	GTAAGTCATCT	TGAATTTTTT	CTTTTGATTG	1260
	AGGCTGACGT	AAACCAGCAG	CTATTTGTGT	GGCAGTATTA	CCACCAGCTC	CCATTGACAC	1320
	CAGGGATAGA	ACAGTTTGTA	CAGACAATGG	GGACATGATG	AGATTGTCTT	TGTTGCCAGA	1380
	AGCAACCGTA	TTGTACAGGC	TTCCAGCAAA	CTGGTTAATA	CTTGTAAGCA	ATTCTTGGGG	1440
5	ATCCGCCATT	GTTGAAATTG	GTATTAACAC	TGATACAAAA	AGAAACACAA	GTCGTGCGTG	1500
	TTGAACTATC	GCGTCAAAC	GAGGACGCGG	CATCAAAACT	CTAGATATTA	GAATCGTTTG	1560
	TGAGTCATAT	TGTATAATAT	AGAAATACTA	GGAATGTTTG	AATAACACAC	TATATACATT	1620
	CAAACATTTG	GCTTCGCGGA	ACAACACACA	TGAATCTGTT	TTATATTTCT	GTCGTCCTTG	1680
	GAATCACACA	GTTTCCAAAA	ATTAGCTTCA	TTTTTATATT	TGTGTTTAAT	TTATTTTGAA	1740
10	CATGATGTCA	TTTGAATGAT	AGTTATTGAA	TTGATTTGTA	TGTATTTTTG	GAGATACTGA	1800
	TTTTAATATT	CTAAATGCGT	AATTTGACTT	TGCACAAAT			1838

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1260 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cdNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

20	ATGCCGCGTC	CTCAGTTTGA	CGCGATAGTT	CAACACGCAC	GACTTGTTGTT	TCTTTTTTGTA	60
	TCAGTGTAA	TACCAATTTC	AACAATGGCG	GATCCCCAGG	AATTGTCTAC	AAGTATTAAAC	120
	CAGTTTGCTG	GAAGCCTGTA	CAATACGGTT	GCTTCTGGCA	ACAAAGACAA	TCTCATCATG	180
	TCCCCATTGT	CTGTACAAAC	TGTTCTATCC	CTGGTGTCAC	TGGGAGCTGG	TGGTAATACT	240
	GCCACACAAA	TAGCTGCTGG	TTTACGTCAG	CCTCAATCAA	AAGAAAAAAT	TCAAGATGAC	300
25	TACCATGCAT	TGATGAACAC	TCTTAATACA	CAAAAAGGTG	TAAGTCTGGA	AATTGCCAAC	360
	AAAGTTTACG	TTATGGAAGG	CTATACATTG	AAACCCACCT	TCAAAGAAGT	TGCCACCAAC	420
	AAATTCTTAG	CTGGAGCAGA	AAACTTGAAC	TTTGCCCAAA	ATGCTGAAAG	CGCTAAAGTT	480
	ATCAACACTT	GGGTGAAGA	AAAACTCAT	GACAAAAATC	ATGATTTGAT	CAAAGCCGGT	540
	GATCTAGACC	AAGATTCAAG	AATGGTTCTT	GTCAATGCAT	TGTAATTCAC	GGGTCTTTGG	600
30	GAGAAACAAT	TCAAGAAGGA	AAACACTCAA	GACAAACCTT	TCTATGTTAC	TGAAACAGAG	660
	ACAAAGAATG	TACGAATGAT	GCACATTAAG	GATAAATTC	GTTATGGAGA	ATTTGAAGAA	720
	TTAGATGCCA	AGGCTGTAGA	ATTGCCCTAC	AGGAACTCAG	ATTTGGCCAT	GTTAATCAAT	780
	TTGCCAAACA	GCAAACTGG	TCTCCCCGCT	CTTGAAGAAA	AATTACAAAA	TGTTGACTTG	840
	CAAACTTGA	CTCAACGCAT	GTAATCTGTT	GAAGTTATTT	TGGATCTGCC	TAAATTCAAG	900
35	ATTGAATCTG	AAATTAATTT	GAATGATCCT	CTGAAAAAGT	TGGGTATGTC	TGATATGTTT	960
	GTTCTTGGA	AAGCTGATTT	CAAAGGATTT	CTTGAAGGAT	CTGATGAGAT	GTTATATATT	1020
	TCTAAAGTAA	TTCAAAAAGC	TTTCATTGAA	GTAATGAAG	AAGGTGCTGA	AGCTGCAGCT	1080
	GCCACAGCGG	TGCTTTTAGT	AACGGAATCT	TATGTACCTG	AGGAAGTATT	CGAAGCTAAT	1140
	CATCCCTTTT	ATTTTGCACT	CTATAAATCT	GCACAAAATC	CAGTAGAATC	TGAAAATGAA	1200
40	AGCTCTGAAA	ATGAAAACCC	TGAAAATGTT	GAAGTACTAT	TCTCTGGGAG	ATTTACCAAT	1260

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1260 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cdNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

50	ATTGGTAAAT	CTCCCAGAGA	ATAGTACTTC	AACATTTTCA	GGGTTTTTCAT	TTTCAGAGCT	60
	TTCATTTTCA	GATTCTACTG	GATTTTGTGC	AGATTTATAG	AGTGCAAAAT	AAAAGGGATG	120
	ATTAGCTTCG	AATACTTCCT	CAGGTACATA	AGATTCGGTT	ACTAAAAGCA	CCGCTGTGGC	180
	AGCTGCAGCT	TCAGCACCTT	CTTCATTTAC	TTCAATGAAA	GCTTTTTTGA	TTACTTTAGA	240

-113-

	AATATATAAC	ATCTCATCAG	ATCCTTCAAG	CAATCCTTTG	AAATCAGCTT	TTCCAGGAAC	300
	AAACATATCA	GACATACCCA	ACTTTTTCAG	AGGATCATTC	AAATTAATTT	CAGATTCAAT	360
	CTTGAATTTA	GGCAGATCCA	AAATAACTTC	AACAGAGTAC	ATGCGTTGAG	TCAAGTTTTG	420
	CAAGTCAACA	TTTTGTAATT	TTTCTTCAAG	AGCGGGGAGA	CCAGTTTTCG	TGTTTGGCAA	480
5	AATGATTAAC	ATGGCCAAAT	CTGAGTTCCT	GTAGGGCAAT	TCTACAGCCT	TGGCATCTAA	540
	TTCTTCAAAT	TCTCCATAAC	GGAATTTATC	CTTAATGTGC	ATCATTCGTA	CATTCTTTGT	600
	CTCTGTTTCA	GTAACATAGA	AAGGTTTGTC	TTGAGTGTTC	TCCTTCTTGA	ATTGTTTCTC	660
	CCAAAGACCC	TTGAAGTACA	ATGCATTGAC	AAGAACCATT	CTTGAATCCT	GGTCTAGATC	720
	ACCGGCTTTG	ATCAAATCAT	GAATTTTGTC	ATGAGTTTTT	TCTTCAACCC	AAGTGTGTGAT	780
10	AACTTTAGCG	CTTTCAGCAT	TTTGGGCAAA	GTTCAAGTTT	TCTGCTCCAG	CTAAGAATTT	840
	GTTGGTGGCA	ACTTCTTTGA	AGGTGGGTTT	CAATGTATAG	CCTTCCATAA	CGTAAACTTT	900
	GTTGGCAATT	TCCAGAGTTA	CACCTTTTTC	TGTATTAAGA	GTGTTTCATCA	GTGTTGTTGA	960
	GTCTCTTTGA	ATTTTTTCTT	TTGATTGAGG	CTGACGTAAA	CCAGCAGCTA	TTTGTGTGGC	1020
	AGTATTACCA	CCAGCTCCCA	TTGACACCAG	GGATAGAACA	GTTTGTACAG	ACAATGGGGA	1080
15	CATGATGAGA	TTGTCTTTGT	TGCCAGAAGC	AACCGTATTG	TACAGGCTTC	CAGCAAACCTG	1140
	GTTAATACTT	GTAGACAATT	CCTGGGGATC	CGCCATTGTT	GAAATTGGTA	TTAACACTGA	1200
	TACAAAAAGA	AACACAAGTC	GTGCGTGTTC	AACTATCGCG	TCAAACCTGAG	GACGCGGCAT	1260

(2) INFORMATION FOR SEQ ID NO:18:

	(i)	SEQUENCE CHARACTERISTICS:															
20		(A)	LENGTH: 390 amino acids														
		(B)	TYPE: amino acid														
		(D)	TOPOLOGY: linear														
	(ii)	MOLECULE TYPE: protein															
	(xi)	SEQUENCE DESCRIPTION: SEQ ID NO:18:															
25	Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu	
	1				5					10					15		
	Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro	
				20					25					30			
	Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	
30			35					40					45				
	Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	
		50					55					60					
	Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	
	65					70					75					80	
35	Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	
					85					90					95		
	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	
				100					105					110			
	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	
40			115					120					125				
	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	
		130					135					140					
	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	
	145					150				155						160	
45	Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	
				165						170					175		

-114-

Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr Glu Thr Glu Thr Lys
 180 185 190
 Asn Val Arg Met Met His Ile Lys Asp Lys Phe Arg Tyr Gly Glu Phe
 195 200 205
 5 Glu Glu Leu Asp Ala Lys Ala Val Glu Leu Pro Tyr Arg Asn Ser Asp
 210 215 220
 Leu Ala Met Leu Ile Ile Leu Pro Asn Ser Lys Thr Gly Leu Pro Ala
 225 230 235 240
 10 Leu Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln Arg
 245 250 255
 Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile Glu
 260 265 270
 Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp
 275 280 285
 15 Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser
 290 295 300
 Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu
 305 310 315 320
 20 Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Ala Val Leu Leu
 325 330 335
 Val Thr Glu Ser Tyr Val Pro Glu Glu Val Phe Glu Ala Asn His Pro
 340 345 350
 Phe Tyr Phe Ala Leu Tyr Lys Ser Ala Gln Asn Pro Val Glu Ser Glu
 355 360 365
 25 Asn Glu Ser Ser Glu Asn Glu Asn Pro Glu Asn Val Glu Val Leu Phe
 370 375 380
 Ser Gly Arg Phe Thr Asn
 385 390

(2) INFORMATION FOR SEQ ID NO:19:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1414 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 2..1180

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

40 A CGA CTT GTG TTT CTT TTT GTA TCA GTG TTA ATA CCA ATT TCA ACA
 Arg Leu Val Phe Leu Phe Val Ser Val Leu Ile Pro Ile Ser Thr
 1 5 10 15

46

-115-

	ATG GCG GAT CCC CAG GAA TTG TCT ACA AGT ATT AAC CAG TTT GCT GGA	94
	Met Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly	
	20 25 30	
5	AGC CTG TAC AAT ACG GTT GCT TCT GGC AAC AAA GAC AAT CTC ATC ATG	142
	Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met	
	35 40 45	
	TCC CCA TTG TCT GTA CAA ACT GTT CTA TCC CTG GTG TCA ATG GGA GCT	190
	Ser Pro Leu Ser Val Gln Thr Val Leu Ser Leu Val Ser Met Gly Ala	
	50 55 60	
10	GGT GGT AAT ACT GCC ACA CAA ATA GCT GCT GGT TTA CGT CAG CCT CAA	238
	Gly Gly Asn Thr Ala Thr Gln Ile Ala Ala Gly Leu Arg Gln Pro Gln	
	65 70 75	
15	TCA AAA GAA AAA ATT CAA GAT GAC TAC CAT GCA TTG ATG AAC ACT CTT	286
	Ser Lys Glu Lys Ile Gln Asp Asp Tyr His Ala Leu Met Asn Thr Leu	
	80 85 90 95	
	AAT ACA CAA AAA GGT GTA ACT CTG GAA ATT GCC AAC AAA GTT TAC GTT	334
	Asn Thr Gln Lys Gly Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val	
	100 105 110	
20	ATG GAA GGC TAT ACA TTG AAA CCC ACC TTC AAA GAA GTT GCC ACC AAC	382
	Met Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr Asn	
	115 120 125	
	AAA TTC TTA GCT GGA GCA GAA AAC TTG AAC TTT GCC CAA AAT GCT GAA	430
	Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln Asn Ala Glu	
	130 135 140	
25	AGC GCT AAA GTT ATC AAC ACT TGG GTT GAA GAA AAA ACT CAT GAC AAA	478
	Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu Lys Thr His Asp Lys	
	145 150 155	
30	ATT CAT GAT TTG ATC AAA GCC GGT GAT CTA GAC CAG GAT TCA AGA ATG	526
	Ile His Asp Leu Ile Lys Ala Gly Asp Leu Asp Gln Asp Ser Arg Met	
	160 165 170 175	
	GTT CTT GTC AAT GCA TTG TAC TTC AAG GGT CTT TGG GAG AAA CAA TTC	574
	Val Leu Val Asn Ala Leu Tyr Phe Lys Gly Leu Trp Glu Lys Gln Phe	
	180 185 190	
35	AAG AAG GAA AAC ACT CAA GAC AAA CCT TTC TAT GTT ACT GAA ACA GAG	622
	Lys Lys Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr Glu Thr Glu	
	195 200 205	
	ACA AAG AAT GTA CGA ATG ATG CAC ATT AAG GAT AAA TTC CGT TAT GGA	670
	Thr Lys Asn Val Arg Met Met His Ile Lys Asp Lys Phe Arg Tyr Gly	
	210 215 220	
40	GAA TTT GAA GAA TTA GAT GCC AAG GCT GTA GAA TTG CCC TAC AGG AAC	718
	Glu Phe Glu Glu Leu Asp Ala Lys Ala Val Glu Leu Pro Tyr Arg Asn	
	225 230 235	
45	TCA GAT TTG GCC ATG TTA ATC ATT TTG CCA AAC AGC AAA ACT GGT CTC	766
	Ser Asp Leu Ala Met Leu Ile Ile Leu Pro Asn Ser Lys Thr Gly Leu	
	240 245 250 255	

-116-

	CCC GCT CTT GAA GAA AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT	814
	Pro Ala Leu Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr	
	260 265 270	
5	CAA CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC AAG	862
	Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys	
	275 280 285	
	ATT GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG TTG GGT ATG	910
	Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met	
	290 295 300	
10	TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC AAA GGA TTG CTT GAA	958
	Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu	
	305 310 315	
	GGA TCT GAT GAG ATG TTA TAT ATT TCT AAA GTA ATT CAA AAA GCT TTC	1006
15	Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe	
	320 325 330 335	
	ATT GAA GTA AAT GAA GAA GGT GCT GAA GCT GCA GCT GCC ACA GGC GTG	1054
	Ile Glu Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Gly Val	
	340 345 350	
20	ATG TTA ATG ATG CGT TGT ATG CCA ATG ATG CCA ATG GCC TTC AAT GCT	1102
	Met Leu Met Met Arg Cys Met Pro Met Met Pro Met Ala Phe Asn Ala	
	355 360 365	
	GAG CAT CCA TTC CTG TAC TTC TTA CAC AGC AAA AAT TCT GTT CTA TTC	1150
	Glu His Pro Phe Leu Tyr Phe Leu His Ser Lys Asn Ser Val Leu Phe	
	370 375 380	
25	AAT GGT CGT CTT GTT AAA CCA ACA ACT GAA TAA AAGCCAAATG CACTTCACTA	1203
	Asn Gly Arg Leu Val Lys Pro Thr Thr Glu	
	385 390	
	ATATTTTTTTA ATTGCTTACT GAAACAGTGC CTGTAGAACA TTGTGTTCAA TTTATATTTG	1263
	TCAGCTTTTAA GTATTCAGTA TTTTATATCA TCACTATTTT AGTGGTGGAT CTTAAGTACA	1323
30	AATTTATTGT TATGATATAT ATTTATTTTT TGTGAATATT TTTTAAACAA ATTTTGATAA	1383
	AAAACATAAG ACTAAAAAAA AAAAAAAAAA A	1414

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 393 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

40	Arg Leu Val Phe Leu Phe Val Ser Val Leu Ile Pro Ile Ser Thr Met
	1 5 10 15
	Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser
	20 25 30
	Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser
	35 40 45

-117-

	Pro	Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	
	50						55					60					
	Gly	Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	
	65					70					75					80	
5	Lys	Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	
					85					90					95		
	Thr	Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	
				100					105					110			
10	Glu	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	
			115					120					125				
	Phe	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	
	130						135					140					
	Ala	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	
	145					150					155					160	
15	His	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	
					165					170					175		
	Leu	Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	
				180					185					190			
20	Lys	Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	
			195					200					205				
	Lys	Asn	Val	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	
	210						215					220					
	Phe	Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	
	225					230					235					240	
25	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	
					245					250					255		
	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	
					260				265					270			
30	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp	Leu	Pro	Lys	Phe	Lys	Ile	
			275					280					285				
	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	
	290						295					300					
	Asp	Met	Phe	Val	Pro	Gly	Lys	Ala	Asp	Phe	Lys	Gly	Leu	Leu	Glu	Gly	
	305					310					315					320	
35	Ser	Asp	Glu	Met	Leu	Tyr	Ile	Ser	Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile	
					325					330					335		
	Glu	Val	Asn	Glu	Glu	Gly	Ala	Glu	Ala	Ala	Ala	Ala	Thr	Gly	Val	Met	
					340				345					350			
40	Leu	Met	Met	Arg	Cys	Met	Pro	Met	Met	Pro	Met	Ala	Phe	Asn	Ala	Glu	
					355				360				365				
	His	Pro	Phe	Leu	Tyr	Phe	Leu	His	Ser	Lys	Asn	Ser	Val	Leu	Phe	Asn	
							375					380					

-118-

Gly Arg Leu Val Lys Pro Thr Thr Glu
385 390

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1414 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```

TTTTTTTTTT TTTTTTTTAG TCTTATGTTT TTTATCAAAA TTTGTAAAAA AAATATTCAC 60
AAAAAATAAA TATATATCAT AACAATAAAT TTGTACTTAA GATCCACCAC TGAAATAGTG 120
ATGATAAAAA ATACTGAATA CTAAAGCTG ACAAATATAA ATTGAACACA ATGTTCTACA 180
GGCACTGTTT CAGTAAGCAA TTAAAAAATA TTAGTGAAGT GCATTTGGCT TTTATTTCAGT 240
15 TGTGGTTTA ACAAGACGAC CATTGAATAG AACAGAATTT TTGCTGTGTA AGAAGTACAG 300
GAATGGATGC TCAGCATTGA AGGCCATTGG CATCATTGGC ATACAACGCA TCATTAACAT 360
CACGCTGTG GCAGCTGCAG CTTCAGCACC TTCTTCATTT ACTTCAATGA AAGCTTTTTG 420
AATTACTTTA GAAATATATA ACATCTCATC AGATCCTTCA AGCAATCCTT TGAAATCAGC 480
TTTTCAGGA ACAAACATAT CAGACATACC CAACTTTTTT AGAGGATCAT TCAAATTAAT 540
20 TTCAGATTCA ATCTTGAATT TAGGCAGATC CAAAATAACT TCAACAGAGT ACATGCGTTG 600
AGTCAAGTTT TGCAAGTCAA CATTTTGTA TTTTCTTCA AGAGCGGGGA GACCAGTTTT 660
GCTGTTGGC AAAATGATTA ACATGGCCAA ATCTGAGTTC CTGTAGGGCA ATTCTACAGC 720
CTTGGCATCT AATTCTTCAA ATTCTCCATA ACGGAATTTA TCCTTAATGT GCATCATTCG 780
TACATCTTTT GTCTCTGTTT CAGTAACATA GAAAGGTTTG TCTTGAGTGT TTTCTTCTT 840
25 GAATGTGTTT TCCCAAAGAC CCTTGAAGTA CAATGCATTG ACAAGAACCA TTCTTGAATC 900
CTGGTCTAGA TCACCGGCTT TGATCAAATC ATGAATTTTG TCATGAGTTT TTTCTTCAAC 960
CCAAGTGTTG ATAACTTTAG CGCTTTCAGC ATTTTGGGCA AAGTTCAAGT TTTCTGCTCC 1020
AGCTAAGAAAT TTGTTGGTGG CAACTTCTTT GAAGGTGGGT TTCAATGTAT AGCCTTCCAT 1080
AACGTAAACT TTGTTGGCAA TTTCCAGAT TACACCTTTT TGTGTATTAA GAGTGTTTCA 1140
30 CAATGCATGG TAGTCATCTT GAATTTTTC TTTTGATTGA GGCTGACGTA AACCAGCAGC 1200
TATTGTGTG GCAGTATTAC CACCAGCTCC CATTGACACC AGGGATAGAA CAGTTGTAC 1260
AGACAATGGG GACATGATGA GATTGTCTTT GTTGCCAGAA GCAACCGTAT TGTACAGGCT 1320
TCCAGCAAAC TGGTTAATAC TTGTAGACAA TTCTGGGGA TCCGCCATTG TTGAAATTGG 1380
TATTAACACT GATACAAAAA GAAACACAAG TCGT 1414

```

35 (2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 1179 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

CGACTTGTGT TTCTTTTTGT ATCAGTGTGA ATACCAATTT CAACAATGGC GGATCCCCAG 60
GAATGTGCTA CAAGTATTAA CCAGTTTGCT GGAAGCCTGT ACAATACGGT TGCTTCTGGC 120
45 AACAAAGACA ATCTCATCAT GTCCCCATTG TCTGTACAAA CTGTTCTATC CTTGGTGTCA 180
ATGGGAGCTG GTGGTAATAC TGCCACACAA ATAGCTGCTG GTTTACGTCA GCCTCAATCA 240
AAAGAAAAAA TTCAAGATGA CTACCATGCA TTGATGAACA CTCTTAATAC AAAAAAGGT 300
GTAACCTCTG AAATTGCCAA CAAAGTTTAC GTTATGGAAG GCTATACATT GAAACCCACC 360
TTCAAAGAAG TTGCCACCAA CAAATTCCTT GCTGGAGCAG AAAACTTGAA CTTTGCCCAA 420
50 AATGCTGAAA GCGCTAAAGT TATCAACACT TGGGTGGAAG AAAAACTCA TGACAAAATT 480
CATGATTGTA TCAAAGCCGG TGATCTAGAC CAGGATTCAA GAATGGTTCT TGTCAATGCA 540
TTGTACTTCA AGGGTCTTTG GGAGAAACAA TTCAAGAAGG AAAACACTCA AGACAAACCT 600

```

-119-

	TTCTATGTTA	CTGAAACAGA	GACAAAGAAT	GTACGAATGA	TGCACATTAA	GGATAAATTC	660
	CGTTATGGAG	AATTTGAAGA	ATTAGATGCC	AAGGCTGTAG	AATTGCCCTA	CAGGAAGTCA	720
	GATTTGGCCA	TGTTAATCAT	TTTGCCAAAC	AGCAAACTG	GTCTCCCCGC	TCTTGAAGAA	780
	AAATTACAAA	ATGTTGACTT	GCAAACTTG	ACTCAACGCA	TGTACTCTGT	TGAAGTTATT	840
5	TTGGATCTGC	CTAAATTCAA	GATTGAATCT	GAAATTAATT	TGAATGATCC	TCTGAAAAAG	900
	TTGGGTATGT	CTGATATGTT	TGTTCTTGGG	AAAGCTGATT	TCAAAGGATT	GCTTGAAGGA	960
	TCTGATGAGA	TGTTATATAT	TTCTAAAGTA	ATTCAAAAAG	CTTTCATTGA	AGTAAATGAA	1020
	GAAGGTGCTG	AAGCTGCAGC	TGCCACAGGC	GTGATGTTAA	TGATGCGTTG	TATGCCAATG	1080
	ATGCCAATGG	CCTTCAATGC	TGAGCATCCA	TTCCTGTACT	TCTTACACAG	CAAAAATTCT	1140
10	GTTCTATTCA	ATGGTCGTCT	TGTTAAACCA	ACAAGTAA			1179

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

	(A) LENGTH:	1179 nucleotides
	(B) TYPE:	nucleic acid
15	(C) STRANDEDNESS:	single
	(D) TOPOLOGY:	linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

	TTCAGTTGTT	GGTTTAACAA	GACGACCATT	GAATAGAACA	GAATTTTTGC	TGTGTAAGAA	60
20	GTACAGGAAT	GGATGCTCAG	CATTGAAGGC	CATTGGCATC	ATTGGCATAC	AACGCATCAT	120
	TAACATCACG	CCTGTGGCAG	CTGCAGCTTC	AGCACCTTCT	TCATTTACTT	CAATGAAAGC	180
	TTTTTTGAATT	ACTTTAGAAA	TATATAACAT	CTCATCAGAT	CCTTCAAGCA	ATCCTTTGAA	240
	ATCAGCTTTT	CCAGGAACAA	ACATATCAGA	CATACCCAAC	TTTTTCAGAG	GATCATTCAA	300
	ATTAATTTCA	GATTCAATCT	TGAATTTAGG	CAGATCCAAA	ATAACTTCAA	CAGAGTACAT	360
25	GCGTTGAGTC	AAGTTTTGCA	AGTCAACATT	TTGTAATTTT	TCTTCAAGAG	CGGGGAGACC	420
	AGTTTTGCTG	TTTGGCAAAA	TGATTAACAT	GGCCAAATCT	GAGTTCCTGT	AGGGCAATTC	480
	TACAGCCTTG	GCATCTAATT	CTTCAAATTC	TCCATAACGG	AATTTATCCT	TAATGTGCAT	540
	CATTTCGTACA	TTCTTTGTCT	CTGTTTCAGT	AACATAGAAA	GGTTTGTCTT	GAGTGTTTTC	600
	CTTCTTGAAT	TGTTTCTCCC	AAAGACCCTT	GAAGTACAAT	GCATTGACAA	GAACCATTCT	660
30	TGAATCCTGG	TCTAGATCAC	CGGCTTTGAT	CAAATCATGA	ATTTTGTGAT	GAGTTTTTTC	720
	TTCAACCCAA	GTGTTGATAA	CTTTAGCGCT	TTCAGCATTT	TGGGCAAGT	TCAAGTTTTC	780
	TGCTCCAGCT	AAGAATTTGT	TGGTGGCAAC	TTCTTTGAAG	GTGGGTTTCA	ATGTATAGCC	840
	TTCCATAACG	TAAACTTTGT	TGGCAATTTT	CAGAGTTACA	CCTTTTGTG	TATTAAGAGT	900
	GTTTCATCAAT	GCATGGTAGT	CATCTTGAAT	TTTTTCTTTT	GATTGAGGCT	GACGTAAACC	960
35	AGCAGCTATT	TGTGTGGCAG	TATTACCACC	AGCTCCCAT	GACACCAGGG	ATAGAACAGT	1020
	TTGTACAGAC	AATGGGGACA	TGATGAGATT	GTCTTTGTTG	CCAGAAGCAA	CCGTATTGTA	1080
	CAGGCTTCCA	GCAAACTGGT	TAATACTTGT	AGACAAATTC	TGGGGATCCG	CCATTGTTGA	1140
	AATTGGTATT	AACACTGATA	CAAAAAGAAA	CACAAGTCG			1179

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

	(A) LENGTH:	376 amino acids
	(B) TYPE:	amino acid
40	(D) TOPOLOGY:	linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu
1				5					10					15	

Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro
			20				25						30		

-120-

	Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	
			35					40					45				
	Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	
		50					55					60					
5	Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	
	65					70					75					80	
	Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	
					85					90					95		
10	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	
				100					105					110			
	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	
			115					120					125				
	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	
		130					135					140					
15	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	
	145					150					155					160	
	Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	
				165					170						175		
20	Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	
				180					185					190			
	Asn	Val	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	
			195					200					205				
	Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	
		210					215					220					
25	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	
	225					230					235					240	
	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	
				245						250					255		
30	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp	Leu	Pro	Lys	Phe	Lys	Ile	Glu	
				260					265						270		
	Ser	Glu	Ile	Asn	Leu	Asn	Asp	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	
			275					280					285				
	Met	Phe	Val	Pro	Gly	Lys	Ala	Asp	Phe	Lys	Gly	Leu	Leu	Glu	Gly	Ser	
		290					295					300					
35	Asp	Glu	Met	Leu	Tyr	Ile	Ser	Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile	Glu	
	305					310					315					320	
	Val	Asn	Glu	Glu	Gly	Ala	Glu	Ala	Ala	Ala	Ala	Thr	Gly	Val	Met	Leu	
				325						330					335		
40	Met	Met	Arg	Cys	Met	Pro	Met	Met	Pro	Met	Ala	Phe	Asn	Ala	Glu	His	
				340					345					350			
	Pro	Phe	Leu	Tyr	Phe	Leu	His	Ser	Lys	Asn	Ser	Val	Leu	Phe	Asn	Gly	
			355					360					365				

-121-

Arg Leu Val Lys Pro Thr Thr Glu
370 375

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:
5 (A) LENGTH: 1492 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 3..1196

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

15	CG ATA GTT CAA CAC GCA CGA CTT GTG TTT CTT TTT GTA TCA GTG TTA	47
	Ile Val Gln His Ala Arg Leu Val Phe Leu Phe Val Ser Val Leu	
	1 5 10 15	
	ATA CCA ATT TCA ACA ATG GCG GAT CCC CAG GAA TTG TCT ACA AGT ATT	95
	Ile Pro Ile Ser Thr Met Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile	
	20 25 30	
20	AAC CAG TTT GCT GGA AGC CTG TAC AAT ACG GTT GCT TCT GGC AAC AAA	143
	Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys	
	35 40 45	
	GAC AAT CTC ATC ATG TCC CCA TTG TCT GTA CAA ACT GTT CTA TCC CTG	191
25	Asp Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu Ser Leu	
	50 55 60	
	GTG TCA ATG GGA GCT GGT GGT AAT ACT GCC ACA CAA ATA GCT GCT GGT	239
	Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile Ala Ala Gly	
	65 70 75	
	TTA CGT CAG CCT CAA TCA AAA GAA AAA ATT CAA GAT GAC TAC CAC GCA	287
30	Leu Arg Gln Pro Gln Ser Lys Glu Lys Ile Gln Asp Asp Tyr His Ala	
	80 85 90 95	
	TTG ATG AAC ACT CTT AAT ACA CAA AAA GGT GTA ACT CTG GAA ATT GCC	335
	Leu Met Asn Thr Leu Asn Thr Gln Lys Gly Val Thr Leu Glu Ile Ala	
	100 105 110	
35	AAT AAA GTT TAT GTT ATG GAA GGC TAT ACA TTA AAA CCC ACC TTC AAA	383
	Asn Lys Val Tyr Val Met Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys	
	115 120 125	
	GAA GTT GCC ACC AAC AAA TTC TTA GCT GGA GCA GAA AAC TTG AAC TTT	431
40	Glu Val Ala Thr Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe	
	130 135 140	
	GCC CAA AAT GCT GAA AGC GCT AAA GTT ATC AAC ACT TGG GTT GAA GAA	479
	Ala Gln Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu	
	145 150 155	
45	AAA ACT CAT GAC AAA ATT CAT GAT TTG ATC AAA GCC GGT GAT CTA GAC	527
	Lys Thr His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp Leu Asp	
	160 165 170 175	

-122-

	CAG GAT TCA AGA ATG GTT CTT GTC AAT GCA TTG TAC TTC AAG GGT CTT	575
	Gln Asp Ser Arg Met Val Leu Val Asn Ala Leu Tyr Phe Lys Gly Leu	
	180 185 190	
5	TGG GAG AAA CAA TTC AAG AAG GAA AAC ACC CAA GAC AAA CCT TTC TAT	623
	Trp Glu Lys Gln Phe Lys Lys Glu Asn Thr Gln Asp Lys Pro Phe Tyr	
	195 200 205	
	GTT ACT GAA ACA GAG ACA AAG AAT GTA CGA ATG ATG CAC ATT AAG GAT	671
	Val Thr Glu Thr Glu Thr Lys Asn Val Arg Met Met His Ile Lys Asp	
	210 215 220	
10	AAA TTC CGT TAT GGA GAA TTT GAA GAA TTA GAT GCC AAG GCT GTA GAA	719
	Lys Phe Arg Tyr Gly Glu Phe Glu Glu Leu Asp Ala Lys Ala Val Glu	
	225 230 235	
15	TTG CCC TAC AGG AAC TCA GAT TTG GCC ATG TTA ATC ATT TTG CCA AAC	767
	Leu Pro Tyr Arg Asn Ser Asp Leu Ala Met Leu Ile Ile Leu Pro Asn	
	240 245 250 255	
	AGC AAA ACT GGT CTC CCC ACT CTT GAA GAA AAA TTA CAA AAT GTT GAT	815
	Ser Lys Thr Gly Leu Pro Thr Leu Glu Glu Lys Leu Gln Asn Val Asp	
	260 265 270	
20	TTG CAA AAC TTG ACT CAA CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT	863
	Leu Gln Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp	
	275 280 285	
	CTG CCT AAA TTC AAA ATT GAG TCT GAA ATT AAT TTG AAT GAT CCT CTG	911
	Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu	
	290 295 300	
25	AAA AAG TTG GGT ATG TCT GAT ATG TTC ATG CCT GGA AAA GCT GAT TTC	959
	Lys Lys Leu Gly Met Ser Asp Met Phe Met Pro Gly Lys Ala Asp Phe	
	305 310 315	
30	AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT AAA GTA	1007
	Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val	
	320 325 330 335	
	ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT GCT GAA GCT GCA	1055
	Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu Ala Ala	
	340 345 350	
35	GCT GCC ACA GGC GTG ATG TTA ATG ATG CGT TGT ATG CCA ATG ATG CCA	1103
	Ala Ala Thr Gly Val Met Leu Met Met Arg Cys Met Pro Met Met Pro	
	355 360 365	
	ATG GCC TTC AAT GCT GAG CAT CCA TTC CTG TAC TTC TTA CAC AGC AAA	1151
	Met Ala Phe Asn Ala Glu His Pro Phe Leu Tyr Phe Leu His Ser Lys	
	370 375 380	
40	AAT TCT GTT CTA TTC AAT GGT CGT CTT GTT AAA CCA ACA ACT GAA TAA	1199
	Asn Ser Val Leu Phe Asn Gly Arg Leu Val Lys Pro Thr Thr Glu	
	385 390 395	
45	AAGCCAAATG CACTTCACTA ATATTTTTTA ATTGCTTACT GAAACAGTGC CTGTAGAACA	1259
	TTGTGTTCAA TTTATATTTG TCAGCTTTAA GTATTCAGTA TTTTATATCA TCACTATTTT	1319
	AGTGGTGGAT CTTAAGTACA AATTTATTGT TATGATATAT ATTTATTTTT TGTGAATATT	1379
	TTTTTAACAA ATTTTGATAA AAAACATAAG ACTAAAAATA AAAGAAAAAT TAAAATTTAT	1439
	GTATAATTGT TGTATACTAA ATTATATCTT TAAGAAAAAA AAAAAAAAAA AAA	1492

-123-

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 398 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

```

Ile Val Gln His Ala Arg Leu Val Phe Leu Phe Val Ser Val Leu Ile
 1             5             10             15
10 Pro Ile Ser Thr Met Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn
    20             25             30
    Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp
        35             40             45
15 Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu Ser Leu Val
    50             55             60
    Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile Ala Ala Gly Leu
    65             70             75             80
    Arg Gln Pro Gln Ser Lys Glu Lys Ile Gln Asp Asp Tyr His Ala Leu
        85             90             95
20 Met Asn Thr Leu Asn Thr Gln Lys Gly Val Thr Leu Glu Ile Ala Asn
    100             105             110
    Lys Val Tyr Val Met Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu
    115             120             125
25 Val Ala Thr Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala
    130             135             140
    Gln Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu Lys
    145             150             155             160
    Thr His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp Leu Asp Gln
    165             170             175
30 Asp Ser Arg Met Val Leu Val Asn Ala Leu Tyr Phe Lys Gly Leu Trp
    180             185             190
    Glu Lys Gln Phe Lys Lys Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val
    195             200             205
35 Thr Glu Thr Glu Thr Lys Asn Val Arg Met Met His Ile Lys Asp Lys
    210             215             220
    Phe Arg Tyr Gly Glu Phe Glu Glu Leu Asp Ala Lys Ala Val Glu Leu
    225             230             235             240
    Pro Tyr Arg Asn Ser Asp Leu Ala Met Leu Ile Ile Leu Pro Asn Ser
        245             250             255
40 Lys Thr Gly Leu Pro Thr Leu Glu Glu Lys Leu Gln Asn Val Asp Leu
    260             265             270

```

-124-

Gln Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu
275 280 285

Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys
290 295 300

5 Lys Leu Gly Met Ser Asp Met Phe Met Pro Gly Lys Ala Asp Phe Lys
305 310 315 320

Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val Ile
325 330 335

10 Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu Ala Ala Ala
340 345 350

Ala Thr Gly Val Met Leu Met Met Arg Cys Met Pro Met Met Pro Met
355 360 365

Ala Phe Asn Ala Glu His Pro Phe Leu Tyr Phe Leu His Ser Lys Asn
370 375 380

15 Ser Val Leu Phe Asn Gly Arg Leu Val Lys Pro Thr Thr Glu
385 390 395

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 1492 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

25	TTTTTTTTTT	TTTTTTTTTC	TTAAAGATAT	AATTTAGTAT	ACAACAATTA	TACATAAATT	60
	TTAATTTTTT	TTTATTTTTT	AGTCTTATGT	TTTTATCAA	AATTTGTTAA	AAAAATATTC	120
	ACAAAAATA	AATATATATC	ATAACAATA	ATTTGTA	ACTTCCACC	ACTGAAATAG	180
	TGATGATAAA	AAATACTGAA	TACTTAAAGC	TGACAAATAT	AAATTGAACA	CAATGTTCTA	240
	CAGGCAC	TGTTCAGTAAGC	AATTAAAAA	TATTAGTGAA	GTGCATTGG	CTTTTATTCA	300
30	GTTGTTGGTT	TAACAAGACG	ACCATTGAAT	AGAACAGAA	TTTTGCTGTG	TAAGAAGTAC	360
	AGGAATGGAT	GCTCAGCATT	GAAGGCCATT	GGCATCATTG	GCATACAACG	CATCATTAAC	420
	ATCACGCC	TGTCAGCTGC	AGCTTCAGCA	CCTTCTTCAT	TTACTTCAAT	GAAAGCTTTT	480
	TGAATTACTT	TAGAAATATA	TAACATCTCA	TCAGATCCTT	CAAGCAATCC	TTTGAAATCA	540
	GCTTTTCCAG	GCATGAACAT	ATCAGACATA	CCCAACTTTT	TCAGAGGATC	ATTCAAATTA	600
35	ATTTTCAGACT	CAATTTTGAA	TTTAGGCAGA	TCCAAAATA	CTTCAACAGA	GTACATGCGT	660
	TGAGTCAAGT	TTTGCAAATC	AACATTTTGT	AATTTTCTT	CAAGAGTGGG	GAGACCAGTT	720
	TTGCTGTTTG	GCAAAATGAT	TAACATGGCC	AAATCTGAGT	TCCTGTAGGG	CAATTC	780
	GCCTTGGCAT	CTAATCTTTC	AAATTCTCCA	TAACGGAATT	TATCCTTAAT	GTGCATCATT	840
	CGTACATTCT	TTGTCTCTGT	TTTAGTAACA	TAGAAAGGTT	TGTCTTGGGT	GTTTCTCTTC	900
40	TTGAATTGTT	TCTCCCAAAG	ACCCTTGAAG	TACAATGCAT	TGACAAGAAC	CATTCTTGAA	960
	TCCTGGTCTA	GATCACCGGC	TTTGATCAAA	TCATGAATTT	TGTCATGAGT	TTTTTCTTCA	1020
	ACCCAAGTGT	TGATAACTTT	AGCGCTTTCA	GCATTTTGGG	CAAAGTTCAA	GTTTCTGCT	1080
	CCAGCTAAGA	ATTTGTTGGT	GGCAACTTCT	TTGAAGGTGG	GTTTAAATGT	ATAGCCTTCC	1140
	ATAACATAAA	CTTTATTGGC	AATTTCCAGA	GTTACACCTT	TTTGTGTATT	AAGAGTGTTC	1200
45	ATCAATGCGT	GGTAGTCATC	TTGAATTTTT	TCTTTTGATT	GAGGCTGACG	TAAACGACA	1260
	GCTATTTGTG	TGGCAGTATT	ACCACCAGCT	CCCATTGACA	CCAGGGATAG	AACAGTTTGT	1320
	ACAGACAATG	GGGACATGAT	GAGATTGTCT	TTGTTGCCAG	AAGCAACCGT	ATTGTACAGG	1380
	CTTCCAGCAA	ACTGGTTAAT	ACTTGTAGAC	AATTCCTGGG	GATCCGCCAT	TGTTGAAATT	1440
	GGTATTAACA	CTGATACAAA	AAGAAACACA	AGTCGTGCGT	GTTGAACTAT	CG	1492

-125-

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1194 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

```

10  ATAGTTCAAC ACGCAGCACT TGTGTTTCTT TTTGTATCAG TGTTAATACC AATTTCAACA 60
    ATGGCGGATC CCCAGGAATT GTCTACAAGT ATTAACCAGT TTGCTGGAAG CCTGTACAAT 120
    ACGGTTGCTT CTGGCAACAA AGACAATCTC ATCATGTCCC CATTGTCTGT ACAAACTGTT 180
    CTATCCCTGG TGTCAATGGG AGCTGGTGGT AATACTGCCA CACAAATAGC TGCTGGTTTA 240
    CGTCAGCCTC AATCAAAAAGA AAAAATTCAA GATGACTACC ACGCATTGAT GAACACTCTT 300
    AATACACAAA AAGGTGTAAC TCTGGAAATT GCCAATAAAG TTTATGTTAT GGAAGGCTAT 360
15  ACATTAAAAC CCACCTTCAA AGAAGTTGCC ACCAACAAAT TCTTAGCTGG AGCAGAAAAAC 420
    TTGAACCTTG CCCAAAATGC TGAAAGCGCT AAAGTTATCA ACACTTGGGT TGAAGAAAAA 480
    ACTCATGACA AAATTCATGA TTTGATCAAA GCCGGTGATC TAGACCAGGA TTCAAGAATG 540
    GTTCTTGTC AATGCATTGTA CTTCAAGGGT CTTTGGGAGA AACAATTCAA GAAGGAAAAAC 600
    ACCCAAGACA AACCTTTCTA TGTACTGAA ACAGAGACAA AGAATGTACG AATGATGCAC 660
20  ATTAAGGATA AATTCCGTTA TGGAGAATTT GAAGAATTAG ATGCCAAGGC TGTAGAATTG 720
    CCCTACAGGA ACTCAGATTT GGCCATGTTA ATCATTTTGC CAAACAGCAA AACTGGTCTC 780
    CCCACTCTTG AAGAAAAATT ACAAATGTT GATTTGCAA ACTTGACTCA ACGCATGTAC 840
    TCTGTTGAAG TTATTTTGG TCTGCCTAAA TTCAAAATTG AGTCTGAAAT TAATTTGAAT 900
    GATCCTCTGA AAAAGTTGGG TATGTCTGAT ATGTTTCATC CTGGAAAAGC TGATTTCAAA 960
25  GGATTGCTTG AAGGATCTGA TGAGATGTTA TATATTTCTA AAGTAATTCA AAAAGCTTTC 1020
    ATTGAAGTAA ATGAAGAAG TGCTGAAGCT GCAGCTGCCA CAGGCGTGAT GTTAATGATG 1080
    CGTTGTATGC CAATGATGCC AATGGCCTTC AATGCTGAGC ATCCATTCCT GTACTTCTTA 1140
    CACAGCAAAA ATTCTGTTCT ATTCAATGGT CGTCTTGTTA AACCAACAAC TGAA 1194
  
```

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1194 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

```

30  TTCAGTTGTT GGTTTAACAA GACGACCATT GAATAGAACA GAATTTTTCG TGTGTAAGAA 60
    GTACAGGAAT GGATGCTCAG CATTTGAAGGC CATTGGCATC ATTGGCATAC AACGCATCAT 120
    TAACATCACG CCTGTGGCAG CTGCAGCTTC AGCACCTTCT TCATTTACTT CAATGAAAGC 180
40  TTTTGTGAAT ACTTTAGAAA TATATAACAT CTCATCAGAT CCTTCAAGCA ATCCTTTGAA 240
    ATCAGCTTTT CCAGGCATGA ACATATCAGA CATACCCAAC TTTTTCAGAG GATCATTCAA 300
    ATTAATTTCA GACTCAATTT TGAATTTAGG CAGATCCAAA ATAACCTCAA CAGAGTACAT 360
    GCGTTGAGTC AAGTTTTGCA AATCAACATT TTGTAATTTT TCTTCAAGAG TGGGGAGACC 420
    AGTTTTGCTG TTTGGCAAAA TGATTAACAT GGCCAAATCT GAGTTCCTGT AGGGCAATTC 480
45  TACAGCCTTG GCATCTAATT CTTCAAATTC TCCATAACGG AATTTATCCT TAATGTGCAT 540
    CATTTCGTACA TTCTTTGTCT CTGTTTCAGT AACATAGAAA GGTGTGCTT GGGTGTTTTC 600
    CTTCTTGAAT TGTTTCTCCC AAAGACCTTT GAAGTACAA GCATTGACAA GAACCATTCT 660
    TGAATCCTGG TCTAGATCAC CGGCTTTGAT CAAATCATGA ATTTTGTCT GAGTTTTTTC 720
    TTCAACCCAA GTGTTGATAA CTTTAGCGCT TTCAGCATTT TGGGCAAGT TCAAGTTTTC 780
50  TGCTCCAGCT AAGAATTTGT TGGTGGAAC TTCTTTGAAG GTGGGTTTTC ATGTATAGCC 840
    TTCCATAACA TAACTTTTAT TGGCAATTTT CAGAGTTACA CTTTTTGTG TATTAAGAGT 900
    GTTCATCAAT GCGTGGTAGT CATCTTGAAT TTTTCTTTT GATTGAGGCT GACGTAAACC 960
    AGCAGCTATT TGTGTGGCAG TATTACCACC AGCTCCCAT TACACCAGG ATAGAACAGT 1020
  
```

-126-

TTGTACAGAC AATGGGGACA TGATGAGATT GTCTTTGTTG CCAGAAGCAA CCGTATTGTA 1080
 CAGGCTTCCA GCAAACTGGT TAATACTTGT AGACAATTCC TGGGGATCCG CCATTGTTGA 1140
 AATTGGTATT AACACTGATA CAAAAAGAAA CACAAGTCGT GCGTGTGAA CTAT 1194

(2) INFORMATION FOR SEQ ID NO:30:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 376 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu	1	5	10	15
Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro	20	25	30	
Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	35	40	45	
Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	50	55	60	
Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	65	70	75	80
Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	85	90	95	
Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	100	105	110	
Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	115	120	125	
Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	130	135	140	
Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	145	150	155	160
Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	165	170	175	
Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	180	185	190	
Asn	Val	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	195	200	205	
Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	210	215	220	
Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Thr	225	230	235	240
Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	245	250	255	

-127-

Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile Glu
260 265 270

Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp
275 280 285

5 Met Phe Met Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser
290 295 300

Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu
305 310 315 320

10 Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Gly Val Met Leu
325 330 335

Met Met Arg Cys Met Pro Met Met Pro Met Ala Phe Asn Ala Glu His
340 345 350

Pro Phe Leu Tyr Phe Leu His Ser Lys Asn Ser Val Leu Phe Asn Gly
355 360 365

15 Arg Leu Val Lys Pro Thr Thr Glu
370 375

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 1454 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
25 (A) NAME/KEY: CDS
(B) LOCATION: 20..1210

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GAGCCGAAAT TTTAGCAAA ATG ATT AAC GCA CGA CTT GTG TTT CTT TTT GTA 52
Met Ile Asn Ala Arg Leu Val Phe Leu Phe Val
30 1 5 10

TCA GTG TTA ATA CCA ATT TCA ACA ATG GCG GAT CCC CAG GAA TTG TCT 100
Ser Val Leu Ile Pro Ile Ser Thr Met Ala Asp Pro Gln Glu Leu Ser
15 20 25

ACA AGT ATT AAC CAG TTT GCT GGA AGC CTG TAC AAT ACG GTT GCT TCT 148
Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser
35 30 35 40

GGC AAC AAA GAC AAT CTC ATC ATG TCC CCA TTG TCT GTA CAA ACT GTT 196
Gly Asn Lys Asp Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val
45 50 55

40 CTA TCC CTG GTG TCA ATG GGA GCT GGT GGT AAT ACT GCC ACA CAA ATA 244
Leu Ser Leu Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile
60 65 70 75

-128-

	GCT	GCT	GGT	TTA	CGT	CAG	CCT	CAA	TCA	AAA	GAA	AAA	ATT	CAA	GAT	GAC	292
	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	Glu	Lys	Ile	Gln	Asp	Asp	
					80					85					90		
5	TAC	CAT	GCA	TTG	ATG	AAC	ACT	CTT	AAT	ACA	CAA	AAA	GGT	GTA	ACT	CTG	340
	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	Gln	Lys	Gly	Val	Thr	Leu	
				95					100					105			
	GAA	ATT	GCC	AAC	AAA	GTT	TAC	GTT	ATG	GAA	GGC	TAT	ACA	TTG	AAA	CCC	388
	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	Gly	Tyr	Thr	Leu	Lys	Pro	
			110					115					120				
10	ACC	TTC	AAA	GAA	GTT	GCC	ACC	AAC	AAA	TTC	TTA	GCT	GGA	GCA	GAA	AAC	436
	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	Leu	Ala	Gly	Ala	Glu	Asn	
		125					130					135					
	TTG	AAC	TTT	GCC	CAA	AAT	GCT	GAA	AGC	GCT	AAA	GTT	ATC	AAC	ACT	TGG	484
15	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	Lys	Val	Ile	Asn	Thr	Trp	
	140					145					150					155	
	GTT	GAA	GAA	AAA	ACT	CAT	GAC	AAA	ATT	CAT	GAT	TTG	ATC	AAA	GCC	GGT	532
	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	Asp	Leu	Ile	Lys	Ala	Gly	
					160					165					170		
20	GAT	CTA	GAC	CAG	GAT	TCA	AGA	ATG	GTT	CTT	GTC	AAT	GCA	TTG	TAC	TTC	580
	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	Val	Asn	Ala	Leu	Tyr	Phe	
				175					180					185			
	AAG	GGT	CTT	TGG	GAG	AAA	CAA	TTC	AAG	AAG	GAA	AAC	ACT	CAA	GAC	AAA	628
	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	Glu	Asn	Thr	Gln	Asp	Lys	
			190					195					200				
25	CCT	TTC	TAT	GTT	ACT	GAA	ACA	GAG	ACA	AAG	AAT	GTA	CGA	ATG	ATG	CAC	676
	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg	Met	Met	His	
		205					210					215					
	ATT	AAG	GAT	AAA	TTC	CGT	TAT	GGA	GAA	TTT	GAA	GAA	TTA	GAT	GCC	AAG	724
30	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu	Glu	Leu	Asp	Ala	Lys	
	220					225					230					235	
	GCT	GTA	GAA	TTG	CCC	TAC	AGG	AAC	TCA	GAT	TTG	GCC	ATG	TTA	ATC	ATT	772
	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	Leu	Ala	Met	Leu	Ile	Ile	
					240					245					250		
35	TTG	CCA	AAC	AGC	AAA	ACT	GGT	CTC	CCC	GCT	CTT	GAA	GAA	AAA	TTA	CAA	820
	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	
				255					260					265			
	AAT	GTT	GAC	TTG	CAA	AAC	TTG	ACT	CAA	CGC	ATG	TAC	TCT	GTT	GAA	GTT	868
	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	
			270					275					280				
40	ATT	TTG	GAT	CTG	CCT	AAA	TTC	AAG	ATT	GAA	TCT	GAA	ATT	AAT	TTG	AAT	916
	Ile	Leu	Asp	Leu	Pro	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	
		285					290					295					
45	GAT	CCT	CTG	AAA	AAG	TTG	GGT	ATG	TCT	GAT	ATG	TTT	GTT	CCT	GGA	AAA	964
	Asp	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	Met	Phe	Val	Pro	Gly	Lys	
	300					305					310					315	

-129-

GCT GAT TTC AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT 1012
 Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile
 320 325 330

TCT AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT GCT 1060
 5 Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala
 335 340 345

GAA GCT GCA GCT GCC ACA GCT ACC TTT ATG GTT ACC TAT GAA CTG GAG 1108
 Glu Ala Ala Ala Ala Thr Ala Thr Phe Met Val Thr Tyr Glu Leu Glu
 350 355 360

GTT TCC CTG GAT GAT CCA ACC GTT TTT AAA GTC GAT CAT CCA TTC AAT 1156
 10 Val Ser Leu Asp Asp Pro Thr Val Phe Lys Val Asp His Pro Phe Asn
 365 370 375

ATT GTT TTG AAG ACA GGT GAT ACT GTA ATT TTT AAT GGG CGA GTT CAA 1204
 15 Ile Val Leu Lys Thr Gly Asp Thr Val Ile Phe Asn Gly Arg Val Gln
 380 385 390 395

ACT CTA TGA AATGGATAGT GTAAGAAAAG AATACAAGAT CTATCTGAAT CTCTGGATTA 1263
 Thr Leu

ATGAAGTAAT TTTTCTACAA TATTTTAA TAGTTATTAG GTCTAAAATA AGTTCATTTT 1323
 TTAGTATGTG GTATAAATCG TGTAGACGAA AAATGTTTTG TTTTAGTTT CACTTTTTAT 1383
 20 GAATGTAATC ACCTATATAA TGTGTAGTT TATGTAATAA AAATGTTAAA TGTGAAAAAA 1443
 AAAAAAAAAA A 1454

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 397 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

30 Met Ile Asn Ala Arg Leu Val Phe Leu Phe Val Ser Val Leu Ile Pro
 1 5 10 15

Ile Ser Thr Met Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln
 20 25 30

Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn
 35 40 45

35 Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu Ser Leu Val Ser
 50 55 60

Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile Ala Ala Gly Leu Arg
 65 70 75 80

40 Gln Pro Gln Ser Lys Glu Lys Ile Gln Asp Asp Tyr His Ala Leu Met
 85 90 95

Asn Thr Leu Asn Thr Gln Lys Gly Val Thr Leu Glu Ile Ala Asn Lys
 100 105 110

Val Tyr Val Met Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val
 115 120 125

-130-

Ala Thr Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln
 130 135 140

Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu Lys Thr
 145 150 155 160

5 His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp Leu Asp Gln Asp
 165 170 175

Ser Arg Met Val Leu Val Asn Ala Leu Tyr Phe Lys Gly Leu Trp Glu
 180 185 190

10 Lys Gln Phe Lys Lys Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr
 195 200 205

Glu Thr Glu Thr Lys Asn Val Arg Met Met His Ile Lys Asp Lys Phe
 210 215 220

Arg Tyr Gly Glu Phe Glu Glu Leu Asp Ala Lys Ala Val Glu Leu Pro
 225 230 235 240

15 Tyr Arg Asn Ser Asp Leu Ala Met Leu Ile Ile Leu Pro Asn Ser Lys
 245 250 255

Thr Gly Leu Pro Ala Leu Glu Glu Lys Leu Gln Asn Val Asp Leu Gln
 260 265 270

20 Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro
 275 280 285

Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys
 290 295 300

Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly
 305 310 315 320

25 Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln
 325 330 335

Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Ala
 340 345 350

30 Thr Ala Thr Phe Met Val Thr Tyr Glu Leu Glu Val Ser Leu Asp Asp
 355 360 365

Pro Thr Val Phe Lys Val Asp His Pro Phe Asn Ile Val Leu Lys Thr
 370 375 380

Gly Asp Thr Val Ile Phe Asn Gly Arg Val Gln Thr Leu
 385 390 395

35 (2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1454 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40

- (ii) MOLECULE TYPE: cDNA

-131-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

	TTTTTTTTTT	TTTTTTTCAC	ATTTAACATT	TTTATTACAT	AAACTACAAC	ATTATATAGG	60
	TGATTACATT	CATAAAAAGT	GAAAACATAA	ACAAAACATT	TTTCGTCTAC	ACGATTTATA	120
	CCACATACTA	AAAAATGAAC	TTATTTTAGA	CCTAATAACT	ATTAAAAAAT	ATTGTAGAAA	180
5	AATTACTTCA	TTAATCCAGA	GATTCAGATA	GATCTTGAT	TCTTTTCTTA	CACTATCCAT	240
	TTCATAGAGT	TTGAACTCGC	CCATTAATAA	TTACAGTATC	ACCTGTCTTC	AAAACAATAT	300
	TGAATGGATG	ATCGACTTTA	AAAACGGTTG	GATCATCCAG	GGAAACCTCC	AGTTCATAGG	360
	TAACCATAAA	GGTAGCTGTG	GCAGCTGCAG	CTTCAGCACC	TTCTTCATTT	ACTTCAATGA	420
	AAGCTTTTTG	AATTACTTTA	GAAATATATA	ACATCTCATC	AGATCCTTCA	AGCAATCCTT	480
10	TGAAATCAGC	TTTTCCAGGA	ACAAACATAT	CAGACATACC	CAACTTTTTC	AGAGGATCAT	540
	TCAAATTAAT	TTTCAAGTCA	ATCTTGAATT	TAGGCAGATC	CAAAATAACT	TCAACAGAGT	600
	ACATGCGTTG	AGTCAAGTTT	TGCAAGTCAA	CATTTTGTA	TTTTTCTTCA	AGAGCGGGGA	660
	GACCAGTTTT	GCTGTTTGGC	AAAATGATTA	ACATGGCCAA	ATCTGAGTTC	CTGTAGGGCA	720
	ATTCTACAGC	CTTGGCATCT	AATTCCTTCA	ATTCTCCATA	ACGGAATTTA	TCCTTAATGT	780
15	GCATCATTTC	TATTTCTTTT	GTCTCTGTTT	CAGTAACATA	GAAAGGTTTG	TCTTGAGTGT	840
	TTTCCTTCTT	GAATTGTTTC	TCCCAAAGAC	CCTTGAAGTA	CAATGCATTG	ACAAGAACCA	900
	TTCTTGAATC	CTGGTCTAGA	TCACCGGCTT	TGATCAAATC	ATGAATTTTG	TCATGAGTTT	960
	TTTCTTCAAC	CCAAGTGTTC	ATAACTTTAG	CGCTTTCAGC	ATTTTGGGCA	AAGTTCAGAT	1020
	TTTCTGCTCC	AGCTAAGAAT	TTGTTGGTGG	CAACTTCCTT	GAAGGTGGGT	TTCAATGTAT	1080
20	AGCCTTCCAT	AACGTAAACT	TTGTTGGCAA	TTTCCAGAGT	TACACCTTTT	TGTGTATTAA	1140
	GAGTGTTTAT	CAATGCATGG	TAGTCATCTT	GAATTTTTTC	TTTTGATTGA	GGCTGACGTA	1200
	AACGACGAGC	TATTTGTGTG	GCAGTATTAC	CACCAGCTCC	CATTGACACC	AGGGATAGAA	1260
	CAGTTTGTAC	AGACAATGGG	GACATGATGA	GATTGTCTTT	GTTGCCAGAA	GCAACCGTAT	1320
	TGTACAGGCT	TCCAGCAAAC	TGGTTAATAC	TTGTAGACAA	TTCTTGGGGA	TCCGCCATTG	1380
25	TTGAAATTGG	TATTAACACT	GATACAAAAA	GAAACACAAG	TCGTGCGTTA	ATCATTTTGC	1440
	TAAAATTTTC	GCTC					1454

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

30	(A) LENGTH: 1191 nucleotides
	(B) TYPE: nucleic acid
	(C) STRANDEDNESS: single
	(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

35	ATGATTAACG	CACGACTTGT	GTTTCTTTTT	GTATCAGTGT	TAATACCAAT	TTCAACAATG	60
	GCGGATCCCC	AGGAATGTGC	TACAAGTATT	AACCAGTTTG	CTGGAAGCCT	GTACAATACG	120
	GTTGCTTCTG	GCAACAAAGA	CAATCTCATC	ATGTCCCAT	TGTCTGTACA	AACTGTTCTA	180
	TCCCTGGTGT	CAATGGGAGC	TGGTGGTAAT	ACTGCCACAC	AAATAGCTGC	TGGTTTACGT	240
	CAGCCTCAAT	CAAAAGAAAA	AATTCAAGAT	GACTACCATG	CATTGATGAA	CACTCTTAAT	300
40	ACACAAAAAG	GTGTAACCTC	GGAAATTGCC	AACAAAGTTT	ACGTTATGGA	AGGCTATACA	360
	TTGAAACCCA	CCTTCAAAGA	AGTTGCCACC	AACAAATCTT	TAGCTGGAGC	AGAAAACTTG	420
	AACTTTGCCC	AAAATGCTGA	AAGCGCTAAA	GTTATCAACA	CTTGGGTTGA	AGAAAAAAT	480
	CATGACAAAA	TTTATGATTT	GATCAAAGCC	GGTGATCTAG	ACCAGGATTC	AAGAATGGTT	540
	CTTGTCAATG	CATTGTACTT	CAAGGGTCTT	TGGGAGAAAC	AATTCAAGAA	GGAAAAACAT	600
45	CAAGACAAAC	CTTTCTATGT	TACTGAAACA	GAGACAAAGA	ATGTACGAAT	GATGCACATT	660
	AAGGATAAAT	TCCGTTATGG	AGAATTTGAA	GAATTAGATG	CCAAGGCTGT	AGAATTGCCC	720
	TACAGGAACT	CAGATTTGGC	CATGTTAATC	ATTTTGCCAA	ACAGCAAAAC	TGGTCTCCCC	780
	GCTCTTGAAG	AAAAATTACA	AAATGTTGAC	TTGCAAAACT	TGACTCAACG	CATGTACTCT	840
	GTTGAAGTTA	TTTTTGATCT	GCCTAAATTC	AAGATTGAAT	CTGAAATTAA	TTTGAATGAT	900
50	CCTCTGAAAA	AGTTGGGTAT	GTCTGATATG	TTTGTTCCTG	GAAAAGCTGA	TTTCAAAGGA	960
	TTGCTTGAAG	GATCTGATGA	GATGTTATAT	ATTTCTAAAG	TAATTCAAAA	AGCTTTTCATT	1020
	GAAGTAAATG	AAGAAGGTGC	TGAAGCTGCA	GCTGCCACAG	CTACCTTTAT	GGTTACCTAT	1080
	GAACCTGGAG	TTTCCCTGGA	TGATCCAACC	GTTTTTAAAG	TCGATCATCC	ATTCAATATT	1140
	GTTTTGAAGA	CAGGTGATAC	TGTAATTTTT	AATGGGCGAG	TTCAAACTCT	A	1191

-132-

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1191 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

```

10 TAGAGTTTGA ACTCGCCCAT TAAAAATTAC AGTATCACCT GTCTTCAAAA CAATATTGAA 60
   TGGATGATCG ACTTTAAAAA CGGTTGGATC ATCCAGGGAA ACCTCCAGTT CATAGGTAAC 120
   CATAAAGGTA GCTGTGGCAG CTGCAGCTTC AGCACCTTCT TCATTTACTT CAATGAAAGC 180
   TTTTGAATT ACTTTAGAAA TATATAACAT CTCATCAGAT CCTTCAAGCA ATCCTTTGAA 240
   ATCAGCTTTT CCAGGAACAA ACATATCAGA CATACCCAAC TTTTTCAGAG GATCATTCAA 300
   ATTAATTTCA GATTCAATCT TGAATTTAGG CAGATCCAAA ATAACCTCAA CAGAGTACAT 360
15 GCGTTGAGTC AAGTTTTGCA AGTCAACATT TTGTAATTTT TCTTCAAGAG CGGGGAGACC 420
   AGTTTTGCTG TTTGGCAAAA TGATTAACAT GGCCAAATCT GAGTTCCTGT AGGGCAATTC 480
   TACAGCCTTG GCATCTAATT CTTCAAATTC TCCATAACGG AATTATCCT TAATGTGCAT 540
   CATTCGTACA TTCTTTGTCT CTGTTTCAGT AACATAGAAA GGTTTGTCTT GAGTGTTTTC 600
   CTTCTTGAAT TGTTTCTCCC AAAGACCTTT GAAGTACAAT GCATTGACAA GAACCATTCT 660
20 TGAATCCTGG TCTAGATCAC CGGCTTTGAT CAAATCATGA ATTTTGTTCAT GAGTTTTTTC 720
   TTCAACCCAA GTGTTGATAA CTTTAGCGCT TTCAGCATTT TGGGCAAAGT TCAAGTTTTTC 780
   TGCTCCAGCT AAGAATTTGT TGGTGCAAC TTCTTTGAAG GTGGGTTTCA ATGTATAGCC 840
   TTCCATAACG TAAACTTTGT TGGCAATTTT CAGAGTTACA CCTTTTGTG TATTAAGAGT 900
   GTTCATCAAT GCATGGTAGT CATCTTGAAT TTTTCTTTT GATTGAGGCT GACGTAAACC 960
25 AGCAGCTAAT TGTGTGGCAG TATTACCACC AGTCCCATT GACACCAGG ATAGAACAGT 1020
   TTGTACAGAC AATGGGGACA TGATGAGATT GTCTTTGTTG CCAGAAGCAA CCGTATTGTA 1080
   CAGGCTTCCA GCAAATGGT TAATACTTGT AGACAATTCC TGGGGATCCG CCATTGTTGA 1140
   AATTGGTATT AACACTGATA CAAAAGAAA CACAAGTCGT GCGTTAATCA T 1191

```

(2) INFORMATION FOR SEQ ID NO:36:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 376 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

```

Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu
 1           5           10           15
Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser Pro
          20           25           30
40 Leu Ser Val Gln Thr Val Leu Ser Leu Val Ser Met Gly Ala Gly Gly
          35           40           45
Asn Thr Ala Thr Gln Ile Ala Ala Gly Leu Arg Gln Pro Gln Ser Lys
          50           55           60
45 Glu Lys Ile Gln Asp Asp Tyr His Ala Leu Met Asn Thr Leu Asn Thr
          65           70           75           80
Gln Lys Gly Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val Met Glu
          85           90           95

```

-133-

Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr Asn Lys Phe
 100 105 110
 Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln Asn Ala Glu Ser Ala
 115 120 125
 5 Lys Val Ile Asn Thr Trp Val Glu Glu Lys Thr His Asp Lys Ile His
 130 135 140
 Asp Leu Ile Lys Ala Gly Asp Leu Asp Gln Asp Ser Arg Met Val Leu
 145 150 155 160
 10 Val Asn Ala Leu Tyr Phe Lys Gly Leu Trp Glu Lys Gln Phe Lys Lys
 165 170 175
 Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr Glu Thr Glu Thr Lys
 180 185 190
 Asn Val Arg Met Met His Ile Lys Asp Lys Phe Arg Tyr Gly Glu Phe
 195 200 205
 15 Glu Glu Leu Asp Ala Lys Ala Val Glu Leu Pro Tyr Arg Asn Ser Asp
 210 215 220
 Leu Ala Met Leu Ile Ile Leu Pro Asn Ser Lys Thr Gly Leu Pro Ala
 225 230 235 240
 20 Leu Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln Arg
 245 250 255
 Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile Glu
 260 265 270
 Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp
 275 280 285
 25 Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser
 290 295 300
 Asp Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu
 305 310 315 320
 30 Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Ala Thr Phe Met
 325 330 335
 Val Thr Tyr Glu Leu Glu Val Ser Leu Asp Asp Pro Thr Val Phe Lys
 340 345 350
 Val Asp His Pro Phe Asn Ile Val Leu Lys Thr Gly Asp Thr Val Ile
 355 360 365
 35 Phe Asn Gly Arg Val Gln Thr Leu
 370 375

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 21 bases
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

-134-

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GTGTTTCTTT TTGTATCAGT G 21

(2) INFORMATION FOR SEQ ID NO:38:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CGGAATTCTT TAAAGGATT TAACAC 26

(2) INFORMATION FOR SEQ ID NO:39:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

CGGAATTCTA ATTGGTAAAT CTC 23

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 25 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

30 CGGAATTCTT TTATTCAGTT GTTGG 25

(2) INFORMATION FOR SEQ ID NO:41:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 23 bases

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

CGGAATTCAT AGAGTTTGAA CTC 23

-135-

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

CAAAACTGGT CTCCCCGCTC 20

10 (2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

ATTACAAAAT GTTGACTTGC 20

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

TAATACGACT CACTATAGGG 20

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 549 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 3..404
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

40 TT GAA GAA AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT CAA 44
 Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln
 1 5 10

-136-

	CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC	86
	Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe	
	15 20 25	
5	AAG ATT GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG	128
	Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys	
	30 35 40	
10	TTG GGT ATG TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC	170
	Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe	
	45 50 55	
15	AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT	212
	Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser	
	60 65 70	
	AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT	254
	Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly	
	75 80	
20	GCT GAA GCT GCA GCT GCC ACA GGA GGT TTC ATA ATG GCC GTA	296
	Ala Glu Ala Ala Ala Ala Thr Gly Gly Phe Ile Met Ala Val	
	85 90 95	
25	TCC TTA CCT TTA CCA CCT GAG ACT TTT AAT GCT GAC CAT CCC	338
	Ser Leu Pro Leu Pro Pro Glu Thr Phe Asn Ala Asp His Pro	
	100 105 110	
30	TTC TAT TTT GTG ATC TTC GAC AAA TCT TCC AAA GTG ACA ATG	380
	Phe Tyr Phe Val Ile Phe Asp Lys Ser Ser Lys Val Thr Met	
	115 120 125	
35	TTC CAT GGT CAA CAC GTT AAT CCT TAA GAGTAACAAG GCAAATTTTG	427
	Phe His Gly Gln His Val Asn Pro	
	130	
	ATAATTAATT GTGATAAATT GCACGTTGTA AAAATGCTTC TTGATGCATA	477
	TTTGATAATA TAATGTAAAG CCAAAAAAAAAA AAAAAAATAA AAAAACTCGA	527
	GGGGGGCCCG GTACCCAATT CG	549

(2) INFORMATION FOR SEQ ID NO:46:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 134 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- 45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:
- | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Glu | Lys | Leu | Gln | Asn | Val | Asp | Leu | Gln | Asn | Leu | Thr | Gln |
| 1 | | | | 5 | | | | 10 | | | | | |
| Arg | Met | Tyr | Ser | Val | Glu | Val | Ile | Leu | Asp | Leu | Pro | Lys | Phe |
| 15 | | | | 20 | | | | 25 | | | | | |
| Lys | Ile | Glu | Ser | Glu | Ile | Asn | Leu | Asn | Asp | Pro | Leu | Lys | Lys |
| 30 | | | | 35 | | | | 40 | | | | | |
| Leu | Gly | Met | Ser | Asp | Met | Phe | Val | Pro | Gly | Lys | Ala | Asp | Phe |
| 45 | | | | 50 | | | | 55 | | | | | |

-137-

Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser
 60 65 70

5 Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly
 75 80

Ala Glu Ala Ala Ala Ala Thr Gly Gly Phe Ile Met Ala Val
 85 90 95

10 Ser Leu Pro Leu Pro Pro Glu Thr Phe Asn Ala Asp His Pro
 100 105 110

15 Phe Tyr Phe Val Ile Phe Asp Lys Ser Ser Lys Val Thr Met
 115 120 125

Phe His Gly Gln His Val Asn Pro
 130

(2) INFORMATION FOR SEQ ID NO:47:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 549 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- 25 (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

	CGAATTGGGT	ACCGGGCCCC	CCTCGAGTTT	TTTTTTTTTT	TTTTTTTTTT	50
	GGCTTTACAT	TATATTATCA	AATATGCATC	AAGAAGCATT	TTTACAACGT	100
	GCAATTTATC	ACAATTAATT	ATCAAAATTT	GCCTTGTTAC	TCTTAAGGAT	150
30	TAACGTGTTG	ACCATGGAAC	ATTGTCACCT	TGGAAGATTT	GTCGAAGATC	200
	ACAAAATAGA	AGGGATGGTC	AGCATTAAAA	GTCTCAGGTG	GTAAAGGTAA	250
	GGATACGGCC	ATTATGAAAC	CTCCTGTGGC	AGCTGCAGCT	TCAGCACCTT	300
	CTTCATTTAC	TTCAATGAAA	GCTTTTGTGA	TTACTTTAGA	AATATATAAC	350
	ATCTCATCAG	ATCCTTCAAG	CAATCCTTTG	AAATCAGCTT	TTCCAGGAAC	400
35	AAACATATCA	GACATACCCA	ACTTTTTCAG	AGGATCATTC	AAATTAATTT	450
	CAGATTCAAT	CTTGAATTTA	GGCAGATCCA	AAATAACTTC	AACAGAGTAC	500
	ATGCGTTGAG	TCAAGTTTGT	CAAGTCAACA	TTTTGTAATT	TTTCTTCAA	549

(2) INFORMATION FOR SEQ ID NO:48:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 549 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- 45 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 3..449
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

	TT	GAA	GAA	AAA	TTA	CAA	AAT	GTT	GAC	TTG	CAA	AAC	TTG	ACT	CAA	44
50	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln		
	1				5							10				

-138-

	CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC	88
	Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe	
	15 20 25	
5	AAG ATT GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG	128
	Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys	
	30 35 40	
10	TTG GGT ATG TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC	170
	Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe	
	45 50 55	
15	AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT	212
	Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser	
	60 65 70	
20	AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT	254
	Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly	
	75 80	
25	GCT GAA GCT GCA GCT GCC ACA GGA GTG CTC ATA GAA TTG GAC	296
	Ala Glu Ala Ala Ala Thr Gly Val Leu Ile Glu Leu Asp	
	85 90 95	
30	TCT TTT ATG CCT GAT CGA GTA TTT GAA GCA AAT CAT CCC TTC	338
	Ser Phe Met Pro Asp Arg Val Phe Glu Ala Asn His Pro Phe	
	100 105 110	
35	TAT TTC GCC CTC TAC ACA AAA TCT GCA CAA AAA CCA GAA CAA	380
	Tyr Phe Ala Leu Tyr Thr Lys Ser Ala Gln Lys Pro Glu Gln	
	115 120 125	
40	TCC AAA AAG CGA GCG CGC TCT AAA ATT GTT ACA GTA CTG TTT	422
	Ser Lys Lys Arg Ala Arg Ser Lys Ile Val Thr Val Leu Phe	
	130 135 140	
45	TCT GGA CGT TTA ACC AAT ATT AAT AAC TAGAATAATA TGGAATTCTA	469
	Ser Gly Arg Leu Thr Asn Ile Asn Asn	
	145	
	TTTTTGTGAA ATAAACAGGA TAAATAATGA AGTAAAAAAA AAAAAAAAAA	519
	AAC TCGAGGG GGGGCCCGGT ACCCAATTCTG	549

(2) INFORMATION FOR SEQ ID NO:49:

45	(i) SEQUENCE CHARACTERISTICS:
	(A) LENGTH: 149 amino acids
	(B) TYPE: amino acid
	(D) TOPOLOGY: linear
	(ii) MOLECULE TYPE: Protein
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:
50	Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln
	1 5 10
	Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe
	15 20 25
55	

30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 549 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

35	CGAATTGGGT	ACCGGGCCCC	CCCTCGAGTT	TTTTTTTTTT	TTTTTTTACT	50
	TCATTATTTA	TCCTGTTTAT	TTCACAAAAA	TAGAATTCCA	TATTATTCTA	100
	GTTATTAATA	TTGGTTAAAC	GTCCAGAAAA	CAGTACTGTA	ACAAATTTAG	150
	AGCGCGCTCG	CTTTTTGGAT	TGTTCTGGTT	TTTGTGCAGA	TTTTGTGTAG	200
	AGGGCGAAAT	AGAAGGGATG	ATTTGCTTCA	AATACTCGAT	CAGGCATAAA	250
40	AGAGTCCAA	TCTATGAGCA	CTCCTGTGGC	AGCTGCAGCT	TCAGCACCTT	300
	CTTCATTTAC	TTCAATGAAA	GCTTTTTTGAA	TTACTTTTGA	AATATATAAC	350
	ATCTCATCAG	ATCCTTCAAG	CAATCCTTTG	AAATCAGCTT	TTCCAGGAAC	400
	AAACATATCA	GACATACCCA	ACTTTTTTCAG	AGGATCATTC	AAATTAATTT	450
	CAGATTTCAAT	CTTGAATTTA	GGCAGATCCA	AAATAACTTC	AACAGAGTAC	500
45	ATGCGTTGAG	TCAAGTTTGT	CAAGTCAACA	TTTTGTAAAT	TTTCTTCAA	549

50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 581 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-140-

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 3..410

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

```

5  TT GAA GAA AAA TTA CAA AAT GTT GAT TTG CAA AAC TTG ACT CAA 44
   Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln
    1             5             10

10 CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC 86
   Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe
    15             20             25

   AAA ATT GAG TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG 128
   Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys
    30             35             40

   TTG GGT ATG TCT GAT ATG TTC ATG CCT GGA AAA GCT GAT TTC 170
   Leu Gly Met Ser Asp Met Phe Met Pro Gly Lys Ala Asp Phe
    45             50             55

20  AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT 212
   Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser
    60             65             70

25  AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT 254
   Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly
    75             80

30  GCT GAA GCT GCA GCT GCC ACA GCT GTC TTA GCA GTG GCT TTT 296
   Ala Glu Ala Ala Ala Ala Thr Ala Val Leu Ala Val Ala Phe
    85             90             95

   TCA CTG AGT TTT CCT GCA GAT CCT GTG CTT TTC ACG GCT GAT 338
   Ser Leu Ser Phe Pro Ala Asp Pro Val Leu Phe Thr Ala Asp
    100            105            110

35  CAT CCT TTC CAT TAT TTG CTA ATA GAT CGA TCT CAA CAT AAT 380
   His Pro Phe His Tyr Leu Leu Ile Asp Arg Ser Gln His Asn
    115            120            125

40  CTA CCT CTT TTT AAA GGA CGA TTT GTG CAA TAA TCCATTGGA 423
   Leu Pro Leu Phe Lys Gly Arg Phe Val Gln
    130            135

45  TTAAACATAT TATTGATCAC TTGTGTGTTT TAATTTAATG CATTTTATT 473
   TGTTAATGTT GCCCCAAATA TTAGCAATTT GTATTAAAT AAATTTATT 523
   CGTGCTTGTT ATAAAAAAAA AAAAAAAAAA CTCGAGGGGG GGCCCGGTAC 573
   CCAATTTCG 581

```

(2) INFORMATION FOR SEQ ID NO:52:

(i) SEQUENCE CHARACTERISTICS:

50 (A) LENGTH: 136 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Protein

-141-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln
 1 5 10

5 Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe
 15 20 25

Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys
 30 35 40

10 Leu Gly Met Ser Asp Met Phe Met Pro Gly Lys Ala Asp Phe
 45 50 55

Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser
 15 60 65 70

Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly
 75 80

20 Ala Glu Ala Ala Ala Thr Ala Val Leu Ala Val Ala Phe
 85 90 95

Ser Leu Ser Phe Pro Ala Asp Pro Val Leu Phe Thr Ala Asp
 100 105 110

25 His Pro Phe His Tyr Leu Leu Ile Asp Arg Ser Gln His Asn
 115 120 125

Leu Pro Leu Phe Lys Gly Arg Phe Val Gln
 30 130 135

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 581 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CGAATTGGGT ACCGGGCCCC CCCTCGAGTT TTTTTTTTTT TTTTATTATAA 50

40 CAAGCACGAA ATAAATTTAT TTAAATACAA ATTGCTAATA TTTGGGGCAA 100

CATTAACAAA TAAAAATGCA TTAAATTAAA ACACACAAGT GATCAATAAT 150

ATGTTAAATC CAAATGGATT ATTGCACAAA TCGTCCTTTA AAAAGAGGTA 200

GATTATGTTG AGATCGATCT ATTAGCAAAT AATGGAAAGG ATGATCAGCC 250

GTGAAAAGCA CAGGATCTGC AGGAAAACCTC AGTGAAAAG CCACTGCTAA 300

45 GACAGCTGTG GCAGCTGCAG CTTCAGCACC TTCTTCATTT ACTTCAATGA 350

AAGCTTTTGT AATTACTTTA GAAATATATA ACATCTCATC AGATCCTTCA 400

AGCAATCCTT TGAAATCAGC TTTTCCAGGC ATGAACATAT CAGACATACC 450

CAACTTTTTC AGAGGATCAT TCAAATTAAT TTCAGACTCA ATTTTGAATT 500

TAGGCAGATC CAAAATAACT TCAACAGAGT ACATGCGTTG AGTCAAGTTT 550

50 TGCAAATCAA CATTTTGTAA TTTTCTTCA A 581

-143-

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 118 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp
 1 5 10
 Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp
 15 20 25
 Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Val Pro Gly
 15 30 35 40
 Lys Ala Asp Phe Lys Gly Leu Glu Gly Ser Asp Glu Met
 45 50 55
 Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val
 60 65 70
 Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Glu Tyr Cys
 75 80
 Ser Leu Asn Trp Ser Arg Ile Leu Tyr Val Leu Leu Gln Arg
 85 90 95
 Phe Ser Lys Leu Ile Thr Pro Phe Pro Phe Tyr His Lys Asp
 100 105 110
 Phe Glu His Thr Phe Val
 115

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 654 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CGAATTGGGT ACCGGGCCCC CCCTCGAGTT TTTTTTTTTT TTTTTCACAT 50
 TTAACATTTT TATTACATAA ACTACAACAT TATATAGGTG ATTACATTCA 100
 TAAAAAGTGA AAACATAAAC AAAACATTTT TCGTCTACAC GATTTATAACC 150
 ACATACTAAA AAATGAACTT ATTTTAGACC TAATAACTAT TAAAAAATAT 200
 TGTAGAAAAA TTAATTCATT AATCCAGAGA TTCAGATAGA TCTTGGAATC 250
 CTTCGTTTAA GAAAATTAGC TTTTCATGGC GTTCTGACGC GCCCATCAAA 300
 CAAAAGTGTG TTCGAAGTCC TTATGATAAA ATGGGAAAGG GGTGATCAAC 350
 TTTGAAAACC TTTGGAGGAG GACGTACAAT ATACGAGACC AGTTCAGGGA 400
 GCAGTACTCT GTGGCAGCTG CAGCTTCAGC ACCTTCTTCA TTTACTTCAA 450
 TGAAAGCTTT TTGAATTACT TTAGAAATAT ATAACATCTC ATCAGATCCT 500
 TCAAGCAATC CTTTGAAATC AGCTTTTCCA GGAACAAACA TATCAGACAT 550
 ACCCAACTTT TTCAGAGGAT CATTCAAATT AATTTAGAT TCAATCTTGA 600

-144-

ATTTAGGCAG ATCCAAAATA ACTTCAACAG AGTACATGCG TTGAGTCAAG 650
TTTT 654

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:
5 (A) LENGTH: 670 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 3..377

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

15	AA AAC TTG ACT CAA CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT	44
	Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp	
	1 5 10	
	CTG CCT AAA TTC AAG ATT GAA TCT GAA ATT AAT TTG AAT GAT	86
	Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp	
	15 20 25	
20	CCT CTG AAA AAG TTG GGT ATG TCT GAT ATG TTC ATG CCT GGA	128
	Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Met Pro Gly	
	30 35 40	
25	AAA GCT GAT TTC AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG	170
	Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu Met	
	45 50 55	
30	TTA TAT ATT TCT AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA	212
	Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val	
	60 65 70	
	AAT GAA GAA GGT GCT GAA GCT GCA GCT GCC ACA GGT GTA ATT	254
	Asn Glu Glu Gly Ala Glu Ala Ala Ala Ala Thr Gly Val Ile	
35	75 80	
	ATG GTT GCA TTT ATG TCG TAT ATC GTA CCA CCT CCT CCA ACC	296
	Met Val Ala Phe Met Ser Tyr Ile Val Pro Pro Pro Pro Thr	
	85 90 95	
40	ATT TTT AAA GTT GAT CAT CCT TTC CAC TTT GTC TTA AAG ACT	338
	Ile Phe Lys Val Asp His Pro Phe His Phe Val Leu Lys Thr	
	100 105 110	
45	TCG GAT ACT GTT TTG TTT GAT GGG AGG GTT CGA CTT CCA TAA	380
	Ser Asp Thr Val Leu Phe Asp Gly Arg Val Arg Leu Pro	
	115 120 125	
	ATGATAATGA TGTGATTTTC TTAAATAAAA GAATACAAGA TCTATCTGAA	430
	TCTCCAGATT AATGAAGTAA TTTTCTACA ATATTTTSTA ATAGTTATTA	480
50	GGTCTAAAAT AAGTTCATTT TTTAGTATGT GGTATAAATC GTGTAGACGA	530
	AAAATGTTTT GTTTTAGTTT TCACTTTTGA TGAATGTAAT CACCTATATA	580
	ATGTTGTAGT TTATGTAATA AAAATGTTAA ATGTGAAAAA AAAAAAAAAA	630
	AAAAAAAAAA AACTCGAGGG GGGGCCCGGT ACCCAATTTCG	670

-145-

(2) INFORMATION FOR SEQ ID NO:58:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 125 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

```

Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp
 1             5             10
10 Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp
   15             20             25
    Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Met Pro Gly
      30             35             40
15 Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu Met
   45             50             55
    Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val
  20             60             65             70
    Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Gly Val Ile
      75             80
25 Met Val Ala Phe Met Ser Tyr Ile Val Pro Pro Pro Pro Thr
   85             90             95
    Ile Phe Lys Val Asp His Pro Phe His Phe Val Leu Lys Thr
      100             105             110
30 Ser Asp Thr Val Leu Phe Asp Gly Arg Val Arg Leu Pro
   115             120             125

```

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 670 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

```

CGAATTGGGT ACCGGGCCCC CCCTCGAGTT TTTT TTTT TTTT TTTT TTTT TTTT 50
TTTTTCACAT TTAACATTTT TATTACATAA ACTACAACAT TATATAGGTG 100
ATTACATTCA TAAAAAGTGA AACTAAAAC AAAACATTTT TCGTCTACAC 150
GATTTATACC ACATACTAAA AAATGAACCT ATTTTAGACC TAATAACTAT 200
45 TAAAAAATAT TGTAGAAAAA TTA CTTTTCATT AATCTGGAGA TTCAGATAGA 250
TCTTGTATTC TTTTATTTAA GAAAATCACA TCATTATCAT TTATGGAAGT 300
CGAACCTCC CATCAAACAA AACAGTATCC GAAGTCTTTA AGACAAAGTG 350
GAAAGGATGA TCAACTTTAA AAATGGTTGG AGGAGGTGGT ACGATATACG 400
ACATAAATGC AACCATAATT ACACCTGTGG CAGCTGCAGC TTCAGCACCT 450
50 TCTTCATTTA CTTCAATGAA AGCTTTTGA ATTACTTTAG AAATATATAA 500
CATCTCATCA GATCCTTCAA GCAATCCTTT GAAATCAGCT TTTCCAGGCA 550
TGAACATATC AGACATACCC AACTTTTCA GAGGATCATT CAAATTAATT 600

```

-146-

TCAGATTCAA TCTTGAATTT AGGCAGATCC AAAATAACTT CAACAGAGTA 650
 CATGCGTTGA GTCAAGTTTT 670

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:
 5 (A) LENGTH: 706 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 10 (A) NAME/KEY: CDS
 (B) LOCATION: 3..410

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

15	TT GAA GAA AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT CAA	44
	Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln	
	1 5 10	
	CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC	86
20	Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe	
	15 20 25	
	AAG ATT GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG	128
	Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys	
	30 35 40	
25	TTG GGT ATG TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC	170
	Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe	
	45 50 55	
30	AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT	212
	Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser	
	60 65 70	
	AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT	254
35	Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly	
	75 80	
	GCT GAA GCT GCA GCT GCC ACA GGA ATC GTT AGT TTT GGC TCA	296
40	Ala Glu Ala Ala Ala Ala Thr Gly Ile Val Ser Phe Gly Ser	
	85 90 95	
	TCT CTG TAT GTC GAC AAT CGT CCT CCA GTT GCT TTT ACC GTA	338
	Ser Leu Tyr Val Asp Asn Arg Pro Pro Val Ala Phe Thr Val	
	100 105 110	
45	GAT CAC CCA TTC TAC TAT ACT TTA AAT ACT TGG GAT ACT CTT	380
	Asp His Pro Phe Tyr Tyr Thr Leu Asn Thr Trp Asp Thr Leu	
	115 120 125	
	TTG TTC AAT GGG CGA GTT ATA TCT CCC AAA TAA AAGGCGTTTA	423
50	Leu Phe Asn Gly Arg Val Ile Ser Pro Lys	
	130 135	
	TTGAGAAGAA TACAAGATCT ATCTGAATCT CTGGATTAAT GAAGTAATTT	473
	TTCTACAATA TTTTAAATA GTTATTAGGT CTAAAAAAG TTCATTTTTT	523
	AGTATGTGGT ATAAATCGTG TAGACGAAAA ATGTTTGTGTT TTAGTTTTC	573

-147-

CTTTTATGA ATGTAATCAC CTATATAATG TTGTAGTTTA TGTAATAAAA	623
ATGTTAAATG TGAAAATATA TTTGATACTA ATAATTAAAA AAAAAAAAAA	673
AAAACTCGAG GGGGGGCCCC GTACCCAATT TCG	706

(2) INFORMATION FOR SEQ ID NO:61:

5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln
1				5					10				
Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp	Leu	Pro	Lys	Phe
15	15			20					25				
Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	Pro	Leu	Lys	Lys
	30				35					40			
Leu	Gly	Met	Ser	Asp	Met	Phe	Val	Pro	Gly	Lys	Ala	Asp	Phe
20		45				50						55	
Lys	Gly	Leu	Leu	Glu	Gly	Ser	Asp	Glu	Met	Leu	Tyr	Ile	Ser
		60			65							70	
Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile	Glu	Val	Asn	Glu	Glu	Gly
			75					80					
Ala	Glu	Ala	Ala	Ala	Ala	Thr	Gly	Ile	Val	Ser	Phe	Gly	Ser
30	85			90					95				
Ser	Leu	Tyr	Val	Asp	Asn	Arg	Pro	Pro	Val	Ala	Phe	Thr	Val
	100				105						110		
Asp	His	Pro	Phe	Tyr	Tyr	Thr	Leu	Asn	Thr	Trp	Asp	Thr	Leu
35		115				120					125		
Leu	Phe	Asn	Gly	Arg	Val	Ile	Ser	Pro	Lys				
		130					135						

(2) INFORMATION FOR SEQ ID NO:62:

40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 706 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

CGAAATTGGG	TACCGGGCCC	CCCCTCGAGT	TTTTTTTTTT	TTTTTTTAAT	50
TATTAGTATC	AAATATATTT	TCACATTTAA	CATTTTTTATT	ACATAAACTA	100
CAACATTATA	TAGGTGATTA	CATTCATAAA	AAGTGAAAAC	TAAAACAAAA	150
CATTTTTCGT	CTACACGATT	TATACCACAT	ACTAAAAAAT	GAAC TTATTT	200
50 TAGACCTAAT	AACTATTAAA	AAATATTGTA	GAAAAATTAC	TTCATTAATC	250

	CAGAGATTCA	GATAGATCTT	GTATTCTTCT	CAATAAACGC	CTTTTATTTG	300
	GGAGATATAA	CTCGCCCAT	GAACAAAAGA	GTATCCCAAG	TATTTAAAGT	350
	ATAGTAGAAT	GGGTGATCTA	CGGTAAAAGC	AAC TGAGAGGA	CGATTGTCGA	400
	CATACAGAGA	TGAGCCAAAA	CTAACGATTC	CTGTGGCAGC	TGCAGCTTCA	450
5	GCACCTTCTT	CATTTACTTC	AATGAAAGCT	TTTTGAATTA	CTTTAGAAAT	500
	ATATAACATC	TCATCAGATC	CTTCAAGCAA	TCCTTTGAAA	TCAGCTTTTC	550
	CAGGAACAAA	CATATCAGAC	ATACCCAAT	TTTTCAGAGG	ATCATTCAAA	600
	TTAATTTTCA	ATTCAATCTT	GAATTTAGGC	AGATCCAAAA	TACATTTCAAC	650
	AGAGTACATG	CGTTGAGTCA	AGTTTTGCAA	GTCAACATTT	TGTAATTTTTT	700
10	CTTCAA					706

(2) INFORMATION FOR SEQ ID NO:63:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 623 nucleotides

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 3..368

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

	AA	AAC	TTG	ACT	CAA	CGC	ATG	TAC	TCT	GTT	GAA	GTT	ATT	TTG	GAT	44
	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp		
	1				5					10						
25	CTG	CCT	AAA	TTC	AAG	ATT	GAA	TCT	GAA	ATT	AAT	TTG	AAT	GAT		86
	Leu	Pro	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp		
	15					20					25					
30	CCT	CTG	AAA	AAG	TTG	GGT	ATG	TCT	GAT	ATG	TTT	GTT	CCT	GGA		128
	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	Met	Phe	Val	Pro	Gly		
	30						35					40				
	AAA	GCT	GAT	TTC	AAA	GGA	TTG	CTT	GAA	GGA	TCT	GAT	GAG	ATG		170
	Lys	Ala	Asp	Phe	Lys	Gly	Leu	Leu	Glu	Gly	Ser	Asp	Glu	Met		
35			45					50					55			
	TTA	TAT	ATT	TCT	AAA	GTA	ATT	CAA	AAA	GCT	TTC	ATT	GAA	GTA		212
	Leu	Tyr	Ile	Ser	Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile	Glu	Val		
				60					65					70		
40	AAT	GAA	GAA	GGT	GCT	GAA	GCT	GCA	GCT	GCC	ACA	GGA	TTA	TTT		254
	Asn	Glu	Glu	Gly	Ala	Glu	Ala	Ala	Ala	Ala	Thr	Gly	Leu	Phe		
					75					80						
45	TTC	TCA	ATA	ACG	TCC	TTC	CAA	GAA	CCG	ACT	TTA	TTC	GAA	GCT		296
	Phe	Ser	Ile	Thr	Ser	Phe	Gln	Glu	Pro	Thr	Leu	Phe	Glu	Ala		
	85					90					95					
	GAC	CGA	CCT	TTT	ATG	TTC	ATC	TTA	CGT	ACT	CAG	GAA	AAT	CCT		338
50	Asp	Arg	Pro	Phe	Met	Phe	Ile	Leu	Arg	Thr	Gln	Glu	Asn	Pro		
		100					105					110				
	ATT	CTA	CTA	TTT	TCC	GGT	CAT	TTT	GTC	GAA	TGA	TGA	ACTTAGA			381
	Ile	Leu	Leu	Phe	Ser	Gly	His	Phe	Val	Glu						
55			115					120								

-149-

	ATACAAGATC TATCTGAATC TCTGGATTAA TGAAGTAATT TTTCTACAAT	431
	ATTTTTTTAAT AGTTATTAGG TCTAAAATAA GTTCATTTTT TAGTATGTGG	481
	TATAAATCGT GTAGACGAAA AATGTTTTGT TTTAGTTTTT ACTTTTATGA	531
	ATGTATCACC TATATAATGT GTAGTTATGT ATAAAATGTT AAATGTGAAA	581
5	AAAAAAAAA AAAAACTCG AGGGGGGGCC GGTACCAATT CG	623

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 122 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

15	Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile Leu Asp	
	1 5 10	
15	Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp	
	15 20 25	
20	Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Val Pro Gly	
	30 35 40	
	Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu Met	
	45 50 55	
25	Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val	
	60 65 70	
	Asn Glu Glu Gly Ala Glu Ala Ala Ala Ala Thr Gly Leu Phe	
	75 80	
30	Phe Ser Ile Thr Ser Phe Gln Glu Pro Thr Leu Phe Glu Ala	
	85 90 95	
	Asp Arg Pro Phe Met Phe Ile Leu Arg Thr Gln Glu Asn Pro	
	100 105 110	
35	Ile Leu Leu Phe Ser Gly His Phe Val Glu	
	115 120	

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 623 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

40	CGAATTGGTA CCGGCCCCC CTCGAGTTT TTTTTTTTT TTTTTCACAT	50
	TTAACATTTT ATACATAACT ACACATTATA TAGGTGATAC ATTCAATAAA	100
	GTGAAACTA AAACAAAACA TTTTTCGTCT ACACGATTTA TACCACATAC	150
	TAAAAAATGA ACTTATTTTA GACCTAATAA CTATTAAAAA ATATTGTAGA	200
50	AAAATTACTT CATTAATCCA GAGATTCAGA TAGATCTTGT ATTCTAAGTT	250
	CATCATTCGA CAAAATGACC GGAAAATAGT AGAATAGGAT TTCCTGAGT	300

-150-

	ACGTAAGATG AACATAAAAG GTCGGTCAGC TTCGAATAAA GTCGGTCTCT	350
	GGAAGGACGT TATTGAGAAA AATAATCCTG TGGCAGCTGC AGCTTCAGCA	400
	CCTTCTTCAT TTACTTCAAT GAAAGCTTTT TGAATTACTT TAGAAATATA	450
	TAACATCTCA TCAGATCCTT CAAGCAATCC TTTGAAATCA GCTTTCCAG	500
5	GAACAAACAT ATCAGACATA CCAACTTTT TCAGAGGATC ATTCAAATTA	550
	ATTTGAGATT CAATCTTGAA TTAGGCAGA TCCAAAATAA CTTCACAGA	600
	GTACATGCGT TGAGTCAAGT TTT	623

(2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 731 nucleotides

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 3..413

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

20	TT GAA GAA AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT CAA	44
	Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln	
	1 5 10	
25	CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC	86
	Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe	
	15 20 25	
30	AAG ATT GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG	128
	Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys	
	30 35 40	
35	TTG GGT ATG TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC	170
	Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe	
	45 50 55	
40	AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT	212
	Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser	
	60 65 70	
45	AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT	254
	Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly	
	75 80	
50	GCT GAA GCT GCA GCT GCC ACA GCC GTG TTT GCG ACT CGT CGT	296
	Ala Glu Ala Ala Ala Ala Thr Ala Val Phe Ala Thr Arg Arg	
	85 90 95	
55	GTG ATC AAG GTG CTG GCG AAA GAA ATT TTC AAT TGC GAC CAT	338
	Val Ile Lys Val Leu Ala Lys Glu Ile Phe Asn Cys Asp His	
	100 105 110	
60	CCG TTC TAC TTC GCC TTG GTT CAT TCG CAA GAA GGT ACC TCG	380
	Pro Phe Tyr Phe Ala Leu Val His Ser Gln Glu Gly Thr Ser	
	115 120 125	

-151-

GCG CCT CTT TTC ACC GGC GCT TTC CGG ACG CCT TGA 416
 Ala Pro Leu Phe Thr Gly Ala Phe Arg Thr Pro
 130 135

5 TAAATGACAG TTCCATTTTC CGCACAATAA GAAAAATCAC GGAAAAGAGA 466
 GAAAGTGGAA AGTAATACAA GATCTATCTG AATCTCTGGA TTAATGAAGT 516
 AATTTTCTA CAATATTTT TAATAGTTAT TAAGTCTAAA ATAAGTTCAA 566
 TTTTAAAGTA TGTGGTATAA ATCGTGTAGA CGAAAAATGT TTTGTTTAA 616
 GTTTCACCTT TAAGAAATGT ATCACCTATA TAATGTTGTA GTTATGTAA 666
 TAAAAATGTT AAATGTGAAA AAAAAAAAAA AAAAAACTCG AGGGGGGGCC 716
 10 CGGTACCCAA TTTCG 731

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 137 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Protein
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln
 1 5 10

20 Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe
 15 20 25

25 Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys
 30 35 40

Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe
 45 50 55

30 Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser
 60 65 70

Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly
 75 80

35 Ala Glu Ala Ala Ala Thr Ala Val Phe Ala Thr Arg Arg
 85 90 95

Val Ile Lys Val Leu Ala Lys Glu Ile Phe Asn Cys Asp His
 100 105 110

40 Pro Phe Tyr Phe Ala Leu Val His Ser Gln Glu Gly Thr Ser
 115 120 125

Ala Pro Leu Phe Thr Gly Ala Phe Arg Thr Pro
 45 130 135

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 731 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA

-152-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

	CGAAATTGGG	TACCGGGCCC	CCCCTCGAGT	TTTTTTTTTT	TTTTTTTTTCA	50
	CATTTAACAT	TTTTATTACA	TAAACTACAA	CATTATATAG	GTGATACATT	100
	TCTTAAAAGT	GAAACTTAAA	ACAAAACATT	TTTCGTCTAC	ACGATTTATA	150
5	CCACATACTT	AAAAATTGAA	CTTATTTTAG	ACTTAATAAC	TATTAAAAAA	200
	TATTGTAGAA	AAATTACTTC	ATTAATCCAG	AGATTCAGAT	AGATCTTGTA	250
	TTACTTTCCA	CTTCTCTCT	TTTCCGTGAT	TTTTCTTATT	GTGCGGAAAA	300
	TGGAACGTGC	ATTTATCAAG	GCGTCCGGAA	AGCGCCGGTG	AAAAGAGGCG	350
	CCGAGGTACC	TTCTTGCGAA	TGAACCAAGG	CGAAGTAGAA	CGGATGGTCG	400
10	CAATTGAAAA	TTTCTTTCGC	CAGCACCTTG	ATCACACGAC	GAGTCGCAAA	450
	CACGGCTGTG	GCAGCTGCAG	CTTCAGCACC	TTCTTCATTT	ACTTCAATGA	500
	AAGCTTTTTG	AATTACTTTA	GAAATATATA	ACATCTCATC	AGATCCTTCA	550
	AGCAATCCTT	TGAAATCAGC	TTTTCAGAGG	ACAAACATAT	CAGACATACC	600
	CAACTTTTTT	AGAGGATCAT	TCAAATTAAT	TTCAGATTCA	ATCTTGAATT	650
15	TAGGCAGATC	CAAAATAACT	TCAACAGAGT	ACATGCGTTG	AGTCAAGTTT	700
	TGCAAGTCAA	CATTTTGTAA	TTTTTCTTCA	A		731

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

	(A) LENGTH:	685 nucleotides
20	(B) TYPE:	nucleic acid
	(C) STRANDEDNESS:	single
	(D) TOPOLOGY:	linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

25	(A) NAME/KEY:	CDS
	(B) LOCATION:	3..407

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

	TT GAA GAA AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT CAA	44
	Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln	
30	1 5 10	
	CGC ATG TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC	86
	Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe	
	15 20 25	
35	AAG ATT GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG	128
	Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys	
	30 35 40	
	TTG GGT ATG TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC	170
40	Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe	
	45 50 55	
	AAA GGA TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT	212
	Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser	
45	60 65 70	
	AAA GTA ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT	254
	Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly	
	75 80	
50	GCT GAA GCT GCA GCT GCC ACA GCT GTC GTG ATG CTT GGA TAT	296
	Ala Glu Ala Ala Ala Ala Thr Ala Val Val Met Leu Gly Tyr	
	85 90 95	

-153-

```

TCC CTA ATT ACG TCT CGG GTA GCT CCA ACT GTT TTT AAC GTC      338
Ser Leu Ile Thr Ser Arg Val Ala Pro Thr Val Phe Asn Val
  100                      105                      110

5  GAT CAT CCA TTC CAT GTT GTA TTA AAA TCA AAT GAT GTT GTT      380
   Asp His Pro Phe His Val Val Leu Lys Ser Asn Asp Val Val
      115                      120                      125

TTA TTT AAT GGA CGC GTT CAG TCA CCA TGA AATGGATATT          420
Leu Phe Asn Gly Arg Val Gln Ser Pro
10                      130                      135

TTTGGTAAAA GAATACAAGA TCTATCTGAA TCTCTGGATT AATGAAGTAA      470
TTTTTCTACA ATATTTTTTA ATAGTTATTA GGTCTAAAAT AAGTTCATTT      520
TTTAGTATGT GGTATAAATC GTGTAGACGA AAAATGTTTT GTTTTAGTTT      570
TCACTTTTTA TGAATGTAAT CACCTATATA ATGTTGTAGT TTATGTAATA      620
15 AAAATGTTAA ATGTGAAAAA AAAAAAAAAA AAAAAAATC GAGGGGGGGC      670
   CCGGTACCCA ATTCG                                     685

```

(2) INFORMATION FOR SEQ ID NO:70:

```

(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 135 amino acids
    (B) TYPE: amino acid
    (D) TOPOLOGY: linear
20

(ii) MOLECULE TYPE: Protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

25  Glu Glu Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln
    1                      5                      10

    Arg Met Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe
    15                      20                      25

    Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys
30    30                      35                      40

    Leu Gly Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe
    45                      50                      55

35  Lys Gly Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser
    60                      65                      70

    Lys Val Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly
    75                      80

40  Ala Glu Ala Ala Ala Ala Thr Ala Val Val Met Leu Gly Tyr
    85                      90                      95

    Ser Leu Ile Thr Ser Arg Val Ala Pro Thr Val Phe Asn Val
45    100                      105                      110

    Asp His Pro Phe His Val Val Leu Lys Ser Asn Asp Val Val
    115                      120                      125

50  Leu Phe Asn Gly Arg Val Gln Ser Pro
    130                      135

```

-154-

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 685 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

	CGAATTGGGT ACCGGGCCCC CCCTCGAGTT TTTTTTTTTT TTTTTTTTTT	50
10	CACATTTAAC ATTTTATTAT CATAAACTAC AACATTATAT AGGTGATTAC	100
	ATTCATAAAA AGTGAAACT AAAACAAAAC ATTTTTCGTC TACACGATTT	150
	ATACCACATA CTAAAAAATG AACTTATTTT AGACCTAATA ACTATTAAAA	200
	AATATTGTAG AAAAATTACT TCATTAAATCC AGAGATTTCAG ATAGATCTTG	250
	TATTCTTTTA CCAAAAATAT CCATTTTCATG GTGACTGAAC GCGTCCATTA	300
15	AATAAAACAA CATCATTTGA TTTTAATACA ACATGGAATG GATGATCGAC	350
	GTTAAAAACA GTTGGAGCTA CCCGAGACGT AATTAGGGAA TATCCAAGCA	400
	TCACGACAGC TGTGGCAGCT GCAGCTTCAG CACCTTCTTC ATTTACTTCA	450
	ATGAAAGCTT TTTGAATTAC TTTAGAAATA TATAACATCT CATCAGATCC	500
	TTCAAGCAAT CCTTTGAAAT CAGCTTTTCC AGGAACAAAC ATATCAGACA	550
20	TACCCAACCTT TTTCAGAGGA TCATTCAAAT TAATTTTCAGA TTCAATCTTG	600
	AATTTAGGCA GATCCAAAAT AACTTCAACA GAGTACATGC GTTGAGTCAA	650
	GTTTTGCAAG TCAACATTTT GTAATTTTTC TTCAA	685

(2) INFORMATION FOR SEQ ID NO:72:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1222 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 3.1220

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

35	AC GCG ATA GTT CAA CAC GCA CGA CTT GTG TTT CTT TTT GTA TCA	44
	Ala Ile Val Gln His Ala Arg Leu Val Phe Leu Phe Val Ser	
	1 5 10	
40	GTG TTA ATA CCA ATT TCA ACA ATG GCG GAT CCC CAG GAA TTG	86
	Val Leu Ile Pro Ile Ser Thr Met Ala Asp Pro Gln Glu Leu	
	15 20 25	
45	TCT ACA AGT ATT AAC CAG TTT GCT GGA AGC CTG TAC AAT ACG	128
	Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr	
	30 35 40	
50	GTT GCT TCT GGC AAC AAA GAC AAT CTC ATC ATG TCC CCA TTG	170
	Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser Pro Leu	
	45 50 55	
50	TCT GTA CAA ACT GTT CTA TCC CTG GTG TCA ATG GGA GCT GGT	212
	Ser Val Gln Thr Val Leu Ser Leu Val Ser Met Gly Ala Gly	
	60 65 70	

-155-

	GGT AAT ACT GCC ACA CAA ATA GCT GCT GGT TTA CGT CAG CCT	254
	Gly Asn Thr Ala Thr Gln Ile Ala Ala Gly Leu Arg Gln Pro	
	75 80	
5	CAA TCA AAA GAA AAA ATT CAA GAT GAC TAC CAT GCA TTG ATG	296
	Gln Ser Lys Glu Lys Ile Gln Asp Asp Tyr His Ala Leu Met	
	85 90 95	
10	AAC ACT CTT AAT ACA CAA AAA GGT GTA ACT CTG GAA ATT GCC	338
	Asn Thr Leu Asn Thr Gln Lys Gly Val Thr Leu Glu Ile Ala	
	100 105 110	
15	AAC AAA GTT TAC GTT ATG GAA GGC TAT ACA TTG AAA CCC ACC	380
	Asn Lys Val Tyr Val Met Glu Gly Tyr Thr Leu Lys Pro Thr	
	115 120 125	
20	TTC AAA GAA GTT GCC ACC AAC AAA TTC TTA GCT GGA GCA GAA	422
	Phe Lys Glu Val Ala Thr Asn Lys Phe Leu Ala Gly Ala Glu	
	130 135 140	
	AAC TTG AAC TTT GCC CAA AAT GCT GAA AGC GCT AAA GTT ATC	464
	Asn Leu Asn Phe Ala Gln Asn Ala Glu Ser Ala Lys Val Ile	
	145 150	
25	AAC ACT TGG GTT GAA GAA AAA ACT CAT GAC AAA ATT CAT GAT	506
	Asn Thr Trp Val Glu Glu Lys Thr His Asp Lys Ile His Asp	
	155 160 165	
30	TTG ATC AAA GCC GGT GAT CTA GAC CAG GAT TCA AGA ATG GTT	548
	Leu Ile Lys Ala Gly Asp Leu Asp Gln Asp Ser Arg Met Val	
	170 175 180	
35	CTT GTC AAT GCA TTG TAC TTC AAG GGT CTT TGG GAG AAA CAA	590
	Leu Val Asn Ala Leu Tyr Phe Lys Gly Leu Trp Glu Lys Gln	
	185 190 195	
40	TTC AAG AAG GAA AAC ACT CAA GAC AAA CCT TTC TAT GTT ACT	632
	Phe Lys Lys Glu Asn Thr Gln Asp Lys Pro Phe Tyr Val Thr	
	200 205 210	
	GAA ACA GAG ACA AAG AAT GTA CGA ATG ATG CAC ATT AAG GAT	674
	Glu Thr Glu Thr Lys Asn Val Arg Met Met His Ile Lys Asp	
	215 220	
45	AAA TTC CGT TAT GGA GAA TTT GAA GAA TTA GAT GCC AAG GCT	716
	Lys Phe Arg Tyr Gly Glu Phe Glu Glu Leu Asp Ala Lys Ala	
	225 230 235	
50	GTA GAA TTG CCC TAC AGG AAC TCA GAT TTG GCC ATG TTA ATC	758
	Val Glu Leu Pro Tyr Arg Asn Ser Asp Leu Ala Met Leu Ile	
	240 245 250	
55	ATT TTG CCA AAC AGC AAA ACT GGT CTC CCC GCT CTT GAA GAA	800
	Ile Leu Pro Asn Ser Lys Thr Gly Leu Pro Ala Leu Glu Glu	
	255 260 265	
60	AAA TTA CAA AAT GTT GAC TTG CAA AAC TTG ACT CAA CGC ATG	842
	Lys Leu Gln Asn Val Asp Leu Gln Asn Leu Thr Gln Arg Met	
	270 275 280	

-156-

	TAC TCT GTT GAA GTT ATT TTG GAT CTG CCT AAA TTC AAG ATT	884
	Tyr Ser Val Glu Val Ile Leu Asp Leu Pro Lys Phe Lys Ile	
	285 290	
5	GAA TCT GAA ATT AAT TTG AAT GAT CCT CTG AAA AAG TTG GGT	926
	Glu Ser Glu Ile Asn Leu Asn Asp Pro Leu Lys Lys Leu Gly	
	295 300 305	
10	ATG TCT GAT ATG TTT GTT CCT GGA AAA GCT GAT TTC AAA GGA	968
	Met Ser Asp Met Phe Val Pro Gly Lys Ala Asp Phe Lys Gly	
	310 315 320	
15	TTG CTT GAA GGA TCT GAT GAG ATG TTA TAT ATT TCT AAA GTA	1010
	Leu Leu Glu Gly Ser Asp Glu Met Leu Tyr Ile Ser Lys Val	
	325 330 335	
20	ATT CAA AAA GCT TTC ATT GAA GTA AAT GAA GAA GGT GCT GAA	1052
	Ile Gln Lys Ala Phe Ile Glu Val Asn Glu Glu Gly Ala Glu	
	340 345 350	
25	GCT GCA GCT GCC ACA GCG GTG CTT TTA GTA ACG GAA TCT TAT	1094
	Ala Ala Ala Ala Thr Ala Val Leu Leu Val Thr Glu Ser Tyr	
	355 360	
30	GTA CCT GAG GAA GTA TTC GAA GCT AAT CAT CCC TTT TAT TTT	1136
	Val Pro Glu Glu Val Phe Glu Ala Asn His Pro Phe Tyr Phe	
	365 370 375	
35	GCA CTC TAT AAA TCT GCA CAA AAT CCA GTA GAA TCT GAA AAT	1178
	Ala Leu Tyr Lys Ser Ala Gln Asn Pro Val Glu Ser Glu Asn	
	380 385 390	
40	GAA AGC TCT GAA AAT GAA AAC CCT GAA AAT GTT GAA GTA CTA	1220
	Glu Ser Ser Glu Asn Glu Asn Pro Glu Asn Val Glu Val Leu	
	395 400 405	
	TT	1222

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 34 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

45	GGAAGATCTA TAAATATGCC GCGTCCTCAG TTTG	34
----	---------------------------------------	----

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 29 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer

-157-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

CGGAATTCTA ATTGGTAAAT CTCCCAGAG

29

(2) INFORMATION FOR SEQ ID NO:75:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1155 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- 10 (A) NAME/KEY: CDs
 (B) LOCATION: 1..1155

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

15	GTG TTT CTT TTT GTA TCA GTG TTA TTA CCA ATT TCA ACA ATG	42
	Val Phe Leu Phe Val Ser Val Leu Leu Pro Ile Ser Thr Met	
	1 5 10	
20	GCC GAT CCC CAG GAA TTG TCT ACA AGT ATT AAC CAG TTT GCT	84
	Ala Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala	
	15 20 25	
25	GGA AGC CTG TAC AAT ACA GTT GCT TCT GGC AAC AAA GAC AAT	126
	Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn	
	30 35 40	
30	CTC ATC ATG TCC CCA TTG TCT GTA CAA ACT GTT CTA TCC CTG	168
	Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu Ser Leu	
	45 50 55	
35	GTG TCA ATG GGA GCT GGT GGC AAT ACT GCC ACA CAA ATA GCT	210
	Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln Ile Ala	
	60 65 70	
40	GCT GGT TTG CGT CAG CCT CAA TCA AAA GAA AAA ATT CAA GAT	252
	Ala Gly Leu Arg Gln Pro Gln Ser Lys Glu Lys Ile Gln Asp	
	75 80	
45	GAC TAC CAC GCA TTG ATG AAC ACT CTT AAT ACA CAA AAA GGT	294
	Asp Tyr His Ala Leu Met Asn Thr Leu Asn Thr Gln Lys Gly	
	85 90 95	
50	GTA ACT CTG GAA ATT GCC AAT AAA GTT TAT GTT ATG GAA GGC	336
	Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val Met Glu Gly	
	100 105 110	
55	TAT ACA TTA AAA CCC ACC TTC AAA GAA GTT GCC ACC AAC AAA	378
	Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr Asn Lys	
	115 120 125	
60	TTC TTA GCT GGA GCA GAA AAC TTG AAC TTT GCC CAA AAT GCT	420
	Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln Asn Ala	
	130 135 140	

-158-

	GAA	AGC	GCT	AAA	GTT	ATC	AAC	ACT	TGG	GTT	GAA	GAA	AAA	ACT	462
	Glu	Ser	Ala	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	
					145					150					
5	CAT	GAC	AAA	ATT	CAT	GAT	TTG	ATC	AAA	GCC	GGT	GAT	CTA	GAC	504
	His	Asp	Lys	Ile	His	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	
	155					160					165				
10	CAG	GAT	TCA	AGA	ATG	GTT	CTT	GTC	AAT	GCA	TTG	TAC	TTC	AAG	546
	Gln	Asp	Ser	Arg	Met	Val	Leu	Val	Asn	Ala	Leu	Tyr	Phe	Lys	
		170					175					180			
15	GGT	CTT	TGG	GAG	AAA	CAA	TTC	AAA	AAG	GAA	AAT	ACC	CAA	GAC	588
	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	Glu	Asn	Thr	Gln	Asp	
			185					190					195		
20	AAA	CCT	TTC	TAT	GTT	ACT	GAA	ACA	GAG	ACA	AAG	AAT	GTA	CGA	630
	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg	
				200					205					210	
25	ATG	ATG	CAC	ATT	AAG	GAT	AAA	TTC	CGT	TAT	GGA	GAA	TTT	GAA	672
	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu	
					215					220					
30	GAA	TTA	GAT	GCC	AAG	GCT	GTA	GAA	TTG	CCC	TAC	AGG	AAC	TCA	714
	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	
		225				230					235				
35	GAT	TTG	GCC	ATG	TTA	ATC	ATT	TTG	CCA	AAC	AGC	AAA	ACT	GGT	756
	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	
		240					245					250			
40	CTC	CCC	GCT	CTT	GAA	GAA	AAA	TTA	CAA	AAT	GTT	GAT	TTG	CAA	798
	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	
			255					260					265		
45	AAC	TTG	ACT	CAA	CGC	ATG	TAC	TCT	GTT	GAA	GTT	ATT	TTG	GAT	840
	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp	
				270				275						280	
50	CTG	CCT	AAA	TTC	AAG	ATT	GAA	TCT	GAA	ATT	AAT	TTG	AAT	GAT	882
	Leu	Pro	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	
					285					290					
55	CCT	CTG	AAA	AAG	TTG	GGT	ATG	TCT	GAT	ATG	TTT	GTT	CCT	GGA	924
	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	Met	Phe	Val	Pro	Gly	
		295				300					305				
60	AAA	GCT	GAT	TTC	AAA	GGA	TTG	CTT	GAA	GGA	TCT	GAT	GAG	ATG	966
	Lys	Ala	Asp	Phe	Lys	Gly	Leu	Leu	Glu	Gly	Ser	Asp	Glu	Met	
		310					315					320			
65	TTA	TAT	ATT	TCT	AAA	GTA	ATT	CAA	AAA	GCT	TTC	ATT	GAA	GTA	1008
	Leu	Tyr	Ile	Ser	Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile	Glu	Val	
			325					330					335		
70	AAT	GAA	GAA	GGT	GCT	GAA	GCT	GCA	GCT	GCC	ACA	GCT	ACC	TTT	1050
	Asn	Glu	Glu	Gly	Ala	Glu	Ala	Ala	Ala	Ala	Thr	Ala	Thr	Phe	
				340					345					350	

-159-

ATG GTT ACC TAT GAA CTG GAG GTT TCC CTG GAT CTT CCC ACT 1092
 Met Val Thr Tyr Glu Leu Glu Val Ser Leu Asp Leu Pro Thr
 355 360

5 GTT TTT AAA GTC GAT CAT CCA TTC AAT ATT GTT TTG AAG ACA 1134
 Val Phe Lys Val Asp His Pro Phe Asn Ile Val Leu Lys Thr
 365 370 375

GGT GAT ACT GTT ATT TTT AAT 1155
 Gly Asp Thr Val Ile Phe Asn
 10 380 385

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 nucleotides
 (B) TYPE: nucleic acid
 15 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

GGAAGATCTA TAAATATGAT TAACGCACGA CTT 33

20 (2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 nucleotides
 (B) TYPE: nucleic acid
 25 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

CCGGAATTCA TAGAGTTTGA ACTCGCCC 28

(2) INFORMATION FOR SEQ ID NO:78:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1065 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 3..1064

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

40 AG TTT GCT GGA AGC CTG TAC AAT ACG GTT GCT TCT GGC AAC AAA 44
 Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys
 1 5 10

		GAC Asp 15	AAT Asn	CTC Leu	ATC Ile	ATG Met	TCC Ser 20	CCA Pro	TTG Leu	TCT Ser	GTA Val	CAA Gln 25	ACT Thr	GTT Val	CTA Leu	86
5		TCC Ser 30	CTG Leu	GTG Val	TCA Ser	ATG Met	GGA Gly	GCT Ala 35	GGT Gly	GGT Gly	AAT Asn	ACT Thr	GCC Ala 40	ACA Thr	CAA Gln	128
10		ATA Ile	GCT Ala	GCT Ala 45	GGT Gly	TTA Leu	CGT Arg	CAG Gln	CCT Pro 50	CAA Gln	TCA Ser	AAA Lys	GAA Glu	AAA Lys 55	ATT Ile	170
		CAA Gln	GAT Asp	GAC Asp	TAC Tyr 60	CAT His	GCA Ala	TTG Leu	ATG Met	AAC Asn 65	ACT Thr	CTT Leu	AAT Asn	ACA Thr	CAA Gln 70	212
15		AAA Lys	GGT Gly	GTA Val	ACT Thr 75	CTG Leu	GAA Glu	ATT Ile	GCC Ala	AAC Asn 80	AAA Lys	GTT Val	TAC Tyr	GTT Val	ATG Met	254
20		GAA Glu 85	GGC Gly	TAT Tyr	ACA Thr	TTG Leu	AAA Lys 90	CCC Pro	ACC Thr	TTC Phe	AAA Lys	GAA Glu 95	GTT Val	GCC Ala	ACC Thr	296
25		AAC Asn 100	AAA Lys	TTC Phe	TTA Leu	GCT Ala	GGA Gly	GCA Ala 105	GAA Glu	AAC Asn	TTG Leu	AAC Asn 110	TTT Phe	GCC Ala	CAA Gln	338
30		AAT Asn	GCT Ala 115	GAA Glu	AGC Ser	GCT Ala	AAA Lys	GTT Val	ATC Ile 120	AAC Asn	ACT Thr	TGG Trp	GTT Val 125	GAA Glu	GAA Glu	380
		AAA Lys	ACT Thr	CAT His	GAC Asp 130	AAA Lys	ATT Ile	CAT His	GAT Asp 135	TTG Leu	ATC Ile	AAA Lys	GCC Ala	GGT Gly	GAT Asp 140	422
35		CTA Leu	GAC Asp	CAG Gln	GAT Asp	TCA Ser 145	AGA Arg	ATG Met	GTT Val	CTT Leu	GTC Val 150	AAT Asn	GCA Ala	TTG Leu	TAC Tyr	464
40		TTC Phe 155	AAG Lys	GGT Gly	CTT Leu	TGG Trp	GAG Glu 160	AAA Lys	CAA Gln	TTC Phe	AAG Lys	AAG Lys 165	GAA Glu	AAC Asn	ACT Thr	506
45		CAA Gln 170	GAC Asp	AAA Lys	CCT Pro	TTC Phe	TAT Tyr	GTT Val 175	ACT Thr	GAA Glu	ACA Thr	GAG Glu 180	ACA Thr	AAG Lys	AAT Asn	548
50		GTA Val	CGA Arg	ATG Met 185	ATG Met	CAC His	ATT Ile	AAG Lys	GAT Asp 190	AAA Lys	TTC Phe	CGT Arg	TAT Tyr	GGA Gly	GAA Glu	590
		TTT Phe	GAA Glu	GAA Glu 200	TTA Leu	GAT Asp	GCC Ala	AAG Lys	GCT Ala 205	GTA Val	GAA Glu	TTG Leu	CCC Pro	TAC Tyr	AGG Arg 210	632
55		AAC Asn	TCA Ser	GAT Asp	TTG Leu	GCC Ala 215	ATG Met	TTA Leu	ATC Ile	ATT Ile 220	TTG Leu	CCA Pro	AAC Asn	AGC Ser	AAA Lys	674

-161-

	ACT GGT CTC CCC GCT CTT GAA GAA AAA TTA CAA AAT GTT GAC	716
	Thr Gly Leu Pro Ala Leu Glu Glu Lys Leu Gln Asn Val Asp	
	225 230 235	
5	TTG CAA AAC TTG ACT CAA CGC ATG TAC TCT GTT GAA GTT ATT	758
	Leu Gln Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile	
	240 245 250	
10	TTG GAT CTG CCT AAA TTC AAG ATT GAA TCT GAA ATT AAT TTG	800
	Leu Asp Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu	
	255 260 265	
15	AAT GAT CCT CTG AAA AAG TTG GGT ATG TCT GAT ATG TTT GTT	842
	Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Val	
	270 275 280	
20	CCT GGA AAA GCT GAT TTC AAA GGA TTG CTT GAA GGA TCT GAT	884
	Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp	
	285 290	
25	GAG ATG TTA TAT ATT TCT AAA GTA ATT CAA AAA GCT TTC ATT	926
	Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile	
	295 300 305	
30	GAA GTA AAT GAA GAA GGT GCT GAA GCT GCA GCT GCC ACA GGC	968
	Glu Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Gly	
	310 315 320	
35	ATT GTC ATG CTT GGT TGC TGT ATG CCA ATG ATG GAT CTT TCT	1010
	Ile Val Met Leu Gly Cys Cys Met Pro Met Met Asp Leu Ser	
	325 330 335	
40	CCA GTA GTT TTT AAT ATT GAT CAC CCA TTT TAT TAC TCA TTG	1052
	Pro Val Val Phe Asn Ile Asp His Pro Phe Tyr Tyr Ser Leu	
	340 345 350	
45	ATG ACT TGG GAT A	1065
	Met Thr Trp Asp	

(2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 40 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

GCGGAATTTCG ATCCCCAGGA ATTGTCTACA AGTATTAACC 40

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 44 nucleotides
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Primer

-162-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

GCGAGATCTT TAAAGGGATT TAACACATCC ACTGAACAAA ACAG 44

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1070 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- 10 (A) NAME/KEY: CDS
 (B) LOCATION: 3..1070

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

15	AG TTT GCT GGA AGC CTG TAC AAT ACG GTT GCT TCT GGC AAC AAA	44
	Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys	
	1 5 10	
	GAC AAT CTC ATC ATG TCC CCA TTG TCT GTA CAA ACT GTT CTA	86
	Asp Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu	
	15 20 25	
20	TCC CTG GTG TCA ATG GGA GCT GGT GGT AAT ACT GCC ACA CAA	128
	Ser Leu Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln	
	30 35 40	
25	ATA GCT GCT GGT TTA CGT CAG CCT CAA TCA AAA GAA AAA ATT	170
	Ile Ala Ala Gly Leu Arg Gln Pro Gln Ser Lys Glu Lys Ile	
	45 50 55	
30	CAA GAT GAC TAC CAT GCA TTG ATG AAC ACT CTT AAT ACA CAA	212
	Gln Asp Asp Tyr His Ala Leu Met Asn Thr Leu Asn Thr Gln	
	60 65 70	
35	AAA GGT GTA ACT CTG GAA ATT GCC AAC AAA GTT TAC GTT ATG	254
	Lys Gly Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val Met	
	75 80	
	GAA GGC TAT ACA TTG AAA CCC ACC TTC AAA GAA GTT GCC ACC	296
	Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr	
	85 90 95	
40	AAC AAA TTC TTA GCT GGA GCA GAA AAC TTG AAC TTT GCC CAA	338
	Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln	
	100 105 110	
45	AAT GCT GAA AGC GCT AAA GTT ATC AAC ACT TGG GTT GAA GAA	380
	Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu	
	115 120 125	
50	AAA ACT CAT GAC AAA ATT CAT GAT TTG ATC AAA GCC GGT GAT	422
	Lys Thr His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp	
	130 135 140	

-163-

	CTA Leu	GAC Asp	CAG Gln	GAT Asp	TCA Ser 145	AGA Arg	ATG Met	GTT Val	CTT Leu	GTC Val 150	AAT Asn	GCA Ala	TTG Leu	TAC Tyr	464
5	TTC Phe 155	AAG Lys	GGT Gly	CTT Leu	TGG Trp 160	GAG Glu	AAA Lys	CAA Gln	TTC Phe	AAG Lys 165	AAG Lys	GAA Glu	AAC Asn	ACT Thr	506
10	CAA Gln 170	GAC Asp	AAA Lys	CCT Pro	TTC Phe	TAT Tyr	GTT Val 175	ACT Thr	GAA Glu	ACA Thr	GAG Glu	ACA Thr 180	AAG Lys	AAT Asn	548
15	GTA Val	CGA Arg	ATG Met 185	ATG Met	CAC His	ATT Ile	AAG Lys 190	GAT Asp	AAA Lys	TTC Phe	CGT Arg	TAT Tyr	GGA Gly 195	GAA Glu	590
20	TTT Phe	GAA Glu	GAA Glu	TTA Leu 200	GAT Asp	GCC Ala	AAG Lys	GCT Ala 205	GTA Val 205	GAA Glu	TTG Leu	CCC Pro	TAC Tyr	AGG Arg 210	632
	AAC Asn	TCA Ser	GAT Asp	TTG Leu 215	GCC Ala	ATG Met	TTA Leu	ATC Ile	ATT Ile 220	TTG Leu 220	CCA Pro	AAC Asn	AGC Ser	AAA Lys	674
25	ACT Thr 225	GGT Gly	CTC Leu	CCC Pro	GCT Ala	CTT Leu 230	GAA Glu	GAA Glu	AAA Lys	TTA Leu 235	CAA Gln	AAT Asn	GTT Val	GAC Asp	716
30	TTG Leu 240	CAA Gln	AAC Asn	TTG Leu	ACT Thr	CAA Gln 245	CGC Arg	ATG Met	TAC Tyr	TCT Ser	GTT Val 250	GAA Glu	GTT Val	ATT Ile	758
35	TTG Leu	GAT Asp	CTG Leu 255	CCT Pro	AAA Lys	TTC Phe	AAG Lys	ATT Ile 260	GAA Glu	TCT Ser	GAA Glu	ATT Ile 265	AAT Asn	TTG Leu	800
40	AAT Asn	GAT Asp	CCT Pro 270	CTG Leu	AAA Lys	AAG Lys	TTG Leu	GGT Gly 275	ATG Met 275	TCT Ser	GAT Asp	ATG Met	TTT Phe 280	GTT Val	842
	CCT Pro	GGA Gly	AAA Lys	GCT Ala	GAT Asp 285	TTC Phe	AAA Lys	GGA Gly	TTG Leu 290	CTT Leu 290	GAA Glu	GGA Gly	TCT Ser	GAT Asp	884
45	GAG Glu 295	ATG Met	TTA Leu	TAT Tyr	ATT Ile	TCT Ser 300	AAA Lys	GTA Val	ATT Ile	CAA Gln 305	AAA Lys	GCT Ala	TTC Phe	ATT Ile	926
50	GAA Glu 310	GTA Val	AAT Asn	GAA Glu	GAA Glu	GGT Gly 315	GCT Ala 315	GAA Glu	GCT Ala	GCA Ala	GCT Ala 320	GCC Ala	ACA Thr	GGC Gly	968
55	GTG Val	ATG Met	TTA Leu 325	ATG Met	ATG Met	CGT Arg	TGT Cys 330	ATG Met	CCA Pro	ATG Met	ATG Met	CCA Pro 335	ATG Met	GCC Ala	1010
60	TTC Phe	AAT Asn	GCT Ala 340	GAG Glu	CAT His	CCA Pro	TTC Phe	CTG Leu 345	TAC Tyr 345	TTC Phe	TTA Leu	CAC His	AGC Ser	AAA Lys 350	1052

-164-

AAT TCT GTT CTA TTC AAT
Asn Ser Val Leu Phe Asn
355

1070

(2) INFORMATION FOR SEQ ID NO:82:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 10 (ii) MOLECULE TYPE: Primer

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

CGCAGATCTT TATTCAGTTG TTGGTTTAAC AAGACGACC

39

(2) INFORMATION FOR SEQ ID NO:83:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 17 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: Primer

- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

ATTAACCCTC ACTAAAG

17

(2) INFORMATION FOR SEQ ID NO:84:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: Primer

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

30 ATAGGATCCC CAGGAATTGT C

21

(2) INFORMATION FOR SEQ ID NO:85:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: Primer

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

GCGAGATCTC TAGTTATTAA TATTGGTTAA

30

-165-

(2) INFORMATION FOR SEQ ID NO:86:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

GCGGAATTCT CATGGTGACT GAACGCG

27

10 (2) INFORMATION FOR SEQ ID NO:87:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

GCGGAATTCA ACAAAGTGT GTTC

24

(2) INFORMATION FOR SEQ ID NO:88:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

Asp Pro Gln Glu Leu Ser Thr Ser Ile Asn Gln Phe Ala Gly
 1 5 10

Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys Asp Asn Leu
 15 20 25

30 Ile Met
 30

(2) INFORMATION FOR SEQ ID NO:89:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

40 Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr
 1 5 10

-166-

Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met
15 20 25

(2) INFORMATION FOR SEQ ID NO:90:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

10 Ser Thr Ser Ile Asn Gln Phe Ala Gly Ser Leu Tyr Asn Thr
1 5 10

Val Ala Ser Gly Asn Lys Asp Asn Leu Ile Met Ser Pro
15 20 25

(2) INFORMATION FOR SEQ ID NO:91:

- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

GCGGAATTCT TATTTGGGAG ATATAACTCG

30

(2) INFORMATION FOR SEQ ID NO:92:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

CGCGAATTCT CATTCGACAA AATGACC

27

(2) INFORMATION FOR SEQ ID NO:93:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 nucleotides
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

40 GCGGAATTCT TAAGGATTAA CGTGTGTAAC

30

-167-

(2) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 nucleotides
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Primer

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

GGAATTCTTA TTGCACAAAT CATCC

25

10 (2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 406 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

Ala	Ile	Val	Gln	His	Ala	Arg	Leu	Val	Phe	Leu	Phe	Val	Ser	
1				5					10					
Val	Leu	Ile	Pro	Ile	Ser	Thr	Met	Ala	Asp	Pro	Gln	Glu	Leu	
15				20					25					
Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala	Gly	Ser	Leu	Tyr	Asn	Thr	
	30				35					40				
Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Leu	Ile	Met	Ser	Pro	Leu	
	45				50					55				
Ser	Val	Gln	Thr	Val	Leu	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	
30			60				65					70		
Gly	Asn	Thr	Ala	Thr	Gln	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	
			75				80							
Gln	Ser	Lys	Glu	Lys	Ile	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	
35			85		90				95					
Asn	Thr	Leu	Asn	Thr	Gln	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	
	100				105					110				
Asn	Lys	Val	Tyr	Val	Met	Glu	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	
	115				120					125				
Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys	Phe	Leu	Ala	Gly	Ala	Glu	
45			130				135					140		
Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala	Glu	Ser	Ala	Lys	Val	Ile	
			145				150							
Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr	His	Asp	Lys	Ile	His	Asp	
50			155		160				165					

-168-

	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val
	170						175					180		
5	Leu	Val	Asn	Ala	Leu	Tyr	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln
			185					190					195	
	Phe	Lys	Lys	Glu	Asn	Thr	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr
				200					205					210
10	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg	Met	Met	His	Ile	Lys	Asp
					215					220				
	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu	Glu	Leu	Asp	Ala	Lys	Ala
	225					230					235			
15	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser	Asp	Leu	Ala	Met	Leu	Ile
	240						245					250		
	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly	Leu	Pro	Ala	Leu	Glu	Glu
20			255					260					265	
	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln	Asn	Leu	Thr	Gln	Arg	Met
				270					275					280
25	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp	Leu	Pro	Lys	Phe	Lys	Ile
					285					290				
	Glu	Ser	Glu	Ile	Asn	Leu	Asn	Asp	Pro	Leu	Lys	Lys	Leu	Gly
	295					300					305			
30	Met	Ser	Asp	Met	Phe	Val	Pro	Gly	Lys	Ala	Asp	Phe	Lys	Gly
		310					315					320		
	Leu	Leu	Glu	Gly	Ser	Asp	Glu	Met	Leu	Tyr	Ile	Ser	Lys	Val
35			325					330					335	
	Ile	Gln	Lys	Ala	Phe	Ile	Glu	Val	Asn	Glu	Glu	Gly	Ala	Glu
				340					345					350
40	Ala	Ala	Ala	Ala	Thr	Ala	Val	Leu	Leu	Val	Thr	Glu	Ser	Tyr
					355					360				
	Val	Pro	Glu	Glu	Val	Phe	Glu	Ala	Asn	His	Pro	Phe	Tyr	Phe
	365					370					375			
45	Ala	Leu	Tyr	Lys	Ser	Ala	Gln	Asn	Pro	Val	Glu	Ser	Glu	Asn
		380					385					390		
	Glu	Ser	Ser	Glu	Asn	Glu	Asn	Pro	Glu	Asn	Val	Glu	Val	Leu
50			395					400					405	

(2) INFORMATION FOR SEQ ID NO:96:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 385 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: Protein

-169-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

	Val	Phe	Leu	Phe	Val	Ser	Val	Leu	Leu	Pro	Ile	Ser	Thr	Met
	1				5					10				
5	Ala	Asp	Pro	Gln	Glu	Leu	Ser	Thr	Ser	Ile	Asn	Gln	Phe	Ala
	15					20					25			
	Gly	Ser	Leu	Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys	Asp	Asn
		30					35					40		
10	Leu	Ile	Met	Ser	Pro	Leu	Ser	Val	Gln	Thr	Val	Leu	Ser	Leu
			45					50					55	
	Val	Ser	Met	Gly	Ala	Gly	Gly	Asn	Thr	Ala	Thr	Gln	Ile	Ala
15				60					65					70
	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	Glu	Lys	Ile	Gln	Asp
					75					80				
20	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	Gln	Lys	Gly
	85					90					95			
	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met	Glu	Gly
		100					105					110		
25	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr	Asn	Lys
			115					120					125	
	Phe	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln	Asn	Ala
30				130					135					140
	Glu	Ser	Ala	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu	Lys	Thr
					145					150				
35	His	Asp	Lys	Ile	His	Asp	Leu	Ile	Lys	Ala	Gly	Asp	Leu	Asp
	155					160					165			
	Gln	Asp	Ser	Arg	Met	Val	Leu	Val	Asn	Ala	Leu	Tyr	Phe	Lys
		170					175					180		
40	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	Glu	Asn	Thr	Gln	Asp
			185					190					195	
	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn	Val	Arg
45				200					205					210
	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu	Phe	Glu
					215					220				
50	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg	Asn	Ser
	225					230					235			
	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys	Thr	Gly
		240					245					250		
55	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp	Leu	Gln
			255					260					265	
	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile	Leu	Asp
60				270					275					280

-170-

Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu Asn Asp
 285 290
 5 Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Val Pro Gly
 295 300 305
 Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp Glu Met
 310 315 320
 10 Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile Glu Val
 325 330 335
 Asn Glu Glu Gly Ala Glu Ala Ala Ala Thr Ala Thr Phe
 340 345 350
 15 Met Val Thr Tyr Glu Leu Glu Val Ser Leu Asp Leu Pro Thr
 355 360
 20 Val Phe Lys Val Asp His Pro Phe Asn Ile Val Leu Lys Thr
 365 370 375
 Gly Asp Thr Val Ile Phe Asn
 380 385

(2) INFORMATION FOR SEQ ID NO:97:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 354 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: Protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

30 Phe Ala Gly Ser Leu Tyr Asn Thr Val Ala Ser Gly Asn Lys
 1 5 10
 Asp Asn Leu Ile Met Ser Pro Leu Ser Val Gln Thr Val Leu
 15 20 25
 35 Ser Leu Val Ser Met Gly Ala Gly Gly Asn Thr Ala Thr Gln
 30 35 40
 Ile Ala Ala Gly Leu Arg Gln Pro Gln Ser Lys Glu Lys Ile
 45 50 55
 40 Gln Asp Asp Tyr His Ala Leu Met Asn Thr Leu Asn Thr Gln
 60 65 70
 Lys Gly Val Thr Leu Glu Ile Ala Asn Lys Val Tyr Val Met
 75 80
 45 Glu Gly Tyr Thr Leu Lys Pro Thr Phe Lys Glu Val Ala Thr
 85 90 95
 Asn Lys Phe Leu Ala Gly Ala Glu Asn Leu Asn Phe Ala Gln
 100 105 110
 50 Asn Ala Glu Ser Ala Lys Val Ile Asn Thr Trp Val Glu Glu
 115 120 125

-171-

Lys Thr His Asp Lys Ile His Asp Leu Ile Lys Ala Gly Asp
 130 135 140
 5 Leu Asp Gln Asp Ser Arg Met Val Leu Val Asn Ala Leu Tyr
 145 150
 Phe Lys Gly Leu Trp Glu Lys Gln Phe Lys Lys Glu Asn Thr
 155 160 165
 10 Gln Asp Lys Pro Phe Tyr Val Thr Glu Thr Glu Thr Lys Asn
 170 175 180
 Val Arg Met Met His Ile Lys Asp Lys Phe Arg Tyr Gly Glu
 185 190 195
 15 Phe Glu Glu Leu Asp Ala Lys Ala Val Glu Leu Pro Tyr Arg
 200 205 210
 20 Asn Ser Asp Leu Ala Met Leu Ile Ile Leu Pro Asn Ser Lys
 215 220
 Thr Gly Leu Pro Ala Leu Glu Glu Lys Leu Gln Asn Val Asp
 225 230 235
 25 Leu Gln Asn Leu Thr Gln Arg Met Tyr Ser Val Glu Val Ile
 240 245 250
 Leu Asp Leu Pro Lys Phe Lys Ile Glu Ser Glu Ile Asn Leu
 255 260 265
 30 Asn Asp Pro Leu Lys Lys Leu Gly Met Ser Asp Met Phe Val
 270 275 280
 35 Pro Gly Lys Ala Asp Phe Lys Gly Leu Leu Glu Gly Ser Asp
 285 290
 Glu Met Leu Tyr Ile Ser Lys Val Ile Gln Lys Ala Phe Ile
 295 300 305
 40 Glu Val Asn Glu Glu Gly Ala Glu Ala Ala Ala Ala Thr Gly
 310 315 320
 Ile Val Met Leu Gly Cys Cys Met Pro Met Met Asp Leu Ser
 325 330 335
 45 Pro Val Val Phe Asn Ile Asp His Pro Phe Tyr Tyr Ser Leu
 340 345 350
 Met Thr Trp Asp

(2) INFORMATION FOR SEQ ID NO:98:

- 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 356 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: Protein

-172-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

	Phe	Ala	Gly	Ser	Leu	Tyr	Asn	Thr	Val	Ala	Ser	Gly	Asn	Lys
	1				5					10				
5	Asp	Asn	Leu	Ile	Met	Ser	Pro	Leu	Ser	Val	Gln	Thr	Val	Leu
	15				20					25				
	Ser	Leu	Val	Ser	Met	Gly	Ala	Gly	Gly	Asn	Thr	Ala	Thr	Gln
		30				35					40			
10	Ile	Ala	Ala	Gly	Leu	Arg	Gln	Pro	Gln	Ser	Lys	Glu	Lys	Ile
		45					50					55		
	Gln	Asp	Asp	Tyr	His	Ala	Leu	Met	Asn	Thr	Leu	Asn	Thr	Gln
			60					65					70	
15	Lys	Gly	Val	Thr	Leu	Glu	Ile	Ala	Asn	Lys	Val	Tyr	Val	Met
					75					80				
	Glu	Gly	Tyr	Thr	Leu	Lys	Pro	Thr	Phe	Lys	Glu	Val	Ala	Thr
20		85				90					95			
	Asn	Lys	Phe	Leu	Ala	Gly	Ala	Glu	Asn	Leu	Asn	Phe	Ala	Gln
		100				105						110		
25	Asn	Ala	Glu	Ser	Ala	Lys	Val	Ile	Asn	Thr	Trp	Val	Glu	Glu
			115				120						125	
	Lys	Thr	His	Asp	Lys	Ile	His	Asp	Leu	Ile	Lys	Ala	Gly	Asp
			130					135					140	
30	Leu	Asp	Gln	Asp	Ser	Arg	Met	Val	Leu	Val	Asn	Ala	Leu	Tyr
					145					150				
	Phe	Lys	Gly	Leu	Trp	Glu	Lys	Gln	Phe	Lys	Lys	Glu	Asn	Thr
35		155				160					165			
	Gln	Asp	Lys	Pro	Phe	Tyr	Val	Thr	Glu	Thr	Glu	Thr	Lys	Asn
		170					175					180		
40	Val	Arg	Met	Met	His	Ile	Lys	Asp	Lys	Phe	Arg	Tyr	Gly	Glu
			185					190					195	
	Phe	Glu	Glu	Leu	Asp	Ala	Lys	Ala	Val	Glu	Leu	Pro	Tyr	Arg
			200						205				210	
45	Asn	Ser	Asp	Leu	Ala	Met	Leu	Ile	Ile	Leu	Pro	Asn	Ser	Lys
				215						220				
	Thr	Gly	Leu	Pro	Ala	Leu	Glu	Glu	Lys	Leu	Gln	Asn	Val	Asp
50		225				230					235			
	Leu	Gln	Asn	Leu	Thr	Gln	Arg	Met	Tyr	Ser	Val	Glu	Val	Ile
		240				245						250		
55	Leu	Asp	Leu	Pro	Lys	Phe	Lys	Ile	Glu	Ser	Glu	Ile	Asn	Leu
			255				260						265	
	Asn	Asp	Pro	Leu	Lys	Lys	Leu	Gly	Met	Ser	Asp	Met	Phe	Val
60				270					275				280	

-173-

	Pro	Gly	Lys	Ala	Asp	Phe	Lys	Gly	Leu	Leu	Glu	Gly	Ser	Asp
					285					290				
5	Glu	Met	Leu	Tyr	Ile	Ser	Lys	Val	Ile	Gln	Lys	Ala	Phe	Ile
	295					300					305			
	Glu	Val	Asn	Glu	Glu	Gly	Ala	Glu	Ala	Ala	Ala	Ala	Thr	Gly
		310					315						320	
10	Val	Met	Leu	Met	Met	Arg	Cys	Met	Pro	Met	Met	Pro	Met	Ala
			325					330					335	
	Phe	Asn	Ala	Glu	His	Pro	Phe	Leu	Tyr	Phe	Leu	His	Ser	Lys
				340					345					350
15	Asn	Ser	Val	Leu	Phe	Asn								
					355									

While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims.

What is claimed is:

1. An isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene.
2. An isolated nucleic acid molecule selected from the group consisting of: a
5 nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID
10 NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID
15 NO:81, a nucleic acid sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90; and a nucleic acid molecule comprising an allelic variant of a nucleic acid molecule comprising any of said nucleic acid sequences.
3. An isolated protein encoded by a nucleic acid molecule that hybridizes
20 under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene.
4. An isolated flea protein selected from the group consisting of: a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID
25 NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97 and SEQ ID NO:98; and a protein encoded by an allelic variant of a nucleic acid
30 molecule encoding a protein comprising any of said amino acid sequences.

5. A therapeutic composition that, when administered to an animal, reduces hematophagous ectoparasite infestation, said therapeutic composition comprising a protective compound selected from the group consisting of: an isolated flea serine protease inhibitor protein; a mimetope of a flea serine protease inhibitor protein; an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene; an isolated antibody that selectively binds to a flea serine protease inhibitor protein; and an inhibitor of serine protease inhibitor activity identified by its ability to inhibit the activity of a flea serine protease inhibitor protein.
6. An inhibitor of serine protease inhibitor protein activity identified by its ability to inhibit the activity of a flea serine protease inhibitor protein.
7. A mimetope of a flea serine protease inhibitor protein identified by its ability to inhibit flea serine protease activity.
8. A method to reduce hematophagous ectoparasite infestation comprising administering to an animal a therapeutic composition comprising a protective compound selected from the group consisting of: an isolated flea serine protease inhibitor protein; a mimetope of a flea serine protease inhibitor protein; an isolated nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene; an isolated antibody that selectively binds to a flea serine protease inhibitor protein; and an inhibitor of serine protease inhibitor activity identified by its ability to inhibit the activity of a flea serine protease inhibitor protein.
9. A method to produce a flea serine protease inhibitor protein, said method comprising culturing a cell transformed with a nucleic acid molecule that hybridizes under stringent hybridization conditions with a *Ctenocephalides felis* serine protease inhibitor gene.
10. A method to identify a compound capable of inhibiting flea serine protease inhibitor activity, said method comprising:
- (a) contacting an isolated flea serine protease inhibitor protein with a putative inhibitory compound under conditions in which, in the absence of said compound, said protein has serine protease inhibitor activity; and

(b) determining if said putative inhibitory compound inhibits said activity.

11. A test kit to identify a compound capable of inhibiting flea serine protease inhibitor activity, said test kit comprising an isolated flea serine protease inhibitor protein having serine protease inhibitor activity and a means for determining the extent of inhibition of said activity in the presence of a putative inhibitory compound.

12. The nucleic acid molecule of Claim 1, wherein said *Ctenocephalides felis* serine protease inhibitor gene comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90.

13. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule comprises a nucleic acid sequence that encodes a serine protease inhibitor protein.

14. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule is a flea nucleic acid molecule.

15. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule is selected from the group consisting of *Ctenocephalides*, *Ceratophyllus*, *Diamanus*, *Echidnophaga*, *Nosopsyllus*, *Pulex*, *Tunga*, *Oropsylla*, *Orchopeus* and *Xenopsylla* nucleic acid molecules.

16. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule is selected from the group consisting of *Ctenocephalides felis*, *Ctenocephalides canis*, *Ceratophyllus pulicidae*, *Pulex irritans*, *Oropsylla (Thrassis) bacchi*, *Oropsylla*

(*Diamanus*) *montana*, *Orchopeus howardi*, *Xenopsylla cheopis* and *Pulex simulans* nucleic acid molecules.

17. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule comprises a *Ctenocephalides felis* nucleic acid molecule.

5 18. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule hybridizes under stringent hybridization conditions with a nucleic acid molecule selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:45, SEQ ID
10 NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:69 and SEQ ID NO:71.

19. The invention of Claims 1 or 9, wherein said nucleic acid molecule is
15 selected from the group consisting of nfSPI1₁₅₈₄, nfSPI1₁₁₉₁, nfSPI1₃₇₆, nfSPI2₁₃₅₈, nfSPI2₁₁₉₇, nfSPI2₃₇₆, nfSPI3₁₈₃₈, nfSPI3₁₂₆₀, nfSPI3₃₉₁, nfSPI4₁₄₁₄, nfSPI4₁₁₇₉, nfSPI4₃₇₆, nfSPI5₁₄₉₂, nfSPI5₁₁₉₄, nfSPI5₃₇₆, nfSPI6₁₄₅₄, nfSPI6₁₁₉₁, nfSPI6₃₇₆, nfSPI7₅₄₉, nfSPI8₅₄₉, nfSPI9₅₈₁, nfSPI10₆₅₄, nfSPI11₆₇₀, nfSPI12₇₀₆, nfSPI13₆₂₃, nfSPI14₇₃₁, nfSPI15₆₈₅, nfSPI3₁₂₂₂, nfSPI6₁₁₅₅, nfSPI2₁₀₆₅, nfSPI4₁₀₇₀, nfSPIC4:V7₁₁₆₈, nfSPIC4:V8₁₂₂₂,
20 nfSPIC4:V9₁₁₇₄, nfSPIC4:V10₁₁₅₉, nfSPIC4:V12₁₁₇₁, nfSPIC4:V13₁₁₇₁, and nfSPIC4:V15₁₁₇₉.

20. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule is selected from the group consisting of: a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ
25 ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:45, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:50, SEQ ID
30 NO:51, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:65, SEQ ID

NO:66, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:75, SEQ ID NO:78, SEQ ID NO:81, a nucleic acid sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90; and a nucleic acid molecule comprising an allelic variant of any of said
5 nucleic acid molecules.

21. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule encodes a protein comprising an amino acid sequence that is at least about 40% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:8, SEQ ID NO:14, SEQ ID NO:20, SEQ ID NO:26, SEQ ID NO:32, SEQ ID
10 NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90.

22. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule hybridizes under stringent hybridization conditions with a nucleic acid sequence
15 encoding a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID
20 NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90.

23. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule is selected from the group consisting of: a nucleic acid molecule comprising a nucleic acid sequence that encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ
25 ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89 and SEQ ID NO:90; and a nucleic acid molecule comprising an allelic variant of a nucleic acid sequence encoding
30 a protein comprising any of said amino acid sequences.

24. The nucleic acid molecule of Claim 1, wherein said nucleic acid molecule comprises an oligonucleotide.

25. A recombinant molecule comprising a nucleic acid molecule as set forth in Claim 1 operatively linked to a transcription control sequence.

5 26. A recombinant virus comprising a nucleic acid molecule as set forth in Claim 1.

27. A recombinant cell comprising a nucleic acid molecule as set forth in Claim 1.

28. The protein of Claim 3, wherein said nucleic acid molecule hybridizes
10 under stringent hybridization conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:15, SEQ ID NO:21, SEQ ID NO:27, and SEQ ID NO:33, SEQ ID NO:47, SEQ ID NO:50, SEQ ID NO:53, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:62, SEQ ID NO:65, SEQ ID NO:68 and SEQ ID NO:71.

15 29. The protein of Claim 3, wherein said protein, when administered to an animal, elicits an immune response against a serine protease inhibitor protein.

30. The protein of Claim 3, wherein said protein is a flea protein.

31. The protein of Claim 3, wherein said protein is selected from the group consisting of: a protein encoded by a nucleic acid molecule having a nucleic acid
20 sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:19, SEQ IS NO:22, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:45, SEQ ID NO:48, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:57, SEQ ID NO:60, SEQ ID NO:63, SEQ IS NO:66, SEQ ID NO:69, SEQ ID NO:72, SEQ ID NO:75, SEQ ID
25 NO:78 and SEQ ID NO:81; and a protein encoded by a nucleic acid molecule comprising an allelic variant of a nucleic acid molecule comprising any of said nucleic acid sequences.

32. The protein of Claim 3, wherein said protein is selected from the group consisting of: a protein comprising an amino acid sequence selected from the group
30 consisting of SEQ ID NO:2, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:26, SEQ ID

-180-

NO:30, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:49, SEQ ID NO:52, SEQ ID NO:55, SEQ ID NO:58, SEQ ID NO:61, SEQ ID NO:64, SEQ ID NO:67, SEQ ID NO:70, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:95, SEQ ID NO:96, SEQ ID NO:97 and SEQ ID NO:98; and a protein encoded by
5 an allelic variant of a nucleic acid molecule encoding a protein comprising any of said amino acid sequences.

33. An isolated antibody that selectively binds to a protein as set forth in Claims 3 or 4.

34. The invention of Claims 5 or 8, wherein said flea serine protease inhibitor
10 protein comprises a peptide of a flea serine protease inhibitor protein capable of inhibiting serine protease activity.

35. The invention of Claims 5 or 8, wherein said composition further comprises a component selected from the group consisting of an excipient, an adjuvant, and a carrier.

15 36. The invention of Claims 5 or 8, wherein said composition further comprises a compound that reduces hematophagous ectoparasite burden by a method other than by reducing flea serine protease inhibitor activity.

37. The invention of Claims 5 or 8, wherein said protective compound is selected from the group consisting of a naked nucleic acid vaccine, a recombinant virus
20 vaccine and a recombinant cell vaccine.

38. The invention of Claims 5 or 6 or 8, wherein said inhibitor of serine protease inhibitor protein activity comprises a substrate analog of a flea serine protease inhibitor protein.

39. The invention of Claims 6, wherein said inhibitor comprises a
25 peptidomimetic compound.

40. The mimetope of Claim 7, wherein said mimetope comprises a peptidomimetic compound.

41. The method of Claim 8, wherein said hematophagous ectoparasite is a flea.

42. The method of Claim 8, wherein said flea is of a genus selected from the group consisting of *Ctenocephalides*, *Ceratophyllus*, *Diamanus*, *Echidnophaga*, *Nosopsyllus*, *Pulex*, *Tunga*, *Oropsylla*, *Orchopeus* and *Xenopsylla*.

43. The method of Claim 8, wherein said flea is of a species selected from the
5 group consisting of *Ctenocephalides felis*, *Ctenocephalides canis*, *Ceratophyllus pulicidae*, *Pulex irritans*, *Oropsylla (Thrassis) bacchi*, *Oropsylla (Diamanus) montana*, *Orchopeus howardi*, *Xenopsylla cheopis* and *Pulex simulans*.

44. The method of Claim 8, wherein said animal is selected from the group consisting of adult hematophagous ectoparasites, hematophagous ectoparasite larvae and
10 animals susceptible to hematophagous ectoparasite infestation.

45. The method of Claim 8, wherein said animal is selected from the group consisting of adult fleas, flea larvae and animals susceptible to flea infestation.

46. The method of Claim 8, wherein said animal is selected from the group consisting of mammals and birds.

15 47. The method of Claim 8, wherein said animal is selected from the group consisting of felids and canids.

48. The method of Claim 9, wherein said cell is selected from the group consisting of *E.coli*HB:p λ P_R-nfSPI2₁₁₃₉, *E.coli*HB:p λ P_R-nfSPI3₁₁₇₉, *E.coli*HB:p λ P_R-nfSPI4₁₁₄₀, *E.coli*HB:p λ P_R-nfSPI5₁₄₉₂, *E.coli*HB:p λ P_R-nfSPI6₁₁₃₆, *E.coli*:p λ P_R-nfSPIC4:V7₁₁₆₈, *E.coli*:p λ P_R-nfSPIC4:V8₁₂₂₂, *E.coli*:p λ P_R-nfSPIC4:V9₁₁₇₄, *E.coli*:p λ P_R-nfSPIC4:V10₁₁₅₉, *E.coli*:p λ P_R-nfSPIC4:V12₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V13₁₁₇₁, *E.coli*:p λ P_R-nfSPIC4:V15₁₁₇₉, *S. frugiperda*:pVL-nfSPI3₁₂₂₂, *S. frugiperda*:pVL-nfSPI6₁₁₅₅, *S. frugiperda*:pAcG-nfSPI2₁₀₆₅ and *S. frugiperda*:pAcG-nfSPI4₁₀₇₀.

20

1/1




Fig. 1